

Citation for published version:

N. Yousif, J. Cole, J. Rothwell, and J. Diedrichsen,
'Proprioception in motor learning: lessons from a deafferented
subject', *Experimental Brain Research*, Vol. 233 (8): 2449-2459,
August 2015.

DOI:

<https://doi.org/10.1007/s00221-015-4315-8>

Document Version:

This is the Accepted Manuscript version.

The version in the University of Hertfordshire Research Archive
may differ from the final published version. **Users should
always cite the published version.**

Copyright and Reuse:

© Springer-Verlag Berlin Heidelberg 2015.

This manuscript version is distributed under the terms of the
Creative Commons Attribution licence

(<http://creativecommons.org/licenses/by/4.0/>), which permits
unrestricted re-use, distribution, and reproduction in any
medium, provided the original work is properly cited.

Enquiries

If you believe this document infringes copyright, please contact the
Research & Scholarly Communications Team at rsc@herts.ac.uk

Proprioception in motor learning: lessons from a deafferented subject

N Yousif^{1*}, J Cole², J Rothwell³ & J Diedrichsen⁴

1 Division of Brain Sciences, Imperial College London, Charing Cross Campus, London, W6 8RF,
UK

2 University of Bournemouth, Poole Hospital, Longfleet Road, Poole, BH15 2JB, UK

3 Institute of Neurology, University College London, Queen Square, London, WC1N 3BG, UK

4 Institute of Cognitive Neuroscience, University College London, 17 Queen Square, London, WC1N
3AR, UK

* Address for correspondence: Dr Nada Yousif, n.yousif@imperial.ac.uk, tel: +44 (0)20 33117042,
fax: +44 (0)20 33117577

Abstract

1
2
3 Proprioceptive information arises from a variety of channels, including muscle, tendon and skin
4 afferents. It tells us where our static limbs are in space and how they are moving. It remains unclear
5 however, how these proprioceptive modes contribute to motor learning. Here we studied a subject
6 (IW) who has lost large myelinated fibres below the neck and found that he was strongly impaired in
7 sensing the static position of his upper limbs, when passively moved to an unseen location. When
8 making reaching movements however, his ability to discriminate in which direction the trajectory had
9 been diverted was unimpaired. This dissociation allowed us to test the involvement of static and
10 dynamic proprioception in motor learning. We found that IW showed a preserved ability to adapt to
11 force fields when visual feedback was present. He was even sensitive to the exact form of the force
12 perturbation, responding appropriately with a velocity- or position-dependent force after a single
13 perturbation. The ability to adapt to force fields was also preserved when visual feedback about the
14 lateral perturbation of the hand was withdrawn. In this experiment, however, he did not exhibit a form
15 of use-dependent learning, which was evident in the control participants as a drift of the intended
16 direction of the reaching movement in the perturbed direction. This suggests that this form of learning
17 may depend on static position sense at the end of the movement. Our results indicate that dynamic and
18 static proprioception play dissociable roles in motor learning.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43 **Keywords:** Proprioception; motor control; deafferentation; force field learning.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

1
2
3 Proprioception is a collective term that refers to non-visual input that tells us where our body is in
4 space. Proprioception has an important function in normal motor control and motor learning. It is well
5 documented that patients who have suffered loss of proprioception make significant errors in
6 movement (Forget and Lamarre, 1987; Rothwell et al., 1982; Sainburg et al., 1995; Sanes et al.,
7 1985). Proprioception can be further divided into static versus (active or passive) dynamic
8 proprioception (Fuentes and Bastian, 2010; Gritsenko et al., 2007; Jones et al., 2010). Static refers to
9 sensing limb position while stationary, whereas dynamic proprioception involves the estimation of
10 limb position and velocity during either a self-generated active movement or during a passive
11 displacement.
12
13

14 Proprioception relies on various types of mechanoreceptors. Muscle spindle receptors signal both the
15 length of the muscle and the rate of change, hence contributing to both static and dynamic
16 proprioception (Edin and Vallbo, 1990; Matthews, 1933). These receptors are innervated by large,
17 myelinated primary afferent fibres (A α) (Gilman, 2002). Golgi tendon organs respond to muscle
18 tension (contraction) and are principally active during voluntary movements (Houk and Henneman,
19 1967; Prochazka and Wand, 1980). These are subserved by large, myelinated group Ib, A α fibres. In
20 addition, there are also receptors found in the joint capsules. These are predominantly innervated by
21 small afferent fibres signalling nociception, as well as mechanoreceptors signalling flexion or
22 extension. Finally, stretch sensitive receptors in the skin, relayed through large myelinated afferents,
23 also play a role in signalling proprioceptive information (Aimonetti et al., 2007).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 While previous work has shown that these subtypes of proprioception have different properties in
49 motor control (Adamovich et al., 1998; Fuentes and Bastian, 2010; Gritsenko et al., 2007; Ostry et al.,
50 2010; Wong et al., 2012), it is unclear how these different channels contribute to motor learning. To
51 test this we examined the motor behaviour of a deafferented subject (IW), who has lost large
52 myelinated sensory fibres below the neck (Cole and Katifi, 1991; Cole and Sedgwick, 1992; Cole et
53 al., 1995). As expected he was significantly impaired in a static proprioceptive task. However, his
54
55
56
57
58
59
60
61
62
63
64
65

1 ability to detect the direction of a force perturbation to an active, on-going movement was remarkably
2 preserved.

3
4
5 This pattern of deficit and performance allowed us to ask what role active, dynamic and static
6 proprioception plays in error-driven force field adaptation and use-dependent learning. We first tested
7 whether IW would be able to adapt normally to force fields, and specifically if he would be able to
8 shape his adaptive response according to whether a position or velocity-dependent force field was
9 presented (Sing et al., 2009). We also studied him under conditions where visual information about
10 the lateral perturbation was withdrawn to test if force field learning was preserved without vision.
11
12
13
14
15
16
17
18

19 This experiment allowed us to test whether dynamic or static proprioceptive information underlies
20 use-dependent learning. When lateral deviations of the hand from the target cannot be seen and are
21 not penalised as an error, normal participants alter their intended reaching movements in the direction
22 of the perturbation (Diedrichsen et al., 2010). We hypothesized that this form of learning may rely on
23 a static sense of position and therefore predicted that IW would not show the normal response seen in
24 healthy individuals.
25
26
27
28
29
30
31
32
33
34
35
36

37 **Materials and Methods**

38 *Deafferented subject*

39
40 The deafferented subject IW (60 years old at last time of testing, left handed) lost the sense of touch
41 and proprioception as a result of an acute onset sensory neuropathy caused by an autoimmune
42 response to a viral infection when he was 19 years old (Cole and Sedgwick, 1992).
43
44
45
46
47
48

49 Neurophysiological tests have confirmed the loss of large myelinated fibres below the neck (Cole and
50 Katifi, 1991), though he has regained good motor control with cognitive control and with the use of
51 vision and muscle power is normal. He has perception of pain, heat and cold suggesting that small
52 fibres are intact (Cole et al., 1995).
53
54
55
56
57
58
59
60
61
62
63
64
65

Participants

1
2 A total of 16 age-matched control participants were also tested, with two self-reported left-handers
3
4 and the remainder right handed. Nine volunteers participated in the static proprioceptive test (six
5
6 male, mean age 56 ± 10 years) and six different participants took part in the active proprioceptive test
7
8 (three male, mean age 55 ± 12 years) and these six plus an additional participant completed the force
9
10 field adaptation experiments (four male, mean age 54 ± 12 years). Experimental and consent
11
12 procedures were approved by the University College London Ethics Committee.
13
14

Apparatus

15
16
17 Participants sat in front of a virtual-environmental setup, with their head on a forehead rest. They
18
19 made 15cm reaching movements to a single target with their dominant hand while holding onto a
20
21 robotic device. The static proprioceptive task was performed using a robotic device that was equipped
22
23 with an armrest, which brought the elbow into the same plane as the hand. The remainder of the
24
25 experiments were performed on a device without elbow support. Both devices recorded the position of
26
27 the hand with a sampling frequency of 200Hz. A monitor displayed a white square (0.5cm x 0.5cm)
28
29 indicating the start position. The screen was viewed via a mirror, such that participants did not have
30
31 direct vision of their arm, but sometimes received continuous and calibrated visual feedback of their
32
33 hand position either via a 0.3cm diameter white cursor circle or via an arc (see individual experiments
34
35 below).
36
37
38
39
40

Experiment 1: Static proprioception

41
42 To measure the participants' static proprioceptive sense, subjects held onto the robotic manipulandum
43
44 with their dominant hand and were asked to refrain from making any active movements. The robotic
45
46 device then moved the participant's hand to an unknown position, while direct vision of the hand and
47
48 arm was prevented and no visual feedback was displayed on the screen. When the participant's hand
49
50 was within 0.5cm of the target position, a cursor was displayed at a random location on the screen
51
52 (Fig. 1). By moving the cursor with a mouse held in their non-dominant hand, the participant was
53
54 asked to match the cursor location to the location they felt their target hand to be. We tested six
55
56 locations (two at ± 1.3 cm either side of the midline, two at the bottom left of the workspace and two at
57
58
59
60
61
62
63
64
65

1 the top right of the workspace) presented randomly 25 times (total 150 trials divided into six blocks).

2 IW and the one other left-handed participant performed the task with their left hands, and the
3
4 locations of the targets were mirror-reversed in the y-axis. The sequence was constrained such that
5
6 each location was located in a different region of the workspace than the last one. Because the
7
8 sequence was otherwise random, participants needed to rely on a sense of static position at the end of
9
10 the movement. Possible dynamic information of change in hand position was only informative in
11
12 combination with accurate information about the starting location of the hand, which was not
13
14 provided to the participants. The arm was supported such that the elbow and wrist were in the same
15
16 horizontal plane.
17
18

19
20 We analysed the endpoint errors by plotting the 95% confidence ellipse of the responses for each
21
22 target and for each subject. To compare performance quantitatively, we calculate the SD of the
23
24 responses in each direction, and then averaged these across x- and the y-direction. IW's scores were
25
26 expressed as a z-score relative to the mean and standard deviation of the control group's performance.
27
28
29

30 ***Experiment 2: Active proprioception***

31
32 To quantify proprioception to perturbations during active movements, we asked subjects to reach
33
34 straight ahead, while their reach was deviated by the robotic device either to the left or to the right
35
36 (Fig. 1). They had no vision of their hand or arm and no visual feedback about their movement
37
38 direction. On each trial (n=105 in two blocks), we constrained the hand to move an oblique trajectory
39
40 via a force channel and when the participant reached the target we asked them if they had ended up to
41
42 the left or to the right of midline. However, we presented an arced cursor that indicated the distance
43
44 moved by the hand – and an arced target that signalled the required movement amplitude. Both the
45
46 target and the cursor were a 60° segment of a circle, centred on the starting location of the movement.
47
48 Therefore participants only had feedback about how far their arm had reached, and no feedback about
49
50 the endpoint or angle of their reach. We tested 21 angles for the channel that deviated the hand
51
52 movement (0°, ±3°, ±6°, ±9°, ±15°, ±20°, ±25°, ±30°, ±35°, ±40°, ±45°). Each angle was tested five
53
54 times and the presentation of the trials was randomised. We analysed the responses using logistic
55
56 regression. The responses (left/right) were categorised as zeros (left) or ones (right). We first fitted the
57
58
59
60
61
62
63
64
65

1 data using the channel angle and an intercept, which gave two regression weights. The regression
2 weight for the channel angle expresses the steepness of the curve, from which we calculated the inter-
3 quartile range (IQR) as a measure of discrimination ability: the difference in angle for which the
4 participants made 25% rightward or 75% rightward responses. To determine how the force produced
5 in the channel influenced the response of participants, we also added the force measured at 200ms as a
6 third regressor into the logistic model .
7
8
9
10
11
12

13 ***Experiment 3: Force field perturbation***

14 In this experiment, we tested how participants adapted their reaching movements to position and
15 velocity dependent force fields. At the beginning of each trial, participants moved the cursor into the
16 start box. When the target appeared 15cm from the start position, participants were instructed to move
17 the cursor to the target in the straight-ahead direction and stop at the target (Fig. 1). In all experiments,
18 the arm was pushed back to the start without visual feedback - therefore they should have had no
19 knowledge of results from visual information. Movements had to have a peak speed between 55cm/s
20 and 80cm/s. Furthermore the movement needed to stop within 800ms at a distance of less than 0.65cm
21 from the centre of the target. If all these criteria were met, the target turned red and exploded, and
22 participants scored a point. If the movement was not fast enough or was not completed within 800ms,
23 the target turned blue or pink; if participants moved too fast, the target turned yellow; if participants
24 moved at the right speed but stopped the movement too far from the target, the target turned green.
25 However, all trials were included in the analysis, and only trials where participants moved less than
26 half the distance to the target, or showed a large deviation ($>80^\circ$) in the initial movement direction
27 were excluded.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 On force field trials, participants were exposed to a dynamic force field, exerted orthogonally to the
48 actual movement direction. The force [N] could depend either on the position [in cm] or the velocity
49 [cm/s] of the hand.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

$$\begin{bmatrix} Force_x \\ Force_y \end{bmatrix} = C_p * s_1 * \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} * \begin{bmatrix} Position_x \\ Position_y \end{bmatrix} \\ + C_v * s_2 * \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} * \begin{bmatrix} Velocity_x \\ Velocity_y \end{bmatrix}$$

The position-dependent force field ($C_p=\pm 1$, $C_v=0$) pushed the hand either to the left or the right and increased monotonically from the start of the movement. The velocity-dependent force field ($C_p=0$, $C_v=\pm 1$) acted either in a clockwise or anticlockwise direction relative to the movement direction and reached its highest force at peak velocity (Fig. 1). We chose the coefficients s_1 (0.225N/cm) and s_2 (0.075Ns/cm), to achieve a peak force of ~4N on force field trials.

To probe the state of the adaptation response, we used force-channel trials that were inserted throughout the blocks. During channel trials, the robot applied a spring-like force in the x-direction (6000 N/m), which forced the hand onto a 0° trajectory from the starting position. No force was applied on the y-direction. To reduce vibration, we also applied a small viscous damping force (75 Ns/m). If participants expected to be pushed by a force field, they would exert a compensatory force into the wall of the channel during the movement.

Following 80 practice trials (not used for analysis), participants were exposed to eight short blocks of either the position- or velocity-dependent force field in either a clock-wise or anticlockwise direction. These four different conditions were presented in randomised order. Each condition consisted of a 21-trial blocks: one null trial, one channel trial, one force field trial, one channel trial, two force field trials, one channel trial, three force field trials, one channel trial, four force field trials, one channel trial and five null trials (Fig. 1). During all trials subjects had full vision of their hand location via a cursor on the screen.

Experiment 4: Force field adaptation without visual feedback of movement direction

In this experiment, we withdrew all visual feedback about the lateral deviation of the hand. The target was a wide circle segment rather than a square (Fig. 1) and the cursor was not a dot but also a circle segment centred on the starting point of the reach, which expanded as the participant reached towards

1 the target. It therefore displayed the extent of the reach, but gave no information about the lateral
2 position of the hand. Subjects were instructed that they could reach to any angle within the arc target
3 as long as the extent of their reach brought the cursor within the target. Therefore, lateral deviations of
4 the hand were not task relevant (Diedrichsen et al., 2010).
5
6
7

8
9 The experiment consisted of two blocks (one adaptation to a clockwise force field and one
10 anticlockwise, with the order counterbalanced across participants), each 150 trials long, consisting of
11 10 null trials, 30 velocity-dependent force field trials with no channels, 68 force field trials of which
12 27 were randomly presented channel trials and 42 null trials with 15 random channels. The angle of
13 the 27 channel trials in the latter stages of adaptation and in the washout phase was determined by the
14 average endpoint of the participant once they had adapted to the force field. This angle was found
15 from the mean endpoint angle over 15 trials. Each block was split into three parts and participants
16 could have a short break in between parts.
17
18
19
20
21
22
23
24
25
26
27

28 ***Force field analysis***

29 For the force field experiments we were primarily interested in the force with which participants
30 pressed into the force channel. To quantify the strength and time course of the force responses, we
31 regressed the x-force (F_x) of each individual trial from 300ms before movement start until movement
32 end against the y-position (P_y) and the y-velocity (V_y) of the same trial:
33
34
35
36
37
38
39

$$40 \quad F_x - \bar{F} = b_1 s_1 P_y + b_2 s_2 V_y + \varepsilon$$

41
42
43 Before the regression we subtracted a common baseline (\bar{F}) from each trial, which was calculated
44 from the average force profile of the channel trials when the target was at 0° and the channel forced
45 the hand on a 0° trajectory. These trials came from the practice blocks and the pre-adaptation and post
46 washout channel trials in experiments 3 and 4. The multiple regression model therefore did not
47 contain an intercept. The position and velocity traces were scaled by factors s_1 (0.225N/cm) and s_2
48 (0.075Ns/cm) to convert them into units of force. Thus, the two resultant regression coefficients (b_1
49 and b_2) express the adaptation relative to the strength of the position and velocity-dependent force
50 fields. Therefore a regression coefficient of 1 indicates a force that is exactly equal and opposite to the
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 imposed force field, i.e. it implies that the system fully adapted to the force field. Since there was no
2 significant difference between the left and right acting force fields, we sign-reversed the regression
3 coefficients for leftward force fields and then averaged over the left/right trials of the same condition.
4 These regression coefficients are first plotted in (position, velocity) coefficient space. The origin
5 represents no adaptive response, full adaptation to the pure velocity or position-dependent force fields
6 would be evidenced by position/velocity regression coefficients of (0,1) and (1,0).
7
8
9
10
11
12
13
14
15
16

17 **Results**

18 *Static proprioceptive sense is impaired in the deafferented subject.*

19 We first aimed to establish the level of static proprioceptive impairment in IW compared to the
20 controls. Fig. 2A shows the results from a single representative participant performing the task. This
21 participant had a tendency to report the location of their hand to be closer to their body and more to
22 the right, and their mean signed error (Euclidean distance of the reported hand position to the true
23 hand position) between actual hand position and indicated hand position was 2.3cm. IW's responses
24 are shown in Fig. 2B. His responses for the different hand locations were highly overlapping in space,
25 indicating a lack of specificity for the target locations. His error was slightly larger for the central
26 locations (mean = 9.01cm) compared to the targets at the bottom left (mean = 7.55cm) and the top
27 right (mean = 8.34cm), but a one-way ANOVA showed no significant effect of target location
28 (F=1.22, p=0.250). Comparing the spread of responses, IW had mean SD of 5.6cm in the x-direction
29 and 3.6cm in the y-direction. IW's SD lay outside of the range of the normal controls in the x-
30 direction ($z=-6.03$ ($p<0.001$)) but not in the y-direction ($z=-1.2$, $p=0.2297$). This difference may be the
31 result of a range effect (Stevens, 1957) – the observation that participants tend to respond in the
32 middle of a limited scale. In our experiment, the screen used was longer in the x- than in the y-
33 direction, providing a reference frame for the responses. Alternatively, this may relate to the
34 observation in a previous study by van Beers et al (1998). They showed that in a similar
35 proprioceptive static task, localisation was more precise in the radial compared to the azimuthal
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 direction. Averaged over the two directions, the spatial standard deviation for IW lay clearly outside
2 of the range of the control group ($z=-3.21$, $p=0.0013$, Fig. 2C). Interestingly in the static task, we find
3 that the force applied to move the hand to the target was significantly larger for IW (13.3N) than the
4 control group (6.88N, $z=3.16$, $p=0.0016$), indicating that he is likely to stiffen his arm more than the
5 controls. Hence, this proprioceptive task demonstrates that IW has an impaired sense of the static
6 location of his arm is when it is passively moved to an unseen location.
7
8
9
10
11
12

13 *Active proprioception is preserved in deafferented subject.*

14 Next we considered whether dynamic active proprioception was also affected in IW. Based on
15 previous work showing that this patient had lost all large-fibre proprioceptive sense in his upper limbs
16 (Cole and Katifi, 1991; Cole and Sedgwick, 1992; Cole et al., 1995), a similar deficit to that seen in
17 the static task might have been expected. Participants executed point-to-point reaching movements
18 without visual feedback, during which tilted force channels deviated their arm sideways. After each
19 movement, participants reported whether they had been deviated to the left or to the right. Based on
20 the participants' responses we constructed individual psychometric curves using logistic regression
21 (see methods).
22
23
24
25
26
27
28
29
30
31
32
33

34 Fig. 3A (solid line) shows the average psychometric curve for the control participants (after fitting
35 each participant individually) and the same plot for IW (dashed line). IW's curve is steeper than the
36 control's, indicating that he performed better on this task. To quantify this we then computed the
37 angle range which would produce a 25 to 75% change in correct identification, the interquartile range
38 (IQR, Fig. 3B). The mean IQR (based on the individual IQR per control participant) was 48.6°
39 ($\pm 44.4^\circ$) and for IW was 16.4° , which demonstrates his better performance but did not differ
40 significantly ($z=-0.725$, $p=0.468$).
41
42
43
44
45
46
47
48
49
50

51 One possible reason for the poorer performance of control participants may have been a response bias
52 that we observed in the control participants' data, presumably induced by the force that they sensed on
53 their hand in the channel. Of course, the lateral force correlated highly with the direction of the
54 channel – i.e for a rightward pointing channel the participants experienced a rightward pointing force.
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

However, the correlation was not perfect, presumably because the order of the channels was random and participants adapted on a trial-by-trial basis to these tilted channels (Diedrichsen et al., 2010). Hence the correlation between the angle of the channel and experienced force was only $r^2 = 0.848$. When we used the force as an additional variable in the design matrix for the logistic regression, we found that control participants were biased by the force that they exerted over and above the physical angle of the channel. The regression weight for the channel force was significantly less than zero (mean = -0.22, $t(5) = -4.34$ $p = 0.007$), meaning that even in a straight channel, participants were more likely to respond that they were pushed to the left if they produced a leftward force. In contrast, IW did not seem to show the same bias, his channel force regression weight was slightly greater than zero at 0.171, which differed significantly from the controls ($z=3.12$, $p=0.002$). This indicates that the bias in the normal participants arose from a discrepancy between the direction of the force that they exert and the direction that they were pushed in. IW seemed not to be influenced by this bias.

Force field learning with vision is not impaired in the deafferented subject.

The dissociation between static and dynamic active proprioception in IW allowed us to test the importance of these modes of proprioception when adapting to a novel perturbation. Therefore, we let participants adapt to short blocks of either a position-dependent or a velocity-dependent force field. We hypothesised that with visual feedback of the hand, IW may be able to correct for the perturbation, but would be unable to shape the adaptive response appropriately to account for the shape of the applied force field, as he is missing detailed proprioceptive information. Participants had visual feedback of their hand position via a cursor at all times. We measured the amount of adaptation by the force exerted in channel trials. As previously shown (Sing et al., 2009), we found that participants initially responded with a force that was a mixture of position- and velocity-dependent components (Fig. 4). As adaptation progressed, the shape of this response became better matched to the applied perturbation, such that, in the position-dependent force field, more force was applied at the end of the trial, and in the velocity-dependent force field, more force was applied in the middle of the trial. This can be quantified by the path of the regression coefficients of the produced force against velocity and position (Fig. 5, see methods). The path in adaptation space began by a step towards the

1 diagonal and subsequently curved towards the appropriate axis. We found that IW performed
2 similarly to the control subjects. Although his adaptation path in the position-dependent force field
3 was less biased towards the diagonal at the start of adaptation (angle = 6.90° as compared to controls
4 with 25.7°), this difference was not significant ($z=-1.24$, $p=0.216$). Furthermore, the average angle of
5 all his trials (for channel number > 0) was 7.45° , and this was significantly different to zero
6 ($t(38)=2.4$, $p=0.021$). Therefore, IW was able to adapt to force field perturbations, not simply by
7 changing his motor plan to aim to a different location, but by shaping the force he exerts against the
8 force field in a temporally specific manner. Note that IW's baseline performance (during the practice
9 trials without perturbations) was largely matched to the control group in mean peak speed, points
10 gained and movement time, however his absolute endpoint accuracy error was increased compared to
11 controls (1.49cm compared to 0.81cm) and this difference was significant ($z=4.22$, $p<0.01$).
12 Therefore, IW's ability to adapt differentially to two different dynamic conditions clearly
13 demonstrates his preserved ability to deal with dynamic loads, not simply relying on a stereotypical
14 adjustment for perturbations in one direction.

31 *Force field learning without task-relevant visual feedback*

32 We next looked at the case when no visual feedback in the task relevant dimension was given.
33 Subjects reached to an arc target, visual feedback was given by an arc cursor and reaches were
34 rewarded based on the accuracy of the reach extent alone. In a previous study (Diedrichsen et al.,
35 2010) participants exhibited two effects in this experimental paradigm. First, they adapted to the force
36 field as expected from normal error-based learning. Secondly, the intended endpoint also drifted in the
37 direction of the force field. For example, if they reached normally to 0° within the arc, when a
38 clockwise force field was applied their movement would terminate at $+20^\circ$. This was partly due to
39 participants not correcting for the lateral deviation (as it is task irrelevant). A more important factor
40 however, was that the participants also shifted their desired endpoint in the direction of the force field.
41 This was demonstrated by applying force channels that deviated the hand to the left or right of the
42 average end position (Diedrichsen et al., 2010). In the end of the movement, participants would push
43 towards the new end position, indeed confirming that this was the planned final position of the hand.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 We hypothesised that these changes are driven by sensing the location of the movement endpoint
2 using proprioception. Because the movement was generally successful, as the lateral deviation is task
3 irrelevant, participants may adjust their desired endpoint in this direction. With his strong impairment
4 of static proprioception, we predicted that IW would not exhibit these effects.
5
6
7

8
9 Indeed, we found that the endpoints of IW's movements after adaptation to a velocity-
10 dependent force field were much closer to his original endpoint than for control subjects (Fig. 6A).
11 The change from baseline in the angle of the overall movement (start point to endpoint), averaged
12 across clockwise and counter-clockwise force fields, was $17.6^\circ \pm 7.59^\circ$ for controls, but 2.70° for IW,
13 a significant difference, $z=-1.96$, $p=0.050$. Also, IW made large online corrections when perturbed, as
14 can be seen in the sharp hooks in the end of the trajectory. To quantify this observation we calculated
15 the difference between the maximum angle of the movement in the direction of the perturbation and
16 the angle of the endpoint (the angle subtended by the hook at the end of the trajectory). This measure
17 of online correction was 11.4° for IW and $1.79^\circ (\pm 1.95^\circ)$ for the control, again a significant
18 difference, $z=4.89$, $p<0.001$. Thus, IW showed a reduced drift of his endpoint in the direction of the
19 force field, and exhibited increased online corrections compared to controls.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 However, Figure 6a also reveals a strong asymmetry of the effect with respect to the direction
36 of the force field. While the change in IW's anti-clockwise endpoint (5.16°) was significantly
37 different to the controls ($-22.0^\circ \pm 11.1^\circ$, $z=2.45$ $p=0.01$), the change in his clockwise endpoint (10.6°)
38 did not differ significantly ($13.2^\circ \pm 6.37^\circ$). Similarly, the increase in online correction was mostly
39 apparent in the anti-clockwise force field ($z=-6.36$, $p<0.01$). While they were also slightly higher than
40 control values in the clockwise direction (5.76° vs. $0.59^\circ \pm 3.55^\circ$), this difference was not significant
41 ($z=1.46$, $p=0.15$). One possible reason for this asymmetry is that IW used his left hand to perform the
42 task, while nearly all the control participants used their right hand. Following this idea, both groups
43 would show an increased drift towards flexion of the shoulder joint, possibly explained by stronger
44 flexor than extensor muscles. While our one left-handed control participant did not show this effect,
45 more left-handed participants would need to be tested to test this explanation further.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 To test for the adaptation to the force fields themselves, we applied channel trials tilted to the
3 participant's measured overall movement direction. The forces exerted during these channel trials
4 (Fig. 6B), demonstrate that control subjects exhibited a force profile with a typical velocity-dependent
5 profile as expected. IW's data showed a very similar pattern to the controls. Quantifying the force
6 profiles as a mixture of position- and velocity-dependent force as above, we found that the velocity
7 regression weights for the controls are significantly greater than zero (mean = 0.0564, $t(6) = 12.8$, $p <$
8 0.001) and for IW this regression weight was 0.0397, which based on a Z-score of -1.43 was not
9 significantly different from the controls ($p=0.152$). Similarly, the controls had a non-significant
10 position regression weight (mean = 0.008, $t(6) = 0.805$, $p = 0.452$), and IW's position regression
11 weight (0.0485) was not significantly different to the controls ($Z=1.54$, $p = 0.123$). Therefore, both the
12 control group and IW adapted to the velocity force field without task-relevant feedback.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 **Discussion**

29
30
31 The study of deafferented subjects has contributed greatly to our insight about the role of
32 proprioception in motor control and learning. However, such studies have relied on a small number of
33 subjects with differing degrees of motor control and differing levels of deficit. Our study demonstrates
34 that rather than relying on the fact that proprioception is generally impaired in these individuals, it is
35 important to quantify the impairments of such subjects carefully for different tasks.
36
37
38
39
40
41
42

43 We have shown that the deafferented subject IW has lost most of his proprioceptive static position
44 sense, as when the arm is moved passively to a new location, he was significantly impaired compared
45 to the controls in localizing his arm. This is in line with previous work which has shown that some
46 deafferented subjects (GL) have inaccurate endpoints during reaching (Forget and Lamarre, 1987) as
47 well as an inability to sustain constant muscle contraction without vision (Rothwell et al., 1982),
48 although IW has previously been shown to maintain a small force without feedback (Cole and
49 Sedgwick, 1992). The differences may reflect IW's long period of rehabilitation compared with the
50 subject studied by Rothwell et al., and highlights the importance of differences between subjects with
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

apparently similar conditions. While it has previously been shown that IW is able to actively perform simple repetitive movements without vision when position was not crucial (Cole and Sedgwick, 1992), our task showed that IW was inaccurate at detecting the static position of his hand, i.e. when it had stopped after a passive movement to an unseen location.

The time elapsed after reaching could play a role in the accuracy of the localisation, since the static proprioceptive sense drifts with the time that a limb has been stationary (Paillard, 1968). However, IW did not spend longer than our controls making his decision ($Z=-0.1708$, $p=0.8644$) and therefore this is unlikely to explain IW's impairment. We did not give subjects any feedback about hand position, once the static test had started, and therefore subjects could have accumulated errors on hand position and become increasingly worse. However, we did not find that error increased with increasing number of trials and so this cannot account for differences in performance with this factor.

In contrast, in an active and dynamic proprioceptive test, where subjects made constrained reaches and had to discriminate perturbations to the right from the left, IW performed better than control participants. In controls, it has been suggested that the activation of muscles improves proprioception due to increased discharge from spindles (and Golgi tendon organs) during voluntary contraction (Gandevia et al., 1992; Laufer et al., 2001). It was recently shown that healthy controls are better at discriminating arm position when actively moving compared to when their arm is passively moved (Bhanpuri et al., 2013). Conversely, (Capaday et al., 2013) showed that when performing an active pointing movement subjects were no more accurate at localising their finger position than if their arm was passively moved. Similarly, our control participants did not perform better in the active compared to the static task and seemed to exhibit a form of force illusion, which impaired their performance during the active task. Subjects confused the direction in which their hand was pushed with the direction in which they pushed into the force channel, while IW did not show this effect.

The origin of IW's perception of perturbations under dynamic conditions is unclear. His perception of touch in the hand is poor, his von Frey monofilament threshold has previously been shown to be 200mN compared with 6mN for controls (Cole et al., 2006). Though we did not measure joint

1 angulation during this task, most movement seemed to occur at the shoulder and elbow, with differing
2 amounts of movement of each for different positions, especially for the static task. It is possible that
3
4 IW has some remaining afferents from these joints and their muscles, undetectable from conventional
5
6 testing of joint position sense, either by perception or matching between arms. However, the
7
8 perturbations were more abrupt during the active task and it is difficult to exclude afferent information
9
10 being picked up through the neck, head, or possibly the inner ear.
11
12

13
14 One additional limitation is that the active and static perceptual tests of proprioception are different in
15
16 difficulty and response (categorical versus continuous) and therefore it is difficult to compare these
17
18 tasks directly. Finally, although during the static task, the arm was moved slowly to a new random
19
20 position, we cannot rule out the possibility that subjects used dynamic information from the passive
21
22 movement. Future work could employ the use of vibration during the passive movement to mask any
23
24 proprioceptive information.
25
26

27
28 The observed dissociation allowed us to test the involvement of static and dynamic proprioception in
29
30 force field learning. It has previously been shown that deafferented subjects can adapt as well as
31
32 healthy controls to visual rotations (Bernier et al., 2006). More recently, it has been shown that
33
34 deafferented patients can adapt to force fields with visual feedback (Sarlegna et al., 2010) and healthy
35
36 controls can force field adapt when proprioceptive and visual information are conflicting (Melendez-
37
38 Calderon et al., 2011). Our results mirror these findings, such that IW was able to adapt to the force
39
40 field when full vision was present (experiment 3) and when there was no task relevant visual feedback
41
42 (experiment 4). This use of visual information of the moving body part has been previously described
43
44 as “visual proprioception” and our results suggest that this remains preserved in IW. Furthermore,
45
46 when analysing the forces produced in the channel trials, not only was the force field counteracted,
47
48 but the temporal shape of the force field (velocity- or position-dependent) was also reproduced. This
49
50 indicates that IW not only learnt how to reach the target in the presence of a force field, but did so in a
51
52 way specific to the force field encountered. Given this result, we propose that the error-based learning
53
54 required for force field adaptation uses active and dynamic rather than static proprioceptive
55
56 information to improve performance.
57
58
59
60
61
62

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Another learning modality which has recently been demonstrated for both force field learning (Diedrichsen et al., 2010), spatial errors (Verstynen and Sabes, 2011) and movement speed (Hammerbeck et al., 2014), has been referred to as use-dependent learning. This learning mechanism refers to the phenomenon that the next movement plan is updated to become more similar to the last experienced movement, if this movement was reasonably successful. When exposed to a force field during a task when the lateral deviation did not affect success, control subjects showed use-dependent learning in their endpoints, such that these participants shifted their desired endpoint in the direction of the perturbation. IW did not demonstrate this shift, and maintained a central desired endpoint, as evidenced by the strong corrections at the end of the movement. One possibility is that IW pre-planned his movements in terms of a desired endpoint and increased his arm stiffness to achieve this position using equilibrium-point control (Polit and Bizzi, 1978). The finding that IW produced adequate forces to resist a velocity-dependent force field in a force channel, however, argues against pure equilibrium-point control.

In the experiment, however, IW showed a little to no drift and large endpoint correction in the anticlockwise force field, but with nearly normal looking behaviour in the clockwise force field. While the reasons for this marked asymmetry are not fully clear, the explanation may lay in the fact that IW, who is left-handed, also performed the task with the left hand. An anticlockwise force field would have mostly loaded flexor muscles, which are stronger than extensor muscles, which would have allowed him better to resist the force field.

Our results suggest that the shift of the desired endpoint shown by controls may be caused by the proprioceptively sensed endpoint position of the hand becoming the desired or planned endpoint for the next movement. An alternative possibility is that the control group did not correct at the end of the movement, rather than shifting their endpoint. To test for this possibility, we put in channel trials at $\pm 8^\circ$ around the calculated preferred endpoint angle once subjects were force field adapted. We found that subjects would correct towards their new endpoint if pushed away from it by a force channel (Diedrichsen et al., 2010). Therefore, this indicates that our control subjects actively shifted their endpoint rather than failed to correct.

1 In conclusion, we found that a chronically deafferented subject shows significantly worse static
2 proprioception, and although not-significant, slightly better mean dynamic proprioception than
3 healthy, aged match controls. IW was also able to adapt to force fields during a forward reach of the
4 arm as well as healthy controls, but did not show a type of use-dependent learning. The questions of
5 which receptors in which body parts underlay these results and the role of vision and motor learning
6 remains unclear. These results suggest the adoption of caution when ascribing results from force field
7 experiments to peripheral proprioception and to specific afferents or joints.
8
9
10
11
12
13
14
15
16
17
18
19
20
21

22 **Acknowledgements**

23 NY is funded by a medical research grant from the Bupa Foundation.
24
25
26
27
28
29
30
31
32
33
34

35 **Conflicts of Interest**

36 The authors report no conflict of interest.
37
38
39
40
41
42
43
44
45
46

47 **References**

- 48 Adamovich SV, Berkinblit MB, Fookson O, Poizner H (1998) Pointing in 3D space to remembered
49 targets. I. Kinesthetic versus visual target presentation. J Neurophysiol 79: 2833-46
50
51 Aimonetti JM, Hospod V, Roll JP, Ribot-Ciscar E (2007) Cutaneous afferents provide a neuronal
52 population vector that encodes the orientation of human ankle movements. J Physiol 580:
53 649-58
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Bernier PM, Chua R, Bard C, Franks IM (2006) Updating of an internal model without
2 proprioception: a deafferentation study. *Neuroreport* 17: 1421-5
3
- 4 Bhanpuri NH, Okamura AM, Bastian AJ (2013) Predictive modeling by the cerebellum improves
5 proprioception. *J Neurosci* 33: 14301-6
6
7
- 8 Capaday C, Darling WG, Stanek K, Van Vreeswijk C (2013) Pointing to oneself: active versus
9 passive proprioception revisited and implications for internal models of motor system
10 function. *Exp Brain Res* 229: 171-80
11
12
13
- 14 Cole J, Bushnell MC, McGlone F, Elam M, Lamarre Y, Vallbo A, Olausson H (2006) Unmyelinated
15 tactile afferents underpin detection of low-force monofilaments. *Muscle Nerve* 34: 105-7
16
17
- 18 Cole JD, Katifi HA (1991) Evoked potentials in a man with a complete large myelinated fibre sensory
19 neuropathy below the neck. *Electroencephalogr Clin Neurophysiol* 80: 103-7
20
21
22
- 23 Cole JD, Sedgwick EM (1992) The perceptions of force and of movement in a man without large
24 myelinated sensory afferents below the neck. *J Physiol* 449: 503-15
25
26
27
- 28 Cole JD, Merton WL, Barrett G, Katifi HA, Treede RD (1995) Evoked potentials in a subject with a
29 large-fibre sensory neuropathy below the neck. *Can J Physiol Pharmacol* 73: 234-45
30
31
32
- 33 Diedrichsen J, White O, Newman D, Lally N (2010) Use-dependent and error-based learning of motor
34 behaviors. *J Neurosci* 30: 5159-66
35
36
37
- 38 Edin BB, Vallbo AB (1990) Dynamic response of human muscle spindle afferents to stretch. *J*
39 *Neurophysiol* 63: 1297-306
40
41
- 42 Forget R, Lamarre Y (1987) Rapid elbow flexion in the absence of proprioceptive and cutaneous
43 feedback. *Hum Neurobiol* 6: 27-37
44
45
- 46 Fuentes CT, Bastian AJ (2010) Where is your arm? Variations in proprioception across space and
47 tasks. *J Neurophysiol* 103: 164-71
48
49
50
- 51 Gandevia SC, McCloskey DI, Burke D (1992) Kinaesthetic signals and muscle contraction. *Trends*
52 *Neurosci* 15: 62-5
53
54
- 55 Gilman S (2002) Joint position sense and vibration sense: anatomical organisation and assessment. *J*
56 *Neurol Neurosurg Psychiatry* 73: 473-7
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Gritsenko V, Krouchev NI, Kalaska JF (2007) Afferent input, efference copy, signal noise, and biases in perception of joint angle during active versus passive elbow movements. *J Neurophysiol* 98: 1140-54
- Hammerbeck U, Yousif N, Greenwood R, Rothwell JC, Diedrichsen J (2014) Movement speed is biased by prior experience. *J Neurophysiol* 111: 128-34
- Houk J, Henneman E (1967) Responses of Golgi tendon organs to active contractions of the soleus muscle of the cat. *J Neurophysiol* 30: 466-81
- Jones SA, Cressman EK, Henriques DY (2010) Proprioceptive localization of the left and right hands. *Exp Brain Res* 204: 373-83
- Laufer Y, Hocherman S, Dickstein R (2001) Accuracy of reproducing hand position when using active compared with passive movement. *Physiother Res Int* 6: 65-75
- Matthews BH (1933) Nerve endings in mammalian muscle. *J Physiol* 78: 1-53
- Melendez-Calderon A, Masia L, Gassert R, Sandini G, Burdet E (2011) Force field adaptation can be learned using vision in the absence of proprioceptive error. *IEEE Trans Neural Syst Rehabil Eng* 19: 298-306
- Ostry DJ, Darainy M, Mattar AA, Wong J, Gribble PL (2010) Somatosensory plasticity and motor learning. *J Neurosci* 30: 5384-93
- Paillard JB, M., 1968. Active and passive movements in the calibration of position sense. In: *The Neuropsychology of Spatially Oriented Behaviour*. Vol., S.J. Freeman, ed. eds. Dorsey Press, Homewood, IL, pp. 37-55.
- Polit A, Bizzi E (1978) Processes controlling arm movements in monkeys. *Science* 201: 1235-7
- Prochazka A, Wand P (1980) Tendon organ discharge during voluntary movements in cats. *J Physiol* 303: 385-90
- Rothwell JC, Traub MM, Day BL, Obeso JA, Thomas PK, Marsden CD (1982) Manual motor performance in a deafferented man. *Brain* 105 (Pt 3): 515-42
- Sainburg RL, Ghilardi MF, Poizner H, Ghez C (1995) Control of limb dynamics in normal subjects and patients without proprioception. *J Neurophysiol* 73: 820-35

1 Sanes JN, Mauritz KH, Dalakas MC, Evarts EV (1985) Motor control in humans with large-fiber
2 sensory neuropathy. *Hum Neurobiol* 4: 101-14
3
4 Sarlegna FR, Malfait N, Bringoux L, Bourdin C, Vercher JL (2010) Force-field adaptation without
5 proprioception: can vision be used to model limb dynamics? *Neuropsychologia* 48: 60-7
6
7 Sing GC, Joiner WM, Nanayakkara T, Brayanov JB, Smith MA (2009) Primitives for motor
8 adaptation reflect correlated neural tuning to position and velocity. *Neuron* 64: 575-89
9
10 Stevens SS (1957) On the psychophysical law. *Psychol Rev* 64: 153-81
11
12 Verstynen T, Sabes PN (2011) How each movement changes the next: an experimental and theoretical
13 study of fast adaptive priors in reaching. *J Neurosci* 31: 10050-9
14
15 Wong JD, Kistemaker DA, Chin A, Gribble PL (2012) Can proprioceptive training improve motor
16 learning? *J Neurophysiol* 108: 3313-21
17
18
19
20
21
22
23
24
25
26
27
28

29 **Figure Captions**

30
31
32 **Fig. 1:** Experimental paradigm. Experiment 1: In the static proprioception test participants' right hand
33 was passively moved to one of six randomly chosen locations. There was no visual feedback about the
34 position of the hand (dashed line). At the end of the movement, the participant indicated the felt
35 position of the hand with the left hand via a cursor controlled by a mouse. Experiment 2: In the active
36 proprioceptive task we asked subjects to reach to an arc target without vision of their hand or arm, and
37 with cursor feedback provided via an arc cursor. At the end of each reach we asked subjects to
38 respond as to whether they had been pushed to the right or left. Experiment 3: Participants were
39 exposed to position- or velocity-dependent force fields for short blocks of 21 trials with full visual
40 feedback (solid line for hand). Experiment 4: Participants adapted to a velocity-dependent force field
41 in long blocks of 150 trials with task irrelevant visual feedback.
42
43
44
45
46
47
48
49
50
51
52
53

54
55 **Fig. 2:** Results of static proprioception test. **A)** The responses (small dots) and 95% confidence
56 ellipses of a single control participant performing the static task for the six targets. **B)** IW's responses
57 were highly overlapping in space, indicating the lack of specificity for the target locations. **C)** Box
58
59
60
61
62
63
64
65

1 plot showing the average standard deviation of the endpoints for the control participants and IW
2 (black circle).
3
4

5 **Fig. 3:** Results of dynamic proprioception task. **A)** The mean psychometric curve obtained from the
6 logistic regression of control participants' responses in the active proprioceptive task (solid line) and
7 the same plot for IW's responses (dashed line). **B)** The performance in the active task was quantified
8 by the IQR, showing that the control group (black bar) had a larger IQR indicating worse performance
9 compared to IW (white bar).
10
11
12
13
14
15
16

17 **Fig. 4:** Results of Experiment 3. During adaptation to clockwise (red) and anti-clockwise (blue) force
18 fields, the amount of adaptation was quantified by the force exerted in the channel trials as shown
19 here for controls (solid) and IW (dashed). Participants initially respond with a force which was made
20 up of a mixture of position and velocity components in both types of force field (1st channel), but this
21 force becomes shaped to match the applied perturbation as participants adapt.
22
23
24
25
26
27
28

29 **Fig. 5:** Force adaptation in Experiment 3, plotted as position and velocity weights for the position
30 dependent force field (grey) and the velocity dependent force field (black) blocks. Adaptation curves
31 were biased towards the diagonal and then curved towards the appropriate axis. IW (dashed lines)
32 performs similarly to the control subjects (solid lines) and is able to shape his responses appropriately
33 to the applied force field.
34
35
36
37
38
39
40

41 **Fig. 6:** Results of experiment 4. **(A)** Average trajectory for a clock-wise (blue) or anticlockwise (red)
42 perturbing force field. The mean trajectory across control subjects (solid lines) demonstrates that they
43 shifted their endpoint in the direction of the applied force field. In contrast, IW (dashed lines)
44 corrected more with the endpoints remaining closer to the midline. **B)** The channel forces (shading
45 indicates between subject standard error) once subjects had adapted to the force field show that the
46 control subjects (solid line) show a velocity dependent force profile as expected. IW shows a very
47 similar profile to the controls (dashed line).
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig 1

[Click here to download Figure: Figure1.tif](#)




Fig 2

[Click here to download Figure: Figure2.tif](#)




Fig 3

[Click here to download Figure: Figure3.tif](#)





Fig 4

[Click here to download Figure: Figure4.tif](#)

Position-dependent force field



Velocity-dependent force field




Fig 5


[Click here to download Figure: Figure5.tif](#)

Fig 6

[Click here to download Figure: Fig6.tif](#)