



Article

Maximal Fat Oxidation during Incremental Upper and Lower Body Exercise in Healthy Young Males

Mike Price ^{1,*}, Lindsay Bottoms ² , Matthew Hill ¹  and Roger Eston ³

¹ Centre for Sport, Exercise and Life Sciences, School of Life Sciences, Coventry University, Coventry CV1 5FB, UK

² Department of Psychology, Sport and Geography, University of Hertfordshire, Hatfield AL10 9AB, UK

³ Alliance for Research in Exercise, Nutrition and Activity, Campus Central—City East, University of South Australia, GPO Box 2471, Adelaide 5001, Australia

* Correspondence: mike.price@coventry.ac.uk

Abstract: The aim of this study is to determine the magnitude of maximal fat oxidation (MFO) during incremental upper and lower body exercise. Thirteen non-specifically trained male participants (19.3 ± 0.5 y, 78.1 ± 9.1 kg body mass) volunteered for this repeated-measures study, which had received university ethics committee approval. Participants undertook two incremental arm crank (ACE) and cycle ergometry (CE) exercise tests to volitional exhaustion. The first test for each mode served as habituation. The second test was an individualised protocol, beginning at 40% of the peak power output (PO_{peak}) achieved in the first test, with increases of 10% PO_{peak} until volitional exhaustion. Expired gases were recorded at the end of each incremental stage, from which fat and carbohydrate oxidation rates were calculated. MFO was taken as the greatest fat oxidation value during incremental exercise and expressed relative to peak oxygen uptake ($\dot{V}O_{2peak}$). MFO was lower during ACE (0.44 ± 0.24 g·min⁻¹) than CE (0.77 ± 0.31 g·min⁻¹; respectively, $p < 0.01$) and occurred at a lower exercise intensity (53 ± 21 vs. $67 \pm 18\%$ $\dot{V}O_{2peak}$; respectively, $p < 0.01$). Inter-participant variability for MFO was greatest during ACE. These results suggest that weight loss programs involving the upper body should occur at lower exercise intensities than for the lower body.

Keywords: Fat_{max}; carbohydrate oxidation; cycle ergometry; arm crank ergometry; variability



Citation: Price, M.; Bottoms, L.; Hill, M.; Eston, R. Maximal Fat Oxidation during Incremental Upper and Lower Body Exercise in Healthy Young Males. *Int. J. Environ. Res. Public Health* **2022**, *19*, 15311. <https://doi.org/10.3390/ijerph192215311>

Academic Editor: Paul B. Tchounwou

Received: 7 October 2022

Accepted: 14 November 2022

Published: 19 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

During the initial intensities of incremental exercise, both carbohydrate and fat oxidation increase [1]. However, in subsequent stages of submaximal exercise, carbohydrate oxidation continues to increase, whereas fat oxidation reaches maximal values and begins to decrease [1,2]. Within such an incremental exercise model, the greatest fat oxidation value observed is termed maximal fat oxidation (MFO). The exercise intensity at which the MFO occurs, typically a percentage of maximal or peak oxygen uptake ($\dot{V}O_{2peak}$), has been termed 'Fat_{max}' [3]. The intensity at which Fat_{max} occurs has been shown to be affected by factors such as endurance training status [4], prior carbohydrate ingestion [5], habitual physical activity and maximal oxygen uptake ($\dot{V}O_{2max}$) [6]. Furthermore, few studies have directly compared males and females, with MFO tending to be greater in males but Fat_{max} tending to be greater in females [7]. Although the determination of the exercise intensity at which Fat_{max} occurs has been reported to be reliable, there is often large inter- and intra-individual variation, as demonstrated by large coefficients of variation and not all participants demonstrating a clear Fat_{max} intensity [6,8].

Previous studies examining Fat_{max} have predominantly considered lower body exercise. However, for the general population, upper body exercise elicits lower $\dot{V}O_{2peak}$ than for lower body exercise modes [9–12]. For example, values for $\dot{V}O_{2peak}$ during

arm crank ergometry (ACE) and cycle ergometry (CE) in non-specifically trained individuals have been reported as 34 and 48 mL·kg⁻¹·min⁻¹, respectively [10], with upper body values generally representing ~70% of those during CE [10,11,13]. When $\dot{V}O_{2peak}$ during ACE (e.g., 28–39 mL·kg⁻¹·min⁻¹) is compared to that during treadmill running (e.g., 44–59 mL·kg⁻¹·min⁻¹) in similar non-specifically trained populations [9,12], the proportional difference between modes is increased, with ACE being ~60–65% of treadmill values [9,12]. Differences in $\dot{V}O_{2peak}$ across exercise modes are predominantly due to differences in the size of the active muscle mass. The physiological range for metabolic values during upper body exercise is thus lower than for other exercise modes and likely includes a reduced range of fat metabolism.

Maximal fat oxidation and Fat_{max} during typical incremental CE and treadmill walking/running protocols have been directly compared in moderately trained men [14]. Maximal oxygen uptake and fat oxidation were lower during CE (64 mL·kg⁻¹·min⁻¹ and 0.47 g·min⁻¹, respectively) than for treadmill exercise (67 mL·kg⁻¹·min⁻¹ and 0.65 g·min⁻¹, respectively), as was fat oxidation over a range of submaximal exercise intensities. However, Fat_{max} was similar between modes (~60% $\dot{V}O_{2peak}$), as was the point where blood lactate increased from rest (i.e., the lactate threshold), suggesting no exercise mode differences for these specific variables. The participants in the above study were club/county standard endurance cyclists and, therefore, of greater aerobic training status than the general population, where fitness indicators such as lactate threshold are lower in ACE when compared to CE and treadmill exercise [12]. Differences in MFO and Fat_{max} across exercise modes may, therefore, be more pronounced for non-specifically trained individuals.

As well as eliciting lower aerobic fitness indicators when compared to lower body exercise modes, the upper body is also considered to possess a greater proportion of fast twitch muscle fibres [15,16]. Considering that fast twitch fibres are more glycolytic than slow twitch fibres and that lower training status elicits lower absolute MFO values at lower relative exercise intensities [4,6], the magnitude of both MFO and Fat_{max} is likely lower for upper body exercise. Studies that have investigated Fat_{max} during upper body exercise have examined individuals with diabetes and obesity, specifically focusing on mitochondrial function [17,18], or those with spinal cord injury, focusing upon substrate utilisation [19]. Within these studies, healthy or non-disabled controls have been utilised and matched for age and body mass [17,18].

Based on the above studies, our knowledge of MFO and Fat_{max} during upper body exercise is from a small number of studies of older (~30–40 years of age) and heavier participants (>90 kg) with low aerobic fitness; all factors associated with poor MFO and Fat_{max} intensities [20]. Therefore, the aim of this study is to determine the magnitude of MFO and Fat_{max} during upper body and lower body exercise in young, healthy, recreationally active males. Due to the likely differences in muscle mass, metabolic capacity and training status between the upper and lower body, it is hypothesised that MFO and Fat_{max} would be lower and more variable during upper body exercise.

2. Materials and Methods

2.1. Participants

Thirteen healthy, active males volunteered to participate in this study (age; 19.3 ± 0.5 y, body mass; 78.1 ± 9.1 kg; 4.5 ± 1.8 sessions per week in predominantly team sports), which had received university ethics committee approval (Ref: P35579) and was in accordance with the Declaration of Helsinki. A sub-group of participants (n = 9; age 19.2 ± 0.4 y; body mass 76.8 ± 8.9 kg) undertook an additional exercise testing session, as outlined below. All participants provided written informed consent. An a priori sample size calculation was undertaken using a desired power value of 0.8, an alpha level of $p = 0.05$ and effect size values based on the MFO and Fat_{max} data from Ara et al. [17] and Larsen et al. [18], who previously compared these variables during upper and lower body exercise. The calculation resulted in a recommended sample of n = 6. Therefore, our sample of n = 13

was sufficient to account for any potentially large variation in MFO and Fat_{max} responses and any participant drop-out, as well as being in line with previous studies ($n = 10$ and $n = 7$, respectively).

The study was a repeated measure design. Inclusion criteria were: young, healthy males who took part in structured exercise sessions at least three times each week. Exclusion criteria were: being specifically cycle or arm ergometry trained or endurance athletes. Recruitment occurred through existing researcher networks and participant friendship groups. Participants visited the university's exercise physiology laboratory on four separate occasions. Testing sessions were undertaken in the morning, at the same time of day, with at least three days between sessions. Participants were requested not to consume any food within the two to three hours preceding each testing session and to keep the same diet on all testing days, which was confirmed verbally. Participants were also asked not to undertake any strenuous exercise or consume any alcohol or caffeinated products in the 24 h prior to testing. Participants were able to drink water *ad libitum* during each testing session. Laboratory conditions were maintained at approximately 19–20 °C and ~40% relative humidity. All participants wore similar attire on each laboratory visit, being: light-weight tracksuit trousers, shorts, sports socks and training shoes. Data were collected by the lead author, with considerable experience in both upper and lower body exercise testing.

2.2. Exercise Tests

The testing sessions involved two ACE tests, two incremental CE tests and, for the sub-group of participants, one submaximal ACE test. The incremental exercise tests were maximal aerobic tests to determine $\dot{V}O_{2peak}$ and power output (PO_{peak}). Lower body exercise tests were undertaken using a cycle ergometer (Monark 814E, Varberg, Sweden), whereas the upper body exercise tests were undertaken on an electronically braked Lode arm crank ergometer (Lode Angio, Groningen, the Netherlands). The first ACE and CE tests were preliminary tests to habituate the participants to each exercise mode and the testing environment. For ACE, the protocol began at 0 W, with increments of 20 W every 2-min [21], whereas the CE protocol began at 35 W and increased by 35 W every 3 min [10]. Power output was progressively increased until volitional exhaustion occurred or when the desired cadence (i.e., 70 $rev \cdot min^{-1}$) could not be maintained for 10 s. The second ACE and CE protocols were individualised protocols based on each participant's performance in the preliminary tests. Participants initially warmed up for 5 min on the unloaded ergometer (0 and 35 W for ACE and CE, respectively) before beginning the main protocol at 40% of the PO_{peak} achieved in the preliminary tests. After this stage, the exercise intensity increased by 10% of each participant's initial PO_{peak} every two minutes for ACE and every three minutes for CE until volitional exhaustion, using the same criteria as in the preliminary tests.

As previous studies of upper body exercise routinely used exercise protocols with stage durations of 2 min [10,21,22], it is possible that metabolic responses may not have adapted to the same extent as observed for CE when using 3 min exercise stage durations. Therefore, a sub-group of participants (detailed above) undertook a submaximal ACE protocol comparing 2- and 3-minute exercise responses. The submaximal protocol involved the participants undertaking the same individual exercise intensities for three minutes at each stage, up to a rating of perceived exertion (RPE, Borg Scale; [23]) for local muscular effort (RPE_L) of 15 (equivalent to 'Hard' exercise on the 6–20 Borg scale). After all exercise tests, participants performed a cool-down on the unloaded ergometer (0 and 35 W for ACE and CE, respectively) at a self-selected cadence, typically 30 to 50 $rev \cdot min^{-1}$ for ACE and 50 to 60 $rev \cdot min^{-1}$ for CE, for at least five minutes to aid recovery. Due to individualised exercise protocols being undertaken, only the order of ACE and CE testing modes could be realistically counterbalanced.

2.3. Measurements

Heart rate (HR) was recorded using a Polar heart rate monitor (Polar beat, Kempele, Finland) consisting of a telemetry belt worn around the chest and a monitor. Values

were recorded during seated rest and in the last 15 s of each exercise stage. Participants also wore a face mask connected to an online expired gas analysis system (Metamax 3b, Biophysik GmbH Leipzig, Germany) for the determination of oxygen uptake ($\dot{V}O_2$) and carbon dioxide production ($\dot{V}CO_2$), minute ventilation ($\dot{V}E$) and the respiratory exchange ratio (RER). Carbohydrate and fat oxidation were subsequently calculated according to standard stoichiometric equations [24] where:

$$\text{Carbohydrate metabolism} = \left(1.67 \times \dot{V}O_2\right) - \left(1.67 \times \dot{V}CO_2\right) \text{ g}\cdot\text{min}^{-1} \text{ and}$$

$$\text{Fat metabolism} = \left(6.21 \times \dot{V}CO_2\right) - \left(4.21 \times \dot{V}O_2\right) \text{ g}\cdot\text{min}^{-1}$$

Fat metabolism was plotted against the exercise stage for each participant's individualised ACE and CE protocols and the exercise intensity ($\% \dot{V}O_{2\text{peak}}$) at which the MFO occurred was determined (i.e., Fat_{max}). Ratings of perceived exertion were recorded at the end of each exercise stage for RPE_L and central (cardiorespiratory) exertion (RPE_C). The RPE scale and verbal anchors were fully explained to each participant. Capillary blood samples (20 μL) were taken from the participant's earlobe at rest, at volitional exhaustion and following 5 min of active recovery for the analysis of blood lactate concentration (Biosen, C_Line analyser, EFK Diagnostics, Cardiff, UK). Blood sampling was undertaken in accordance with the British Association of Sport and Exercise Science guidelines [25].

2.4. Data Analysis

Data are expressed as mean \pm standard deviation (SD). All data were analysed using the Statistical Package for the Social Sciences (v22; IBM Inc., Chicago, USA). Data were checked for normality using the Shapiro–Wilkes test and the homogeneity of variance using Levene's test. Peak physiological and perceptual responses, maximal carbohydrate oxidation, MFO and Fat_{max} were compared between exercise modes using paired t-tests. Data for physiological variables at each exercise stage and between exercise modes were analysed using factorial analysis of variance (ANOVA), with repeated measures on both factors (Mode: ACE, CE; \times intensity; rest, 40, 50, 60, 70, 80, 90, 100% PO_{peak}). Where significance was achieved, Tukey's post-hoc analysis was undertaken to determine the difference required between means for significance at the level of $p < 0.05$ [26].

Due to the nature of the exercise protocols increasing by 10% PO_{peak} at each stage, the data could not be statistically compared at a given power output across exercise modes. Therefore, the data for HR, $\dot{V}O_2$, carbohydrate oxidation and fat oxidation were also plotted for illustrative purposes only against the mean power output representing each 10% increment without statistical analysis. Data for the 2- and 3-minute comparison during ACE for intensities completed by all participants (i.e., up to 70% PO_{peak}) were analysed by factorial analysis of variance (ANOVA), with repeated measures on both factors (intensity; 40, 50, 60, 70% $\dot{V}O_{2\text{peak}}$; \times stage duration; 2-min, 3-min).

3. Results

3.1. Peak Physiological Responses

The peak physiological responses for ACE and CE are shown in Table 1. Peak $\dot{V}O_2$, $\dot{V}E$, power output and heart rate (HR_{peak}) were lower during ACE than for CE ($p < 0.05$). No differences were observed between modes for peak RER, blood lactate concentration or RPE_L , whereas RPE_C approached significance ($p = 0.065$), with RPE_C tending to be lower during ACE compared to CE. Carbohydrate oxidation at volitional exhaustion was similar for ACE and CE (4.91 ± 0.73 and 5.16 ± 1.88 $\text{g}\cdot\text{min}^{-1}$ for ACE and CE, respectively) ($p = 0.661$). Fat oxidation at the end of the exercise was also similar between trials (ACE: 0.06 ± 0.42 $\text{g}\cdot\text{min}^{-1}$; CE: 0.13 ± 0.67 $\text{g}\cdot\text{min}^{-1}$; $p = 0.781$).

Table 1. Peak physiological responses for arm crank ergometry (ACE) and cycle ergometry (CE) in young, healthy males.

	ACE	CE	<i>p</i>
	Mean ± SD	Mean ± SD	
$\dot{V}O_{2\text{peak}}$ (L·min ⁻¹)	2.73 ± 0.64	3.62 ± 0.46	<0.001
$\dot{V}O_{2\text{peak}}$ (mL·kg ⁻¹ ·min ⁻¹)	36 ± 7	48 ± 5	<0.001
$\dot{V}E_{\text{peak}}$ (L·min ⁻¹)	100.4 ± 19.6	120.1 ± 19.4	0.003
PO_{peak} (W)	133 ± 21	239 ± 33	<0.001
HR_{peak} (beats·min ⁻¹)	170 ± 16	182 ± 11	0.020
RPE _C (Borg Scale)	16 ± 3	18 ± 2	0.065
RPE _L (Borg Scale)	19 ± 1	19 ± 1	0.120
BLA _{peak} (mmol·L ⁻¹)	7.6 ± 1.9	9.0 ± 2.9	0.134
BLA+5 (mmol·L ⁻¹)	8.0 ± 1.7	8.4 ± 3.0	0.179
RER _{peak}	1.00 ± 0.10	0.96 ± 0.13	0.328

Note: $\dot{V}O_{2\text{peak}}$ (L·min⁻¹) = absolute peak oxygen uptake; $\dot{V}O_{2\text{peak}}$ (mL·kg⁻¹·min⁻¹) = relative peak oxygen uptake; $\dot{V}E_{\text{peak}}$ (L·min⁻¹) = peak minute ventilation; PO_{peak} (W) = peak power output; HR_{peak} (beats·min⁻¹) = peak heart rate; RPE_C (Borg Scale) = peak rating of perceived exertion (central/cardiorespiratory effort); RPE_L (Borg Scale) = peak rating of perceived exertion (local muscular effort); BLA_{peak} (mmol·L⁻¹) = peak blood lactate concentration, BLA+5 (mmol·L⁻¹) = blood lactate concentration 5-min post-cessation of exercise; RER_{peak} = peak respiratory exchange ratio.

3.2. Submaximal Exercise Responses

Heart rate, $\dot{V}O_2$ and RER responses during incremental ACE and CE are shown in Figure 1. When expressed relative to peak power (i.e., % PO_{peak}), $\dot{V}O_2$ was lower during ACE than for CE at every exercise intensity ($p < 0.05$). No mode × intensity interactions were observed for HR ($p = 0.229$), RER ($p = 0.498$), RPE_C ($p = 0.892$) or RPE_L ($p = 0.969$) when expressed relative to % PO_{peak} . Main effects for intensity were observed for HR, RPE_C, RPE_L and RER ($p < 0.05$), with additional main effects for exercise modes observed for HR and RPE_C ($p < 0.05$). When plotted against absolute power output, $\dot{V}O_2$ for ACE and CE was similar where the power output overlapped (~90 to 140 W), whereas HR, RPE_C, RPE_L and RER demonstrated greater values during ACE.

3.3. Substrate Utilisation

The resting respiratory exchange ratio was similar between trials (0.78 ± 0.11 vs. 0.78 ± 0.06 for ACE and CE, respectively; $p = 0.978$), as were carbohydrates (0.16 ± 0.28 vs. 0.17 ± 0.36 g·min⁻¹, respectively) and fat oxidation (0.17 ± 0.07 vs. 0.16 ± 0.06 g·min⁻¹, respectively). No mode × intensity interaction was observed for carbohydrate oxidation ($p = 0.924$), although values increased with exercise intensity (intensity main effect, $p < 0.01$). When plotted as absolute exercise intensities, carbohydrate oxidation was greater during ACE (Figure 1).

A significant mode × intensity interaction was observed for fat oxidation ($p = 0.013$), with differences observed between modes for each exercise intensity ($p < 0.01$). Maximal fat oxidation for ACE and CE modes is shown in Figure 2. Maximal fat oxidation was lower in ACE (0.44 ± 0.24 g·min⁻¹; range: 0.11–0.79 g·min⁻¹) when compared to CE (0.77 ± 0.31 g·min⁻¹; range: 0.33–1.23 g·min⁻¹; $p < 0.01$), occurring at 53 ± 21 and $67 \pm 18\%$ $\dot{V}O_{2\text{peak}}$, respectively ($p < 0.01$).




Figure 1. Heart rate (HR), oxygen uptake ($\dot{V}O_2$) and respiratory exchange ratio (RER) responses to incremental arm crank (ACE) and cycle ergometry (CE). Data plotted in relation to %PO_{peak} (left panel) and absolute power output (W) during exercise (right panel).

The relationship between resting fat oxidation and MFO for both modes of exercise combined was significant ($R = 0.505$; $R^2 = 0.254$, $p = 0.020$), whereas when considered for ACE and CE separately, only CE elicited a significant relationship (ACE: $R = 0.520$; $R^2 = 0.270$, $p = 0.151$; CE: $R = 0.661$; $R^2 = 0.436$, $p = 0.019$). Based on the coefficient of variation, the inter-participant variability for MFO was greater during ACE (CV = 54.5%) when compared to CE (CV = 40.3%). However, inter-individual variation for MFO was considerably greater than for peak physiological responses such as $\dot{V}O_{2peak}$ during ACE and CE (12.7% and 23.4%, respectively).

3.4. Comparison of 2- and 3-Minute Exercise Durations

The physiological responses to submaximal ACE during 2- and 3-minute exercise stages are shown in Table 2. No stage duration \times intensity interactions were observed for HR, $\dot{V}O_2$, $\dot{V}CO_2$, RER, RPE_C, RPE_L, carbohydrate oxidation or fat oxidation during the submaximal ACE trials (all $p > 0.05$; Table 2). However, with the exception of fat oxidation, all physiological and perceptual responses increased with exercise intensity (main effects for intensity; all $p < 0.05$).




Figure 2. Carbohydrate and fat oxidation responses to incremental arm crank (ACE) and cycle ergometry (CE). Data plotted in relation to %PO_{peak} (left panel) and absolute power output (W) during exercise (right panel). NB: Of the n = 13 participants, only n = 9 completed the final exercise stage at 100%PO_{peak}; therefore, data for ACE at 100% PO_{peak} and absolute power output of 133 W are for the remaining n = 9.

Table 2. Physiological responses (mean ± SD) for submaximal intensities (%PO_{peak}) of arm crank ergometry (ACE) during 2- and 3-minute exercise stages in young, healthy males (n = 9).

	Stage	%PO _{peak}				%PO _{peak}	p Stage	Int.
		40	50	60	70			
HR (beats·min ⁻¹)	2 min	103 ± 22	122 ± 19	135 ± 22	151 ± 22	<0.05	0.556	0.997
	3 min	107 ± 23	125 ± 19	138 ± 22	152 ± 23			
V̇O ₂ (L·min ⁻¹)	2 min	0.65 ± 0.11	1.04 ± 0.30	1.34 ± 0.31	1.58 ± 0.33	<0.05	0.983	0.991
	3 min	0.62 ± 0.11	1.07 ± 0.35	1.34 ± 0.33	1.58 ± 0.35			
V̇CO ₂ (L·min ⁻¹)	2 min	0.57 ± 0.16	0.94 ± 0.21	1.23 ± 0.25	1.43 ± 0.31	<0.05	0.773	0.919
	3 min	0.57 ± 0.15	1.03 ± 0.22	1.23 ± 0.26	1.42 ± 0.41			
RER	2 min	0.83 ± 0.03	0.88 ± 0.03	0.95 ± 0.03	0.96 ± 0.03	<0.05	0.145	0.774
	3-min	0.83 ± 0.03	0.92 ± 0.03	0.96 ± 0.04	0.95 ± 0.04			
FOx (g·min ⁻¹)	2 min	0.16 ± 0.04	0.20 ± 0.06	0.22 ± 0.08	0.31 ± 0.14	0.324	0.963	0.986
	3 min	0.16 ± 0.04	0.17 ± 0.06	0.25 ± 0.10	0.33 ± 0.15			
CHO (g·min ⁻¹)	2 min	0.81 ± 0.78	1.49 ± 0.63	2.00 ± 0.82	2.20 ± 1.57	0.003	0.692	0.918
	3 min	0.92 ± 0.96	1.88 ± 0.83	1.95 ± 1.02	2.15 ± 1.62			
RPE _C (Borg Scale)	2 min	7.2 ± 0.7	8.8 ± 1.0	10.6 ± 1.3	12.6 ± 1.1	<0.05	0.145	0.774
	3 min	7.2 ± 0.7	9.1 ± 1.1	10.2 ± 1.3	13.0 ± 1.6			
RPE _L (Borg Scale)	2 min	8.4 ± 1.2	11.0 ± 1.2	13.1 ± 0.8	14.3 ± 0.9	<0.05	0.088	0.968
	3 min	8.9 ± 1.1	12.0 ± 1.0	14.3 ± 0.8	14.8 ± 0.8			

Note: HR (beats·min⁻¹) = heart rate; V̇O₂ (L·min⁻¹) = absolute oxygen uptake; V̇CO₂ (L·min⁻¹) = absolute carbon dioxide production; RER = respiratory exchange ratio; FOx (g·min⁻¹) = fat oxidation; CHO (g·min⁻¹) = carbohydrate oxidation; RPE_C (Borg Scale) = rating of perceived exertion (central/cardiorespiratory effort); RPE_L (Borg Scale) = rating of perceived exertion (local muscular effort); p = p-value; %PO_{peak} = exercise intensity main effect (40 vs. 50 vs. 60 vs. 70% PO_{peak}); Stage = stage duration main effect (2-min vs. 3-min), Int = %PO_{peak} × Stage interaction.

4. Discussion

This is the first study to evaluate the magnitude of MFO and Fat_{max} during individualised incremental exercise tests for upper and lower body exercise in young, recreationally active males. The magnitudes of both MFO and Fat_{max} were lower during ACE than CE but greater than those for previous studies involving overnight fasts. The variability of MFO and Fat_{max} for ACE was greater than for CE.

4.1. The Peak Physiological Responses

The peak physiological responses to incremental upper and lower body exercise were indicative of those expected for the population studied, both from our laboratory and others [10,12,22,27–29]. In brief, $\dot{V}O_{2peak}$ and PO_{peak} for ACE were approximately 75% and 55% of the values observed for CE, respectively, and within the expected range [10,13,27]. Lower peak physiological values for ACE, when compared to CE, are a result of the smaller muscle mass and maximal cardiac output and a greater local fatigue component during upper body exercise [12,30–32]. As such, these factors prevent the central cardiorespiratory responses from eliciting the greater values observed during lower body exercise. The peak physiological responses were, therefore, as expected for the population studied.

4.2. Incremental Exercise Responses

During incremental exercise, $\dot{V}O_2$ was lower during ACE than CE when expressed in relation to $\%PO_{peak}$. The lower $\dot{V}O_2$ at each stage of the individualised incremental exercise tests is a result of the lower $\dot{V}O_{2peak}$ during ACE and likely systematic. Indeed, the difference between $\dot{V}O_{2peak}$ values for ACE and CE was similar to the difference between modes for each individualised exercise stage ($\sim 1.0 \text{ L}\cdot\text{min}^{-1}$). Conversely, HR was greater during ACE than CE at each individualised stage, although representing a similar systematic exercise mode difference (overall mean between modes across exercise intensities $\sim 8 \text{ beats}\cdot\text{min}^{-1}$). When plotted in relation to absolute power output, $\dot{V}O_2$ was similar at the exercise intensities accomplished during both upper and lower body exercise (~ 90 to 140 W), reflecting the overall energy requirements for a given power output [11].

During ACE at a given power output, HR, RER, carbohydrate metabolism and RPE_L were greater and fat oxidation was lower when compared to CE. Such responses illustrate the greater physiological strain at a given power output when exercising with the smaller muscle mass of the arms during ACE when compared to the legs during CE [10,12,27,28]. Subsequently, the lower RPE_C during both individualised and absolute power outputs for ACE is a result of the resultant peripheral limitation during this exercise mode [10]. The physiological and perceptual responses to incremental ACE and CE exercise are, thus, in line with previous research.

4.3. Maximal Fat Oxidation during CE

The MFO values observed for CE in the current study ($\sim 0.77 \text{ g}\cdot\text{min}^{-1}$) are greater than those reported for previous studies of CE and treadmill exercise (0.43 and $0.65 \text{ g}\cdot\text{min}^{-1}$, respectively; [14]) and those reported in a recent review of MFO (endurance-trained males $\sim 0.53 \text{ g}\cdot\text{min}^{-1}$; recreationally active males $\sim 0.43 \text{ g}\cdot\text{min}^{-1}$; [7]). However, the mean MFO values are well within the wide range of values reported for males in the literature (i.e., 0.17 to $1.27 \text{ g}\cdot\text{min}^{-1}$; mean = $0.60 \text{ g}\cdot\text{min}^{-1}$ [6,33–35]). The greater MFO values in the current study may be a result of the resting metabolic state of the participants. Resting RER values (~ 0.78) were indicative of a predominance of fat metabolism, also evidenced by resting fat oxidation values ($\sim 0.17 \text{ g}\cdot\text{min}^{-1}$) being greater than those previously reported ($0.06 \text{ g}\cdot\text{min}^{-1}$; [36]). Rosenkilde et al. [37] observed that participants with lower resting RER values (0.76) elicited greater MFO than those with greater resting RER values (0.86). Furthermore, Robinson et al. [33] observed a relationship between 24-hour fat oxidation and MFO, suggesting habitual diet may be an important contributor to the maximal rate of fat oxidation during exercise.

The current study demonstrated a correlation between resting fat oxidation and MFO values, confirming that greater resting fat oxidation resulted in greater maximal fat oxidation. Thus, although participants in the current study were asked not to eat for only 2–3 h prior to exercise, it is possible that their habitual diets affected their resting and subsequent exercising metabolic substrates. However, as most individuals are unlikely to exercise after an overnight fast, the current MFO and Fat_{max} values are greater than expected for fasted conditions. The values are still within the expected range of reported values reflecting true maximal fat oxidation rates in daily, ecologically valid conditions.

4.4. Maximal Fat Oxidation during ACE

Maximal fat oxidation rates and Fat_{max} were lower during ACE than CE and are in agreement with the current hypothesis. Few studies have reported Fat_{max} during upper body exercise, except for those that have reported values for control groups matched to diabetic or obese experimental groups [17,18] or those with spinal cord injury [19]. For example, Ara et al. [17] reported lower MFO values for incremental ACE than for CE (0.18 vs. 0.28 $g \cdot min^{-1}$) in previously obese men (31 y, $\dot{V}O_{2peak}$ during CE: 27 $mL \cdot kg^{-1} \cdot min^{-1}$). Larsen et al. [18] also reported lower MFO values in ACE than CE (0.18 vs. 0.30 $g \cdot min^{-1}$) in age- and body-mass-matched controls (43 y, $\dot{V}O_{2peak}$ during CE: 42 $mL \cdot kg^{-1} \cdot min^{-1}$). Both studies reported Fat_{max} at relatively low percentages of $\dot{V}O_{2peak}$ in both ACE and CE trials (~30 vs. ~40% $\dot{V}O_{2peak}$, respectively). However, the exercise intensity at which MFO occurred and the absolute MFO values represent similar ACE:CE ratios to the current study (i.e., 0.74 and 0.75, 0.79 and 0.64, and 0.60 and 0.57 for Ara et al. [17], Larsen et al. [18], and the current study, respectively).

Jacobs et al. [19] observed greater MFO values during ACE in sedentary participants with spinal cord injury compared to a non-disabled control group of men and women, suggesting that the greater training status of the upper body through habitual wheelchair locomotion may have adapted upper body fat metabolism. However, MFO and Fat_{max} values for both groups were still low (0.07 to 0.13 $g \cdot min^{-1}$ at 42% $\dot{V}O_{2peak}$ and 0.02 to 0.06 $g \cdot min^{-1}$ at 14% $\dot{V}O_{2peak}$, respectively). The greater overall training status of the current participants, based on $\dot{V}O_{2peak}$ values during ACE and CE (and no sympathetic nervous system dysfunction), would likely have contributed to the greater absolute MFO values observed. Thus, although our current understanding of MFO during ACE is from older and less aerobically conditioned individuals than in the present study, values across exercise modes appear to occur at similar proportions of maximal capacity and with a large potential for improvement in less active individuals.

4.5. Inter-Individual Variability

Usually, only small amounts of variation can be accounted for when examining factors that influence MFO and Fat_{max} . As with previous studies, the current study demonstrates a significant positive correlation between MFO and $\dot{V}O_{2peak}$ across both modes of exercise; however, the R^2 indicated that only 25% of the variation in MFO could be accounted for. Although lower than the 35% and 47% of variation accounted for in MFO for more than one contributing component by Randell et al. [33] and Venables et al. [6], respectively, our R^2 value is similar in that much of the variation cannot be accounted for. Key factors reported to be related to MFO include maximal $\dot{V}O_2$, body fat percentage, fat-free mass, fast duration, self-reported physical activity and sex [6,33]. Randell et al. [33] recently indicated that genetic or epigenetic factors may account for a large proportion of the unexplained variance and should be explored further.

The inter-individual variation for MFO in the current study, as indicated by the coefficient of variation, was extremely broad, with ACE being greater than CE (54.5 vs. 40.3%, respectively) and much greater than for $\dot{V}O_{2peak}$ per se (23.4 vs. 12.7%, respectively). When compared to CE, the variability of MFO may be further confounded during upper

body exercise when considering individual fibre type differences and the greater range of training statuses.

4.6. Application

The main application of the current data is in determining potential aerobic exercise training intensities for optimising fat metabolism and enabling weight loss. A small number of studies have examined training regimes utilising Fat_{max} exercise intensity [38–41]. These training studies have predominantly involved overweight or obese participants, subsequently focusing upon a range of health-related outcome measures: observing improvements in body composition, cardiovascular function, strength and flexibility [39,40]. Within these studies, Fat_{max} ranged between 44–54% $\dot{V}O_{2max}$ [38–40] for lower body exercise, which is lower than for the current young, recreationally active participants during CE (67% $\dot{V}O_{2peak}$), most likely due to their aerobic fitness status. Although upper body Fat_{max} intensities for obese, previously obese and age- and mass-matched control participants are lower than for lower body exercise [17], the use of Fat_{max} as an upper body training intensity has not been reported.

Using a traditional aerobic exercise training intensity (60% PO_{peak}) in a similar population to the current study, Bottoms [42] observed increased free fatty acid concentrations at the end of 30 min continuous ACE, following 8 weeks of aerobic ACE training. The equivalent exercise intensity post-training was $\sim 57\% \dot{V}O_{2peak}$ and similar to that for ACE Fat_{max} in the current study (53% $\dot{V}O_{2peak}$). Furthermore, Bottoms [42] utilised two continuous training sessions and one interval training session per week. In light of the finding that continuous low-intensity exercise training sessions can improve fat metabolism when compared to interval-type training, the ratio of 2:1 for continuous to interval exercise training sessions appears potentially useful for improving fat metabolism in young, healthy, recreationally active male participants. The continuous- and interval-type exercise training comparison noted above, however, was undertaken in overweight men, so the finding needs to be confirmed for healthy individuals.

4.7. Limitations

Firstly, this study examined MFO and Fat_{max} values in male participants only. Previous studies of trained males and females during prolonged exercise at different intensities (25, 65, 85% $\dot{V}O_{2max}$) observed similar total fat oxidation responses, with the greatest values occurring at 65% $\dot{V}O_{2max}$ [43,44]. More recently, similar MFOs in young, trained men and women, despite major differences in plasma lipid concentrations during graded exercise, were also observed [45].

In contrast, a review comparing MFO and Fat_{max} between men and women reported MFO to be greater in males and Fat_{max} to be greater in females [7], though their study did not take into account the separate effects of age, training status, mode of exercise and weight status [46]. Although there are no reported studies on fat metabolism during incremental ACE in females, it is clear that peak physiological responses during ACE in females are lower, but with a similar ratio between ACE and CE, with $\dot{V}O_{2peak}$ for ACE being $\sim 63\%$ [47] to 68% of CE values [48]. It has also been reported that females are 52% and 66% as strong as their male counterparts in the upper body (elbow flexors) and lower body (knee flexors), respectively, which is due to larger fibre areas in males [49]. Although differences in the strength and aerobic power of females vary to a greater extent between the upper and lower body compared to men, there are no data to suggest sex differences in MFO and Fat_{max} for the upper body. A similar response in MFO and Fat_{max} may occur for upper body exercise, as observed for males, although this remains to be tested, particularly in consideration of the effects of sex hormones observed on metabolism [50].

Secondly, the current study comprised a heterogenous group with respect to training status, more so for upper body exercise capacity. Thus, examining homogenous upper body

trained groups or potential changes in fat oxidation before and after upper body aerobic training may be more enlightening.

Thirdly, although resting fat oxidation and RER values were similar between ACE and CE trials, habitual dietary habits were not recorded, which may help explain why the MFO values reported were at the higher end of the expected physiological range.

Finally, fat oxidation values at volitional exhaustion had not reached zero, with participants potentially not having reached true volitional exhaustion. However, with the exception of RER_{peak} values, typical end criteria for ACE and CE exercise protocols were achieved, with values in line with previous studies. Furthermore, at volitional exhaustion, fat oxidation had returned to similar values as those observed at baseline. If greater peak exercise capacity had been achieved, it is likely that this would only have been one, or part of one, further exercise stage that would not affect the MFO values per se and would only slightly reduce the relative exercise intensity at which it occurred (i.e., Fat_{max}). We are, therefore, confident we can state that both MFO and Fat_{max} during ACE are lower than for CE and demonstrate greater variability.

4.8. Future Research

The following areas for future research are recommended. The short duration pre-exercise fast and non-specific dietary regime of the current study likely contributed to the relationship between resting fat oxidation and MFO, prompting the suggestion that habitual diet composition should be considered. Future examination of MFO during exercise, following participants' habitual diets and standardised diets, as well as ecologically valid fast durations, will aid our understanding of dietary factors affecting MFO. The potential to maximise MFO during both ACE and CE will also be enhanced with this approach, along with our understanding of the individual variation in MFO during both upper and lower body exercise.

Future studies on MFO and Fat_{max} adaptations during upper body exercise training in otherwise healthy individuals will provide an adjunct to traditional training modes from which additional physiological and functional benefits (e.g., activities of daily living) may be achieved. Similarly, the application of upper body exercise training to clinical populations allows the potential for improving fat metabolism in populations either without the use or with limited use of their legs.

Further studies to examine the potential underlying genetic and epigenetic factors across a range of exercise modes and populations are also recommended to help understand the intra- and inter-individual variability of MFO and Fat_{max}.

5. Conclusions

The results of this study show consistently lower MFO and Fat_{max} during upper body exercise in young, healthy, recreationally active male participants compared to lower body exercise. Although inter-individual variation was greater for MFO and Fat_{max} during upper body exercise, these data have important applications for weight loss strategies in healthy male populations and those unable to use their legs or where upper body exercise may be the only available exercise mode.

Author Contributions: Conceptualisation, M.P. and L.B.; methodology, M.P.; formal analysis, M.P. and L.B.; investigation, M.P.; writing—original draft preparation, M.P.; writing—review and editing, L.B., M.H. and R.E.; visualisation, L.B. and M.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Coventry University (P35579; 21/10/2015) for studies involving humans.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: The authors would like to thank the following people for their assistance in participant recruitment and data collection: Jack Dicker, Alfie Dwight, Devann Edwards, Bipin Modha, Gareth Peden and Shaun Phillips.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Knechtle, B.; Müller, G.; Willmann, F.; Eser, P.; Knecht, H. Comparison of fat oxidation in arm cranking in spinal cord-injured people versus ergometry in cyclists. *Eur. J. Appl. Physiol.* **2003**, *90*, 614–619. [[CrossRef](#)] [[PubMed](#)]
2. Achten, J.; Jeukendrup, A.E. Optimizing fat oxidation through exercise and diet. *Nutrition* **2004**, *20*, 716–727. [[CrossRef](#)] [[PubMed](#)]
3. Achten, J.; Gleeson, M.; Jeukendrup, A.E. Determination of the exercise intensity that elicits maximal fat oxidation. *Med. Sci. Sports Exerc.* **2002**, *34*, 92–97. [[CrossRef](#)] [[PubMed](#)]
4. Alkahtani, S.A.; King, N.A.; Hills, A.P.; Byrne, N.M. Effect of interval training intensity on fat oxidation, blood lactate and the rate of perceived exertion in obese men. *SpringerPlus* **2013**, *2*, 532. [[CrossRef](#)] [[PubMed](#)]
5. Achten, J.; Jeukendrup, A.E. The effect of pre-exercise carbohydrate feedings on the intensity that elicits maximal fat oxidation. *J. Sports Sci.* **2003**, *21*, 1017–1024. [[CrossRef](#)]
6. Venables, M.C.; Achten, J.; Jeukendrup, A.E. Determinants of fat oxidation during exercise in healthy men and women: A cross-sectional study. *J. Appl. Physiol.* **2005**, *98*, 160–167. [[CrossRef](#)]
7. Maunder, E.; Plews, D.J.; Kilding, A.E. Contextualising maximal fat oxidation during exercise: Determinants and normative values. *Front. Physiol.* **2018**, *9*, 599. [[CrossRef](#)]
8. Croci, I.; Borrani, F.; Byrne, N.M.; Wood, R.E.; Hickman, I.J.; Chenevière, X.; Malatesta, D. Reproducibility of fatmax and fat oxidation rates during exercise in recreationally trained males. *PLoS ONE* **2014**, *9*, e97930. [[CrossRef](#)] [[PubMed](#)]
9. Helgerud, J.; Øiestad, B.E.; Wang, E.; Hoff, J. Prediction of upper extremity peak oxygen consumption from heart rate during submaximal arm cycling in young and middle-aged adults. *Eur. J. Appl. Physiol.* **2019**, *119*, 2589–2598. [[CrossRef](#)] [[PubMed](#)]
10. Price, M.; Beckford, C.; Dorricott, A.; Hill, C.; Kershaw, M.; Singh, M.; Thornton, I. Oxygen uptake during upper body and lower body Wingate anaerobic tests. *Appl. Phys. Nutr. Metab.* **2013**, *39*, 1345–1351. [[CrossRef](#)] [[PubMed](#)]
11. Sawka, M.N. Physiology of upper body exercise. *Exerc. Sport Sci. Rev.* **1986**, *14*, 175–211. [[CrossRef](#)] [[PubMed](#)]
12. Davis, J.A.; Vodak, P.; Wilmore, J.H.; Vodak, J.; Kurtz, P. Anaerobic threshold and maximal aerobic power for three modes of exercise. *J. Appl. Physiol.* **1976**, *41*, 544–550. [[CrossRef](#)] [[PubMed](#)]
13. Larsen, R.T.; Christensen, J.; Tang, L.H.; Keller, C.; Doherty, P.; Zwisler, A.D.; Taylor, R.S.; Langberg, H. A systematic review and meta-analysis comparing cardiopulmonary exercise test values obtained from the arm cycle and the leg cycle respectively in healthy adults. *Int. J. Sports Phys. Ther.* **2016**, *11*, 1006–1039. [[PubMed](#)]
14. Achten, J.; Venables, M.C.; Jeukendrup, A.E. Fat oxidation rates are higher during running compared with cycling over a wide range of intensities. *Metab. Clin. Exp.* **2003**, *52*, 747–752. [[CrossRef](#)]
15. Sanchis-Moysi, J.; Idoate, F.; Olmedillas, H.; Guadalupe-Grau, A.; Alayón, S.; Carreras, A.; Dorado, C.; Calbet, J.A. The upper extremity of the professional tennis player: Muscle volumes, fiber-type distribution and muscle strength. *Scand. J. Med. Sci. Sports* **2010**, *20*, 524–534. [[CrossRef](#)]
16. Koppo, K.; Bouckaert, J.; Jones, A.M. Oxygen uptake kinetics during high-intensity arm and leg exercise. *Resp Physiol Neurobiol* **2002**, *133*, 241–250. [[CrossRef](#)]
17. Ara, I.; Larsen, S.; Stallknecht, B.; Guerra, B.; Morales-Alamo, D.; Andersen, J.L.; Ponce-González, J.G.; Guadalupe-Grau, A.; Galbo, H.; Calbet, J.A.; et al. Normal mitochondrial function and increased fat oxidation capacity in leg and arm muscles in obese humans. *Int. J. Obes.* **2011**, *35*, 99–108. [[CrossRef](#)] [[PubMed](#)]
18. Larsen, S.; Ara, I.; Rabøl, R.; Andersen, J.L.; Boushel, R.; Dela, F.; Helge, J.W. Are substrate use during exercise and mitochondrial respiratory capacity decreased in arm and leg muscle in type 2 diabetes? *Diabetologia* **2009**, *52*, 1400–1408. [[CrossRef](#)] [[PubMed](#)]
19. Jacobs, K.A.; Burns, P.; Kressler, J.; Nash, M.S. Heavy reliance on carbohydrate across a wide range of exercise intensities during voluntary arm ergometry in persons with paraplegia. *J. Spinal Cord Med.* **2013**, *36*, 427–435. [[CrossRef](#)] [[PubMed](#)]
20. Amaro-Gahete, F.; Sanchez-Delgado, G.; Jurado-Fasoli, L.; De-la-O, A.; Castillo, M.J.; Helge, J.W.; Ruiz, J.R. Assessment of maximal fat oxidation during exercise: A systematic review. *Scand. J. Med. Sci. Sports* **2019**, *29*, 910–921. [[CrossRef](#)]
21. Smith, P.M.; Price, M.J. Upper body exercise. In *Sport and Exercise Physiology Testing Guidelines, the British Association of Sports and Exercise Sciences Guide: Sports Testing*; Winter, E.M., Jones, A.M., Davidson, R.C.R., Bromley, P.D., Mercer, T.H., Eds.; Routledge: London, UK, 2007; Volume 1, pp. 138–144.
22. Lyons, S.; Richardson, M.; Bishop, P.; Smith, J.; Heath, H.; Giesen, J. Excess post-exercise oxygen consumption in untrained men following exercise of equal energy expenditure: Comparisons of upper and lower body exercise. *Diab. Obes. Metab.* **2007**, *9*, 889–894. [[CrossRef](#)] [[PubMed](#)]
23. Borg, G. Perceived exertion as an indicator of somatic stress. *Scand. J. Rehab. Med.* **1970**, *2*, 92–98.
24. Frayn, K.N. Calculation of substrate oxidation rates in vivo from gaseous exchange. *J. Appl. Physiol.* **1983**, *55*, 628–634. [[CrossRef](#)] [[PubMed](#)]

25. Maughan, R.J.; Shirreffs, S.M.; Leiper, J.B. Blood sampling. In *Sport and Exercise Physiology Testing Guidelines, the British Association of Sports and Exercise Sciences Guide: Exercise and Clinical Testing*; Winter, E.M., Jones, A.M., Davidson, R.C.R., Bromley, P.D., Mercer, T.H., Eds.; Routledge: London, UK, 2007; Volume 2, pp. 25–29.
26. Vincent, W.J. *Statistics in Kinesiology*, 2nd ed.; Human Kinetics: Champaign, IL, USA, 1999; p. 163.
27. Sawka, M.N.; Pimental, N.A.; Pandolf, K.B. Thermoregulatory responses to upper body exercise. *Eur. J. Appl. Physiol.* **1984**, *52*, 230–234. [[CrossRef](#)] [[PubMed](#)]
28. Kang, J.; Robertson, R.J.; Goss, F.L.; Dasilva, S.G.; Suminski, R.R.; Utter, A.C.; Zoeller, R.F.; Metz, K.F. Metabolic efficiency during arm and leg exercise at the same relative intensities. *Med. Sci. Sports Exerc.* **1997**, *29*, 377–382. [[CrossRef](#)] [[PubMed](#)]
29. Sedlock, D.A.; Schneider, D.A.; Gass, E.; Gass, G. Excess post-exercise oxygen consumption in spinal cord-injured men. *Eur. J. Appl. Physiol.* **2004**, *93*, 231–236. [[CrossRef](#)] [[PubMed](#)]
30. Calbet, J.A.; González-Alonso, J.; Helge, J.W.; Søndergaard, H.; Munch-Andersen, T.; Saltin, B.; Boushel, R. Central and peripheral hemodynamics in exercising humans: Leg vs. arm exercise. *Scand. J. Med Sci. Sports* **2015**, *25* (Suppl. 4), 144–157. [[CrossRef](#)] [[PubMed](#)]
31. Sanada, K.; Kearns, C.F.; Kojima, K.; Abe, T. Peak oxygen uptake during running and arm cranking normalized to total and regional skeletal muscle mass measured by magnetic resonance imaging. *Eur. J. Appl. Physiol.* **2005**, *93*, 687–693. [[CrossRef](#)] [[PubMed](#)]
32. Eston, R.G.; Brodie, D.A. Responses to arm and leg ergometry. *Br. J. Sports Med.* **1986**, *20*, 4–6. [[CrossRef](#)] [[PubMed](#)]
33. Randell, R.K.; Rollo, I.; Roberts, T.J.; Dalrymple, K.J.; Jeukendrup, A.E.; Carter, J.M. Maximal fat oxidation rates in an athletic population. *Med. Sci. Sports Exerc.* **2017**, *49*, 133–140. [[CrossRef](#)]
34. Robinson, S.L.; Hattersley, J.; Frost, G.S.; Chambers, E.S.; Wallis, G.A. Maximal fat oxidation during exercise is positively associated with 24-hour fat oxidation and insulin sensitivity in young, healthy men. *J. Appl. Physiol.* **2015**, *118*, 1415–1422. [[CrossRef](#)] [[PubMed](#)]
35. Fletcher, G.; Eves, F.F.; Glover, E.I.; Robinson, S.L.; Vernooij, C.A.; Thompson, J.L.; Wallis, G.A. Dietary intake is independently associated with the maximal capacity for fat oxidation during exercise. *Am. J. Clin. Nutr.* **2017**, *105*, 864–872. [[CrossRef](#)] [[PubMed](#)]
36. Karppinen, J.E.; Rottensteiner, M.; Wiklund, P.; Hämäläinen, K.; Laakkonen, E.K.; Kaprio, J.; Kainulainen, H.; Kujala, U.M. Fat oxidation at rest and during exercise in male monozygotic twins. *Eur. J. Appl. Physiol.* **2019**, *119*, 2711–2722. [[CrossRef](#)] [[PubMed](#)]
37. Rosenkilde, M.; Nordby, P.; Nielsen, L.B.; Stallknecht, B.M.; Helge, J.W. Fat oxidation at rest predicts peak fat oxidation during exercise and metabolic phenotype in overweight men. *Int. J. Obes.* **2010**, *34*, 871–877. [[CrossRef](#)] [[PubMed](#)]
38. Venables, M.C.; Jeukendrup, A.E. Endurance training and obesity: Effect on substrate metabolism and insulin sensitivity. *Med. Sci. Sports Exerc.* **2008**, *40*, 495–502. [[CrossRef](#)] [[PubMed](#)]
39. Tan, S.; Wang, X.; Wang, J. Effects of supervised exercise training at the intensity of maximal fat oxidation in overweight young women. *J. Exerc. Sci. Fit.* **2012**, *10*, 64–69. [[CrossRef](#)]
40. Wang, J.; Tan, S.; Cao, L. Exercise training at the maximal fat oxidation intensity improved health-related physical fitness in overweight middle-aged women. *J. Exerc. Sci. Fit.* **2015**, *13*, 111–116. [[CrossRef](#)]
41. Botero, J.P.; Prado, W.L.; Guerra, R.L.; Speretta, G.F.; Leite, R.D.; Prestes, J.; Sanz, A.V.; Lyons, S.; de Azevedo, P.H.; Baldissera, V.; et al. Does aerobic exercise intensity affect health-related parameters in overweight women? *Clin. Physiol. Funct. Imag.* **2014**, *34*, 138–142. [[CrossRef](#)] [[PubMed](#)]
42. Bottoms, L.M. Thermoregulatory Responses during Upper Body Exercise, Thermal Stress, Training and Heat Acclimation. Ph.D. Thesis, Coventry University, Coventry, UK, 2008.
43. Romijn, J.A.; Coyle, E.F.; Sidossis, L.S.; Gastaldelli, A.; Horowitz, J.F.; Endert, E.; Wolfe, R.R. Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am. J. Physiol.* **1993**, *265*, E380–E391. [[CrossRef](#)]
44. Romijn, J.A.; Coyle, E.F.; Sidossis, L.S.; Rosenblatt, J.; Wolfe, R.R. Substrate metabolism during different exercise intensities in endurance-trained women. *J. Appl. Physiol.* **2000**, *88*, 1707–1714. [[CrossRef](#)]
45. Frandsen, J.; Poggi, A.I.; Ritz, C.; Larsen, S.; Dela, F.; Helge, J.W. Peak fat oxidation rate is closely associated with plasma free fatty acid concentrations in Women; Similar to men. *Front. Physiol.* **2021**, *12*, 696261. [[CrossRef](#)] [[PubMed](#)]
46. Amaro-Gahete, F.; Guillermo Sanchez-Delgado, G.; Ruiz, J. Commentary: Contextualising maximal fat oxidation during exercise: Determinants and normative values. *Front. Physiol.* **2018**, *9*, 599. [[CrossRef](#)]
47. Orr, J.L.; Williamson, P.; Anderson, W.; Ross, R.; McCafferty, S.; Fettes, P. Cardiopulmonary exercise testing: Arm crank vs cycle ergometry. *Anaesth* **2013**, *68*, 497–501. [[CrossRef](#)] [[PubMed](#)]
48. Bhambhani, Y.N. Prediction of stroke volume during upper and lower body exercise in men and women. *Arch. Phys. Med. Rehab.* **1995**, *76*, 713–718. [[CrossRef](#)]
49. Miller, A.E.; MacDougall, J.D.; Tarnopolsky, M.A.; Sale, D.G. Gender differences in strength and muscle fiber characteristics. *Eur. J. Appl. Physiol.* **1993**, *66*, 254–262. [[CrossRef](#)] [[PubMed](#)]
50. Devries, M.C. Sex-based differences in endurance exercise muscle metabolism: Impact on exercise and nutritional strategies to optimize health and performance in women. *Exp. Physiol.* **2016**, *101*, 243–249. [[CrossRef](#)] [[PubMed](#)]