

SWIFT IDENTIFICATION OF DARK GAMMA-RAY BURSTS

P. JAKOBSSON,^{1,2} J. HJORTH,¹ J. P. U. FYNBO,¹ D. WATSON,¹ K. PEDERSEN,¹ G. BJÖRNSSON,² AND J. GOROSABEL^{3,4}

Draft version November 6, 2007

ABSTRACT

We present an optical flux vs. X-ray flux diagram for all known gamma-ray bursts (GRBs) for which an X-ray afterglow has been detected. We propose an operational definition of dark bursts as those bursts that are optically subluminal with respect to the fireball model, i.e., which have an optical-to-X-ray spectral index $\beta_{\text{OX}} < 0.5$. Out of a sample of 52 GRBs we identify 5 dark bursts. The definition and diagram serve as a simple and quick diagnostic tool for identifying dark GRBs based on limited information, particularly useful for early and objective identification of dark GRBs observed with the *Swift* satellite.

Subject headings: dust, extinction — galaxies: high-redshift — gamma rays: bursts

1. INTRODUCTION

Dark gamma-ray bursts (GRBs) remain one of the unresolved issues in GRB research. Shortly after the localization of the first GRB afterglows it became clear that not all GRBs were accompanied by detections of optical afterglows (OAs). In fact, a fairly large fraction, about 60–70% of well localized GRBs did not lead to detections at optical wavelengths (Fynbo et al. 2001b; Lazzati, Covino & Ghisellini 2002).

Various scenarios have been suggested in order to shed light on dark bursts. The *obscuration scenario* (e.g., Groot et al. 1998; Taylor et al. 1998) ascribes the failed OA detection to extinction. Although there is evidence from X-rays (Galama & Wijers 2001) and damped Ly α absorbers (e.g., Fig. 4 in Vreeswijk et al. 2004) of high column density of gas around many GRBs, the early high energy radiation from them and their afterglows can destroy the dust in their environment within a radius up to a few tens of parsecs (Waxman & Draine 2000; Fruchter, Krolik & Rhoads 2001; Perna, Lazzati & Fiore 2003). This would pave the way for the afterglow light, but dust in the host galaxy at larger distances could still lead to failure in detecting the OA. In the *high-redshift scenario*, as some fraction of bursts will be located beyond $z \gtrsim 5$ (e.g., Totani 1997; Wijers et al. 1998; Lamb & Reichart 2000), the UV band, which is strongly affected by absorption in the Ly α forest, is redshifted into the optical band. Finally, optical faintness can arise if the OA is intrinsically dark as may happen, e.g., if a relativistic ejecta is decelerated in a *low-density ambient medium* (e.g., Sari, Piran & Narayan 1998; Taylor et al. 2000).

The dark burst fraction places important constraints on the fraction of obscured star formation in the Universe (Djorgovski et al. 2001; Ramirez-Ruiz et al. 2002) and the structure of star-forming regions (Lamb 2001; Reichart & Price 2002). Statistical samples studied up to now are unfortunately quite heterogeneous due to

large differences in localisation accuracies, localisation time since the onset of the burst, and search strategies. Moreover, effects of observing conditions (e.g., lunar phase) have generally not been taken into account in statistical studies. In many cases, GRBs have been considered dark if no OA was detected, irrespective of how inefficient the search was. In fact, there is no generally accepted criterion for when a GRB is considered dark. With the launch of the *Swift* satellite it will be essential to have a quick diagnostic tool to flag dark bursts for immediate and/or detailed follow-up (including the near-IR bands) to ensure homogeneity of samples. In this Letter we present a GRB diagram of the optical flux (F_{opt}) vs. the X-ray flux (F_{X}) and propose that those bursts which are optically subluminal with respect to the fireball model, i.e., which have an optical-to-X-ray spectral index $\beta_{\text{OX}} < 0.5$, be defined as dark.

2. CURRENT STATUS

A popular working definition of dark bursts is to set a brightness limit at a given time after the GRB, e.g., $R > 23$ mag at 1–2 days (Djorgovski et al. 2001). Such definitions are necessarily somewhat arbitrary but catch the notion of darkness very well in that the magnitude limits and times correspond to typical search efforts and reaction times. Another approach has been to invoke a physical definition, specifically to require a dark burst to be a significantly obscured burst. It has been argued that GRB 970828 (Djorgovski et al. 2001) and GRB 000210 (Piro et al. 2002) were most likely dark because of optical obscuration.

Fynbo et al. (2001b) demonstrated that the majority ($\gtrsim 75\%$) of GRBs for which searches for optical afterglow had been unsuccessful were consistent with no detection if they were similar to dim bursts like GRB 000630 in the optical band (see their Fig. 3). Hjorth et al. (2002) found that the dim GRB 980613 had similar properties, i.e., it would have been classified as a dark burst had it not been for the relatively deep search efforts. The afterglow was neither strongly reddened nor at high redshift. This suggests that the classification of the majority of dark bursts was due to searches which simply were not sufficiently sensitive to detect the faint optical afterglows.

Berger et al. (2002) reached a similar conclusion for the dim GRB 020124 and ascribed the faintness to rapid de-

¹ Niels Bohr Institute, Astronomical Observatory, University of Copenhagen, Juliane Maries Vej 30, 2100 Copenhagen, Denmark

² Science Institute, University of Iceland, Dunhaga 3, 107 Reykjavík, Iceland

³ IAA-CSIC, P.O. Box 03004, E-18080 Granada, Spain

⁴ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

cay whereas Hjorth et al. (2003) demonstrated that the faintness was largely due to the fairly large redshift of $z = 3.2$ (although not sufficiently large for the burst to be dark due to Ly α absorption). Several studies of the rapidly localised HETE-2 burst GRB 021211 arrived at a similar result: it would have been classified as a dark burst due to its rapid fading, but was found to be very bright after ten minutes. It was not strongly reddened and at a moderate redshift (Fox et al. 2003; Li et al. 2003; Crew et al. 2003; Pandey et al. 2003).

In a study of all BeppoSAX bursts with Narrow Field Instruments follow-up, De Pasquale et al. (2003, hereafter D03) found that most optically faint bursts are also X-ray faint. Some, however, appear even fainter in the optical than expected from X-rays. In a comprehensive study, Rol (2004) concluded that most GRBs can be fitted with standard fireball models. Only three were inconsistent with all models, i.e., fainter than the faintest optical expectation from X-rays. These were classified as dark. In addition, Pedersen et al. (2004) have proposed that GRB 001025A, along with some other bursts, appear optically dark because their (X-ray) afterglow is faint and their synchrotron cooling break, ν_c , is located close to the X-ray band.

Recently, more homogeneous samples have been constructed based on BeppoSAX and HETE-2. Stratta et al. (2004) find a dark burst fraction of 4/13 for a sample of bright BeppoSAX bursts. The better search conditions offered by HETE-2, in particular since the Soft X-ray Camera started to deliver accurate and fast localisations, have resulted in this fraction decreasing further to of the order of 10% (Lamb et al. 2004) as anticipated by Fynbo et al. (2001b).

3. THE OPTICAL FLUX VS. THE X-RAY FLUX DIAGRAM

Previous working definitions have been motivated by what makes a burst dark: its faintness. However, in view of the results that a faint burst does not by itself belong to a separate class (notably GRBs 980613, 000630, 020124, and 021211) and the study of D03 that some bursts may be optically faint simply because they are intrinsically faint, it is clear that another parameter must be invoked. D03 used the ratio of optical to X-ray flux. Here we will use the optical-to-X-ray spectral index which is more directly related to physical properties of afterglows.

In the simplest fireball models, which have been successfully used to interpret the observed properties of GRB afterglows, the spectral index, β ($F_\nu \propto \nu^{-\beta}$), is governed by the energy distribution of electrons, p , and the location of ν_c (e.g., Sari, Piran & Narayan 1998):

$$\beta = \begin{cases} (p-1)/2 & \nu < \nu_c \\ p/2 & \nu > \nu_c \end{cases}.$$

This result is independent of whether the outflow is collimated or not, or whether the expansion takes place in a constant density or stellar wind environment. In GRB afterglows, the cooling break is frequently found to be located between the optical ($\sim 10^{14}$ Hz) and X-ray ($\sim 10^{18}$ Hz) regimes giving rise to a break in the spectral distribution somewhere between these two frequencies. In some cases, though, it is either positioned below the optical or above the X-rays.

The value of p is usually found to be larger than 2 ($p < 2$ is not ruled out but requires a high-energy cutoff in the electron energy distribution, see e.g., Dai & Cheng 2001) and smaller than 2.5. In a study of 36 BeppoSAX X-ray afterglows, Piro (2004) inferred an average value of $p = 2.26$. In this simple picture, the average β_{OX} (where the subscript 'O' stands for 'optical' and 'X' for 'X-ray') is expected to lie between 0.5 ($p = 2$, $\nu_c > 10^{18}$ Hz) and 1.25 ($p = 2.5$, $\nu_c < 10^{14}$ Hz).

In a plot of F_{opt} vs. F_X , optically subluminal bursts, i.e., bursts fainter than expected from the fireball model, will be situated below the line of constant $\beta_{\text{OX}} = 0.5$. In Fig. 1 we plot the $F_{\text{opt}}-F_X$ diagram for all known GRBs which have an X-ray detection and an optical detection or upper limit (as of August 2004). All data have been interpolated/extrapolated to 11 h (following D03), and are listed in Table 1. For the upper limits we have assumed a decay index of $\alpha = 1$ ($F_\nu \propto t^{-\alpha}$) in the extrapolation. We note that the significance level of reported upper limits vary between bursts, ranging between 2σ and 5σ .

All R -band magnitudes in Table 1 have been corrected for foreground (Galactic) extinction using the reddening maps of Schlegel, Finkbeiner & Davis (1998). At 11 h, the optical afterglow is usually sufficiently bright that the host galaxy contribution is negligible, but whenever possible we have used the host-subtracted magnitudes reported in the literature. For the BeppoSAX bursts we have taken the 1.6–10.0 keV X-ray flux at 11 h from D03 and calculated the flux density at 3 keV. The same procedure was carried out for the Rossi X-ray Timing Explorer (RXTE) bursts, except the 2–10 keV X-ray flux was obtained from various sources in the literature. For the XMM-Newton and Chandra (CXO) data the flux density at 3 keV was derived from the best-fit single power-law with Galactic absorption to the 2–10 keV data. This energy (3 keV) was chosen as it is relatively insensitive to absorption and requires very little extrapolation of the data since it is close to the center of the bandpass with respect to total counts, thus yielding a reliable flux density. Data from XMM-Newton were reduced in a standard way using the XMM-Newton Science Analysis System (SAS, version 6.0.0) and the latest calibration files. The CXO data were reduced in a standard way using the Chandra Interactive Analysis of Observations (CIAO, version 3.0.2) and the latest calibration files (CALDB, version 2.27).

4. DISCUSSION

Bursts that fulfill our criterion $\beta_{\text{OX}} < 0.5$ are classified as dark and are printed in boldface in Table 1. We find 5 certain dark bursts out of a sample of 52 GRBs, consistent with the trend that the dark burst fraction is approaching a level of about 10% (e.g., Lamb et al. 2004). It is clear from Fig. 1 that bursts with no optical counterparts tend to be X-ray faint, as concluded by D03.

As long as a GRB optical and X-ray flux is estimated at the same point in time, the burst can be located in the $F_{\text{opt}}-F_X$ diagram. To the extent that the simple external shock fireball model can be applied,⁵ a burst will either move along constant β_{OX} lines with time (if the optical

⁵ Assuming an unchanged OA spectrum, and that the effect of the reverse shock does not dominate the optical flux (Piran 1999).

TABLE 1
 GAMMA-RAY BURSTS THAT HAVE AN UNAMBIGUOUS DETECTED X-RAY AFTERGLOW AND AN OPTICAL FOLLOW-UP.

GRB	Obs.	β_{OX}	$R(11\text{ h})$	Ref.	GRB	Obs.	β_{OX}	$R(11\text{ h})$	Ref.	GRB	Obs.	β_{OX}	$R(11\text{ h})$	Ref.
970111	SAX	< 0.83	>22.2		990907	SAX	< 0.69	>20.9		020127	CXO	< 1.24	>20.4	(23)
970228	SAX	0.81	19.3	(1)	991014	SAX	< 0.63	>22.4		020322	XMM	0.51	23.3	(24)
970402	SAX	< 0.80	>21.5		991106	SAX	< 0.99	>20.2		020405	CXO	0.75	18.3	(25)
970508	SAX	0.69	21.1	(2)	991216	CXO	0.96	16.9	(10)	020427*	CXO	< 0.87	>19.8	(26)
970828	RXTE	< 0.05	>25.0		000115	RXTE	< 1.06	>15.8	(11)	020813	CXO	0.65	19.1	(27)
971214	SAX	0.64	21.9	(3)	000210	SAX	< 0.54	>23.1	(12)	021004	CXO	0.93	18.4	(28)
971227	SAX	< 0.92	>20.3		000214*	SAX	< 0.92	>19.5	(13)	030226	CXO	0.81	19.5	(29)
980329	SAX	0.54	22.6	(4)	000528	SAX	< 0.69	>22.5		030227	XMM	0.62	21.7	(30)
980519	SAX	1.06	18.8	(5)	000529	SAX	< 1.09	>18.8		030328	CXO	0.80	20.2	(31)
980613	SAX	0.69	22.6	(6)	000615	SAX	< 0.69	>23.1	(14)	030329	RXTE	0.86	14.7	(32)
980703	SAX	0.71	20.1	(7)	000926	SAX	0.87	18.0	(15)	030528*	CXO	0.63	21.1	(33)
981226	SAX	< 0.51	>23.1		001025A	XMM	< 0.43	>24.3	(16)	030723	CXO	1.07	20.9	(34)
990123	SAX	0.65	19.4	(8)	001109*	SAX	< 0.30	>23.2	(17)	031203*	XMM	0.80	21.0	(35)
990506	RXTE	< 0.06	>23.2		010214	SAX	< 0.63	>22.7	(18)	040106	XMM	0.59	21.8	(36)
990510	SAX	0.86	18.1	(9)	010220	XMM	< 0.94	>21.4	(19)	040223*	XMM	< 0.78	>21.4	(37)
990627	SAX	< 1.02	>20.1		010222	SAX	0.64	19.2	(20)	040701	CXO	< 1.17	>18.1	(38)
990704	SAX	< 0.43	>23.4		011030	CXO	< 0.59	>21.7	(21)					
990806	SAX	< 0.51	>23.3		011211	XMM	0.98	20.1	(22)					

REFERENCES. — (1) Galama et al. (2000); (2) Pedersen et al. (1998); (3) Diercks et al. (1998); (4) Reichart et al. (1999); (5) Jaunsen et al. (2001); (6) Hjorth et al. (2002); (7) Holland et al. (2001); (8) Castro-Tirado et al. (1999); (9) Harrison et al. (1999); (10) Halpern et al. (2000); (11) Gorosabel et al. (2000); (12) Piro et al. (2002); (13) Rhoads et al. (2000); (14) Maiorano et al. (2004); (15) Fynbo et al. (2001a); (16) Pedersen et al. (2004); (17) Castro Cerón et al. (2004); (18) Rol et al. (2001); (19) Berger et al. (2001); (20) Stanek et al. (2001); (21) Rhoads et al. (2001); (22) Jakobsson et al. (2003); (23) Lamb et al. (2002); (24) Bloom et al. (2002); (25) Bersier et al. (2003); (26) This work; (27) Urata et al. (2003); (28) Holland et al. (2003); (29) Pandey et al. (2004); (30) Castro-Tirado et al. (2003); (31) Burenin et al. (2003); (32) Lipkin et al. (2004); (33) Rau et al. (2004); (34) Fynbo et al. (2004); (35) Malesani et al. (2004); (36) Masetti et al. (2004); (37) Simoncelli et al. (2004); (38) de Ugarte Postigo et al. (2004)

NOTE. — A burst is marked with an asterisk if the follow-up was not carried out in the R -band, or a deeper limit was available in another band. In these cases, we assumed a spectral index of 0.6 to transform to the R -band. If a burst fulfills our dark burst criteria, i.e., has $\beta_{\text{OX}} < 0.5$, its name is written in boldface. A total of five bursts are classified as dark according to our proposed scheme. The references refer to the optical follow-up; if void they are extrapolated from the R -band magnitudes listed in Fynbo et al. (2001b). The magnitudes have been corrected for Galactic extinction.

and X-ray bands are positioned on the same power-law segment) or along lines with a slightly different slope (if $10^{14}\text{ Hz} < \nu_c < 10^{18}\text{ Hz}$). In the *Swift* era the data will be obtained within the first hour; hence information from the early X-ray light curve or spectrum could be used to estimate p , making it possible to set a limit on β_{OX} for individual bursts (making a universal β_{OX} cutoff unnecessary). However, this relies on instant availability of data and is potentially hampered by, e.g., reverse shocks and light curve fluctuations.

Dark bursts, i.e., bursts located below the line of constant $\beta_{\text{OX}} = 0.5$ in the $F_{\text{opt}}-F_X$ diagram, are guaranteed to be special in the sense that, with respect to the fireball model predictions, they either have a diminished optical flux or an excessive X-ray flux. The former could be due to high redshift or obscuration, while the latter could be caused by X-ray emission lines (e.g., Reeves et al. 2002) or thermal emission. An X-ray faint burst with a low value of $p < 2$ will also be classified as dark in this scheme. It is important to note that, using this definition of dark bursts, there is no assurance that we will catch *all* obscured or high- z bursts. If, for instance, for a particular burst $p = 2.5$ and $\nu_c < 10^{14}\text{ Hz}$, it will have a high intrinsic β_{OX} value and there is no guarantee that high redshift or optical obscuration will shift β_{OX} below 0.5. Moreover, to answer the question why a specific burst is dark it must be modeled in detail; the $F_{\text{opt}}-F_X$ diagram is only a quick diagnostic tool.

We may consider bursts with $0.50 \lesssim \beta_{\text{OX}} \lesssim 0.55$ as

potentially dark. We identify five such bursts, namely GRBs 980329, 981226, 990806, 000210, and 020322. If the value of p is universal (e.g., Waxman 1997), with $p \approx 2.2$, the lower limit on β_{OX} allowed in the fireball model is closer to 0.6. This would shift the aforementioned five bursts into the dark burst category.

The imminent launch of the multi-wavelength observatory *Swift*, expected to detect ~ 100 GRBs/year, offers a unique chance to construct a homogeneous sample with well-understood selection criteria. *Swift* will reach an X-ray limit of $\sim 8\text{ mCrab}$ at 60 s and an optical limit of $R \sim 22\text{ mag}$ at $\sim 300\text{ s}$ (Gehrels et al. 2004). For a *Swift* burst with an X-ray afterglow detected above this flux limit and no detection in the UVOT image, the value of β_{OX} will be below 0.1. This implies that the early (few minutes after the burst) *Swift* data will be adequate to get a rough location of the burst in Fig. 1 and hence to initiate dedicated follow-up observations.

We thank J. Bloom, C. Kouveliotou, D. Lazzati, E. Rol and R. Wijers for discussions on dark GRBs over the years. We thank the anonymous referee for critical reading and useful comments on the paper. PJ and GB gratefully acknowledge support from a special grant from the Icelandic Research Council. KP acknowledges support from the Carlsberg foundation and from the Instrument Center for Danish Astrophysics (IDA). This work was supported by the Danish Natural Science Research Council (SNF). The authors acknowledge benefits from




FIG. 1.— A diagram of optical flux vs. X-ray flux for all bursts in Table 1. Optical fluxes, the corresponding R -band magnitudes shown on the right hand ordinate, and X-ray fluxes have been interpolated/extrapolated to 11 h. The magnitudes have been corrected for Galactic extinction. Filled symbols indicate optical detections while open symbols are upper limits. Lines of constant β_{OX} are shown along with the corresponding value. We define dark bursts as those which have $\beta_{\text{OX}} < 0.5$.

collaboration within the EU FP5 Research Training Net-

work “Gamma-Ray Bursts: An Enigma and a Tool”.

REFERENCES

- Berger, E., et al. 2001, GCN Circ. 958
 Berger, E., et al. 2002, ApJ, 581, 981
 Bersier, D., et al. 2003, ApJ, 583, L63
 Bloom, J. S., et al. 2002, GCN Circ. 1294
 Burenin, R., et al. 2003, GCN Circ. 1990
 Castro Cerón, J. M., et al. 2004, A&A, 424, 833
 Castro-Tirado, A. J., et al. 1999, Science, 283, 2069
 Castro-Tirado, A. J., et al. 2003, A&A, 411, L315
 Crew, G. B., et al. 2003, ApJ, 599, 387
 Dai, Z. G., & Cheng, K. S. 2001, ApJ, 558, L109
 De Pasquale, M., et al. 2003, ApJ, 592, 1018 (D03)
 de Ugarte Postigo, A., et al. 2004, GCN Circ. 2621
 Diercks, A. H., et al. 1998, ApJ, 503, L105
 Djorgovski, S. G., et al. 2001, ApJ, 562, 654
 Fox, D. W., et al. 2003, ApJ, 586, L5
 Fruchter, A., Krolik, J. H., & Rhoads, J. E. 2001, ApJ, 563, 597
 Fynbo, J. P. U., et al. 2001a, A&A, 373, 796
 Fynbo, J. P. U., et al. 2001b, A&A, 369, 373
 Fynbo, J. P. U., et al. 2004, ApJ, 609, 962
 Galama, T., & Wijers, R. A. M. 2001, ApJ, 549, L209
 Galama, T., et al. 2000, ApJ, 536, 185
 Gehrels, N., et al. 2004, ApJ, 611, 1005
 Gorosabel, J., et al. 2000, GCN Circ. 563
 Groot, P. J., et al. 1998, ApJ, 493, L27
 Halpern, J. P., et al. 2000, ApJ, 543, 697
 Harrison, F. A., et al. 1999, ApJ, 523, L121
 Hjorth, J., et al. 2002, ApJ, 576, 113
 Hjorth, J., et al. 2003, ApJ, 597, 699
 Holland, S. T., et al. 2001, A&A, 371, 52
 Holland, S. T., et al. 2003, AJ, 125, 2291
 Jakobsson, P., et al. 2003, A&A, 408, 941
 Jaunsen, A. O., et al. 2001, ApJ, 546, 127
 Lamb, D. Q. 2001, in Gamma-Ray Bursts in the Afterglow Era, ed. E. Costa, F. Frontera, & J. Hjorth (Berlin: Springer), 297
 Lamb, D. Q., & Reichart, D. E. 2000, ApJ, 536, 1
 Lamb, D. Q., et al. 2002, GCN Circ. 1230
 Lamb, D. Q., et al. 2004, NewAR, 48, 423
 Lazzati, D., Covino, S., & Ghisellini, G. 2002, MNRAS, 330, 583
 Lipkin, Y. M., et al. 2004, ApJ, 606, 381
 Li, W., Filippenko, A., Chornock, R., & Jha, S. 2003, ApJ, 586, L9
 Malesani, D., et al. 2004, ApJ, 609, L5
 Masetti, N., et al. 2004, GCN Circ. 2515
 Maiorano, E., et al. 2004, in Gamma-Ray Bursts in the Afterglow Era, ed. M. Feroci, et al. (San Francisco: ASP), 221
 Pandey, S. B., et al. 2003, A&A, 408, L21
 Pandey, S. B., et al. 2004, A&A, 417, 919
 Pedersen, H., et al. 1998, ApJ, 496, 311
 Pedersen, K., et al. 2004, ApJ, submitted
 Perna, R., Lazzati, D., & Fiore, F. 2003, ApJ, 585, 775
 Piran, T. 1999, PhR, 314, 575
 Piro, L. 2004, in Gamma-Ray Bursts in the Afterglow Era, ed. M. Feroci, et al. (San Francisco: ASP), 149
 Piro, L., et al. 2002, ApJ, 577, 680
 Ramirez-Ruiz, E., Trentham, N., & Blain, A. W. 2002, MNRAS, 329, 465
 Rau, A., et al. 2004, A&A, in press (astro-ph/0408210)
 Reeves, J. N., et al. 2002, Nature, 416, 512
 Reichart, D. E., & Price, P. A. 2002, ApJ, 565, 174
 Reichart, D. E., et al. 1999, ApJ, 517, 692
 Rhoads, J., et al. 2000, GCN Circ. 564
 Rhoads, J. E., et al. 2001, GCN Circ. 1140
 Rol, E. 2004, PhD Thesis
 Rol, E., et al. 2001, GCN Circ. 955
 Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
 Simoncelli, A., et al. 2004, GCN Circ. 2549
 Stanek, K. Z., et al. 2001, ApJ, 563, 592
 Stratta, G., et al. 2004, ApJ, 608, 846
 Taylor, G. B., et al. 1998, ApJ, 502, L115
 Taylor, G. B., et al. 2000, ApJ, 537, L17
 Totani, T. 1997, ApJ, 486, L71
 Urata, Y., et al. 2003, ApJ, 595, L21
 Vreeswijk, P. M., et al. 2004, A&A, 419, 927
 Waxman, E. 1997, ApJ, 485, L5
 Waxman, E., & Draine, B. T. 2000, ApJ, 537, 796
 Wijers, R. A. M. J., Bloom, J. S., Bagla, J. S., & Natarajan, P. 1998, MNRAS, 294, L13