

Citation for published version:

Herbert Talwana, et al, 'Agricultural nematology in East and Southern Africa: problems, management strategies and stakeholder linkages', *Pest Management Science* Vol. 72 (2): 226-245, February 2016.

DOI:

<http://dx.doi.org/10.1002/ps.4104>

Document Version:

This is the Accepted Manuscript version.

The version in the University of Hertfordshire Research Archive may differ from the final published version. **Users should always cite the published version of record.**

Copyright and Reuse:

© 2015 Society of Chemical Industry.

This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

Enquiries

If you believe this document infringes copyright, please contact the Research & Scholarly Communications Team at rsc@herts.ac.uk

1 **AGRICULTURAL NEMATOLOGY IN EAST AND SOUTHERN**
2 **AFRICA: PROBLEMS, MANAGEMENT STRATEGIES AND**
3 **STAKEHOLDER LINKAGES**

4		
5	Herbert Talwana ¹	haltalwana@agric.mak.ac.ug
6	Zibusiso Sibanda ²	sibandazibusiso@yahoo.com
7	Waceke Wanjohi ³	wanjohi.waceke@ku.ac.ke
8	Wangai Kimenju ⁴	wkimenju@yahoo.com
9	Nessie Luambano-Nyoni ⁵	mluambano@yahoo.com
10	Cornel Massawe ⁶	massawesa@yahoo.co.uk
11	Keith G Davies ^{7, 12}	k.davies@herts.ac.uk
12	Rosa H Manzanilla-López ^{7,*}	rosa.manzanilla@gmail.com
13	David J Hunt ⁸	d.hunt@cabi.org
14	Richard A Sikora ⁹	ulp40@uni-bonn.de
15	Danny L Coyne ¹⁰	d.coyne@cgair.org
16	Simon R Gowen ¹¹	s.r.gowen@reading.ac.uk
17	Brian R Kerry ⁷	[deceased]

18
19 ¹ Department of Crop Science, Faculty of Agriculture, Makerere University, P.O. Box 7062,
20 Uganda

21 ² Goldengro Pvt Ltd, 26 Langbourne Avenue, Harare, Zimbabwe

22 ³ Agricultural Science and Technology, Kenyatta University, P.O. Box 43844-00100 Nairobi,
23 Kenya.

24 ⁴ Faculty of Agriculture, University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya

25 ⁵ Sugarcane Research Institute, Kibaha, Tanzania

- 1 ⁶ Tengeru Horticultural Research Institute, Arusha, Dar es Salaam, Tanzania
- 2 ⁷ Department of AgroEcology, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ,
- 3 UK
- 4 ⁸ CABI Europe-UK, Bakeham Lane, Egham, Surrey, TW20 9TY, UK
- 5 ⁹ INRES-Phytomedizin, Nassallee 9, 53115 Bonn, Germany
- 6 ¹⁰ IITA-Tanzania Plot 331, Regional Hub, CocaCola Road, Mikocheni A. P.O. Box 34441,
- 7 Dar es Salaam, Tanzania
- 8 ¹¹ School of Agriculture and Policy Development, University of Reading, Reading,
- 9 Berkshire, UK
- 10 ¹² School of Life and Medical Sciences, University of Hertfordshire, College Lane, Hatfield,
- 11 LA10 9AB, UK
- 12 * Corresponding author.
- 13

Abstract

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

BACKGROUND: By 2050, Africa’s population is projected to hit two billion. Africa will have to increase food production more than 50% in the coming 50 years to meet the nutritional requirements of its growing population. Nowhere is the need to increase agricultural productivity more pertinent than in much of sub-Saharan Africa where it is currently static or declining. Optimal pest management will be essential as intensification of any system creates heightened selection pressures for pests.

RESULTS: Plant-parasitic nematodes and their damage potential are intertwined with intensified systems and can be an indicator of unsustainable practices. As soil pests, nematode are commonly overlooked or misdiagnosed, particularly where appropriate expertise and knowledge transfer systems are meager or inadequately funded. Nematode damage to roots produces symptoms similar to nutrient deficiency, often leading to a misdiagnosis and consequent overuse of fertilizers. Damage in subsistence agriculture is exacerbated by growing crops on degraded soils and in areas of low water retention where strong root growth is vital.

CONCLUSION: This review focuses on the current knowledge of economically important nematode pests affecting key crops, control methods and the research and development needs for sustainable management, stakeholder involvement and capacity building in the context of crop security in Kenya, Tanzania, Uganda and Zimbabwe.

Key words: Africa, agricultural nematology, capacity building, development needs, food security, pest management.

1 **1. INTRODUCTION**

2

3 Nematodes, either alone or in combination with other pathogens, constitute an important
4 constraint to world food production.¹ Although not all plant-parasitic nematodes (PPN) are of
5 economic importance, nematode damage to agricultural crops is estimated globally to cause
6 losses in the range of 8.4-20.6% in life sustaining and economically important/commodity
7 crops (Table 1). Although precise data are difficult to calculate, the value of these losses is
8 estimated to exceed US \$125 billion.^{2,3} Amongst the most widespread and economically
9 important nematodes are species of the genera *Helicotylenchus*, *Meloidogyne*, *Pratylenchus*,
10 *Radopholus* and *Scutellonema*, all of which have wide host ranges and are capable of
11 attacking many different crops (see Table 2). In addition, a small number of species of some
12 genera can act as virus vectors (*e.g.*, species in the genera *Longidorus*, *Paralongidorus*,
13 *Trichodorus* and *Xiphinema*), and some are post-harvest pests, including *Meloidogyne*,
14 *Pratylenchus sudanensis* Loof & Yassin, 1971 and *Scutellonema bradys* (Steiner & Le Hew,
15 1933) Andrassy, 1958 on stored yam or potato,^{4,5} thereby reducing the longevity and
16 marketability of the harvested crop.⁶ Nematode infection is also known to compromise plant
17 resistance to other soil-borne plant pathogens. In addition to disruption of the usual root
18 functions, processes such as nitrogen fixation in the nodules of leguminous plants is
19 suppressed by species of *Meloidogyne*, the so-called root-knot nematodes (RKN),^{7,8} even
20 when there is no visual reduction in nodulation.⁹ Reduction in root depth extension caused by
21 nematodes also increases crop susceptibility to water stress leading to wilting and yield loss.

22

23 The damage caused by PPN, their behavior and control have received little attention in many
24 countries in sub-Saharan Africa (SSA), despite indications that crop productivity can be
25 severely reduced.^{10,11} Information concerning PPN problems in much of Africa is all too

1 scarce, often being restricted to gray literature in institute reports rather than globally
2 disseminated and accurate crop loss damage is lacking for many crops and cropping systems,
3 particularly those practised by smallholders. This lack is of particular concern as it is
4 predicted that tropical and subtropical nematology will face ever more complex and
5 economically important problems, especially in subsistence agriculture¹, such problems being
6 exacerbated by the effects of climate change. There is therefore an urgent need in SSA for
7 nematode surveys, accurate pest identification, the establishment of distribution maps, and
8 the gathering of accurate crop damage and yield loss data, not only for commodity crops, but
9 also for staples. Such information will underpin the requirement from agricultural research
10 and development programmers to investigate nematode problems and to develop
11 management systems for sustainable cropping systems. With this in mind it is highly
12 desirable to exploit the good farming practices, including conservation agriculture and
13 integrated crop and pest management, that form a part of ecosystem services and which have
14 been proven to provide optimum production potential, input efficiency, economic benefit and
15 environmental protection. Implementing good practices and applying integrated pest
16 management (IPM) at the smallholder level will benefit the rural poor, increase agricultural
17 productivity and enable farmers to grow more food, thereby leading to better diets and
18 potentially higher farm incomes. With income derived from more efficient and sustainable
19 crop management, farmers are more likely to diversify production and grow higher-value
20 crops, benefiting not only themselves but also the broader economy.¹²

21

22 This review stems from a Gatsby Charitable Foundation initiative aimed at capacity building
23 in agricultural nematology in certain countries in East and Southern Africa (ESA). With
24 particular emphasis being placed on the status quo in Kenya, Tanzania, Uganda and
25 Zimbabwe, the review endeavors to serve as a ‘call to action’ by summarizing the currently

1 available knowledge of: *i*) the economically important nematode problems affecting key
2 crops; *ii*) nematode control/management methods; and *iii*) highlighting the research and
3 development needs for future management, soil health, stakeholder involvement and capacity
4 building in the context of crop security.

5

6 **1.1 Common cropping systems in East and Southern Africa**

7

8 The countries within ESA, as defined above, possibly have the greatest concentration of
9 poverty in the world and, for the foreseeable future, will continue to struggle to overcome
10 widespread food insecurity and malnourishment.¹³ Population growth, drought and degraded
11 soils coupled with the long term impact of climate change, are new and disturbing challenges
12 that are likely to lead to a further reduction in crop security and thereby to increased poverty.
13 Predicted increase in extreme weather patterns, particularly heavy rains and drought, will be
14 detrimental to farmers, their crops, and the rural economy. Although there are many examples
15 of good agricultural practice, the vast majority of the rural poor in ESA are smallholder
16 farmers working in conditions of either static or declining productivity. Agriculture accounts
17 for 60% of the region's GDP, employs 70% of the total labor force and is its main livelihood.
18 These resource poor farmers produce the vast majority of food on the continent and they need
19 to increase production 1-2% per year to offset growth in populations both rural and in the
20 enlarging mega-cities. Present food production in SSA will only be able to meet 13% of the
21 continent's food needs by 2050.^{14,15}

22

23 The major farming systems in ESA are largely determined by the rainfall pattern (total annual
24 amount and distribution). Farming systems cover a wide range of activities, including the
25 production of cash and food crops and livestock. Eleven broad farming systems have been

1 identified² and these share many similarities (Table 3). In each system, production is typically
2 done by subsistence smallholder producers, many of whom are women, using relatively
3 rudimentary technology. However, at the other end of the spectrum there are also
4 industrial/export crops that are grown on a much larger scale with a commensurate increase
5 in inputs and technology. As 40% or more of the soils in SSA are degraded, changes in
6 cropping systems and support for the challenges faced by small farmers are urgently needed.

7

8 Increasing urbanization within the region has produced a more commercial attitude to
9 horticultural enterprises such as export vegetables, fruits, spices, and cut flowers. For
10 example, Kenya, South Africa, Zimbabwe, Zambia, Ethiopia and Uganda have all witnessed
11 the creation of new horticulture industries.¹⁶⁻¹⁸ These operate in open fields and screen-
12 houses, with extensive irrigation and use of pesticides and chemical fertilizers. Upland
13 plateaux are considered the best location for horticulture, hence the selective developments in
14 Kenya, Tanzania, Zambia, Zimbabwe and other countries such as Ethiopia, although lower
15 altitude areas with high humidity and warmer nights can also be suitable, as in Uganda. The
16 horticultural sector is growing rapidly, underscoring its increasing rank as a major, non-
17 traditional, foreign exchange earner.¹⁶⁻¹⁸

18

19 **1.2 Nature of nematode problems and sustainability of cropping systems in East and** 20 **Southern Africa**

21

22 The traditional farming systems in Africa usually consisted of mixed crops (even including
23 trees or woody perennials) that provided an ecosystem in which farmers manipulated and
24 derived advantage from local resources and natural processes. This required the application
25 of agro-ecology principles to manage soil fertility, pests, diseases, weeds and available

1 genetic resources in the context of climate and land use. However, the rapidly increasing
2 human population and increased competition for land has led to the abandonment of
3 traditional farming systems in favor of larger, more intensive and specialized systems
4 characterized by monocultures with new varieties, crop protection aids and inorganic
5 fertilizers. The intensification of agricultural land use is often associated with natural habitat
6 decline at farm and landscape level. This landscape simplification has increasingly negative
7 effects, including a long-term decrease in sustainability of cropping systems due to nutrient
8 extract agricultural practices.

9
10 Crop losses due to pests and diseases appear to be more frequent in simplified landscapes and
11 decrease in sustainability is indicated by increased incidence and higher populations of PPN
12 leading to elevated damage levels. By contrast, in more diverse agro-ecosystems where soils
13 are populated by a broader range of microorganisms, including bacterial and fungal feeding
14 nematodes (important to convert nutrients to forms accessible to plants), PPN can be
15 suppressed to densities below the economic threshold.¹⁹

16
17 PPN are among the major constraints to improvement in crop productivity, attainment of food
18 security and poverty alleviation in SSA. Rarely is any crop free from nematode attack in
19 smallholder farmer fields, orchards and home gardens, yet their presence is generally not
20 recognized due to the cryptic nature of the perpetrators. Their importance in SSA is relatively
21 higher due to a number of factors operating singly or interactively. For example, shorter
22 nematode life-cycles in warmer soils and continuous crop growing seasons result in more
23 rapid population build-up, and negatively impact upon crops and antagonistic
24 microorganisms. In most cases, crop roots are simultaneously parasitized by several
25 economically important nematode species (Table 2), yield losses being estimated from 8-50%

1 (Table 1). Component species of the mixed nematode population often differ from crop to
2 crop, country to country and region to region, making diagnosis and prediction of damage
3 much more difficult and thereby complicating the development of management strategies.
4 Interestingly, one of the major influences on the distribution of nematodes appears to be
5 altitude, several studies having investigated its role in ESA (Table 4). Such complexity also
6 impacts upon the advice given to growers by the extension services.

7

8 **2 NEMATODE PROBLEMS ON KEY STAPLE AND ECONOMICALLY** 9 **IMPORTANT CROPS**

10

11 The majority of nematology research in ESA has, perhaps unsurprisingly, mainly focused on
12 high value crops destined for export (*e.g.*, cotton, maize, sugar cane and tobacco). Research
13 on the staple food crops of an area and awareness of nematode problems in the local farming
14 communities is limited. However, the limited data available in ESA (addressing this
15 limitation is one of the major keys to ameliorating the problem) indicates that PPN are
16 widespread and cause significant losses in bananas,²⁰⁻²⁴ maize,^{25,26} yam,²⁷⁻³⁰ and vegetables.³¹⁻
17 ³⁰ The examples and discussions presented here have been selected to demonstrate that
18 nematodes are frequently a limiting factor on a wide range of crops commonly grown in
19 ESA, particular focus being given to Kenya, Tanzania, Uganda and Zimbabwe.

20

21 **2.1 Cereal crops**

22

23 Cereals constitute ESA's most important source of food and include maize (*Zea mays* L.),
24 sorghum (*Sorghum vulgare* L.) and rice (*Oryza sativa* L.), maize being the most important of
25 these crops on a regional basis.^{1,10}

1

2 Rice has gained in popularity and is grown as a rain-fed lowland and upland crop in much of
3 Uganda and as an irrigated crop in Kenya and Tanzania.¹ Sorghum, a traditional African
4 cereal crop, is drought-tolerant and regularly out-yields maize in semi-arid parts of the region
5 where rainfall is both insufficient and unreliable.¹ It is also more resistant to water-logging,
6 yields well in infertile soils, and can be ratooned. *Meloidogyne acronea* Coetzee, 1956 causes
7 damage on sorghum in Malawi and South Africa.¹⁰

8

9 Maize occupies the pre-eminent position in terms of production, acreage and source of
10 nutrition and is a major staple in many rural and urban communities.^{1,10} Maize has also
11 recently been identified as one of the non-traditional cash crops – traded regionally and
12 purchased by relief organizations for distribution to internally displaced persons. Production
13 of crops such as maize and other cereals will be the most effective in increasing food security
14 as they act as a source of cash for poorer households and benefit farmers indirectly by
15 allowing them to diversify into higher-value crops, thus improving their incomes and
16 reducing urban poverty.

17

18 PPN constitute serious impediments to intensified cereal production in SSA and several
19 species have been associated with cereals in Uganda²⁶ and Kenya.^{32,33} So far, pathogenicity
20 of *Pratylenchus zae* Graham, 1951 and *Meloidogyne* spp. on maize has been
21 demonstrated.^{25,34} These nematodes occur very frequently and abundantly in maize roots and
22 their rhizosphere.^{25,26,35} Analysis of the relationship between nematode densities, soil
23 physical/chemical properties and cropping history revealed that intensified cereal systems
24 with low frequency of non-cereal rotations increased the risk of *P. zae* infestation, especially
25 if the soils were loamy sand, sandy loam or sandy clay loam,²⁶ and could be associated with

1 yield losses up to 37%.³⁴ Symptoms, especially those above ground, are not specific and are
2 characterized by irregular patches of stunted plants randomly distributed throughout the
3 fields. Reduced root/shoot growth, leaf necrosis, proliferation of fibrous roots, and the
4 presence of small blackish root lesions also occur.³⁵ Other important cereal pests are RKN,
5 particularly on maize and rice. RKN are regularly found in combination with *P. zaeae*, the
6 latter apparently being more detrimental to crop productivity.³⁵

7

8 **2.2 Bananas and plantains**

9

10 Bananas and plantains (*Musa* spp.) have received great attention regarding PPN (Table 2) and
11 their control. Starchy bananas and plantains are often grown by small-scale farmers²³ and are
12 of great importance in the highlands of countries such as Burundi, Democratic Republic of
13 Congo (Kivu region), Rwanda, Tanzania (Kagera region) and western Kenya and Uganda,
14 where they are an important staple food and cash crop.²⁴ However, banana yields in the
15 highlands of Kenya and Uganda have shown a steady decline per unit land area due to a
16 range of factors that include infection and damage by a complex of PPN³⁶.

17

18 Plant-parasitic nematodes associated with damage and reduced yields in banana and plantain
19 include: *Helicotylenchus* spp., *Meloidogyne* spp., *Pratylenchus coffeae* (Zimmermann, 1898)
20 Filipjev & Schuurmans Stekhoven, 1941, *P. goodeyi* Sher & Allen, 1953, *Radopholus similis*
21 (Cobb, 1893) Thorne, 1949 and *Rotylenchulus reniformis* Linford & Oliveira, 1940 (Tables 1
22 and 2). Heavy infestations of root nematodes have been associated with toppling,³⁵ a problem
23 initiated by PPN where root lesions become so extensive that secondary fungal infections
24 cause the roots to rot, thereby reducing nutrient and water uptake and compromising
25 anchorage of the plant in the soil. This affects the growing banana plant, especially those in

1 bunch, to such an extent that they topple, the developing fruit being lost, reduced in value or
2 fit only for livestock consumption.

3

4 Although there is little work on interrelationships between nematodes and other organisms,
5 one example is that between *P. goodeyi* and a non-pathogenic strain of *Fusarium oxysporum*
6 in a root-rotting complex of highland bananas in Kenya.³⁷⁻³⁹ Nematodes are found on all
7 varieties of bananas, but there is some evidence to suggest that there are differences in
8 susceptibility among the most commonly grown varieties, two widely known and confirmed
9 sources of resistance to *R. similis* being: 'Pisang Jari Buaya' and 'Yangambi km5'.⁴⁰⁻⁴⁷

10 Breeding resistance into local varieties, supported by other management strategies such as
11 heavy application of organic matter ('trash') can result in substantial yield responses with
12 little financial input from the smallholders.⁴⁸⁻⁵¹

13

14 **2.3 Legumes**

15

16 Legumes are important crops as they have the potential to fix nitrogen and therefore improve
17 the fertility of the soil. They are also an important source of nutrition in that they are a vital
18 source of protein.

19

20 The common bean (*Phaseolus vulgaris* L.) has a centre of origin and genetic diversity in the
21 highland regions of Africa and has become a crop of major importance in subsistence
22 agriculture and food security in the countries of the highlands of ESA.⁵² The most serious
23 nematode pests include the RKN species *M. incognita* and *M. javanica* and losses up to 60%
24 have been attributed to RKN in experimental work done in Kenya.^{8,53}

25

1 The cultivated groundnut (*Arachis hypogaea* L.) is a legume originating in South America
2 although it is now widely grown throughout Africa and is an important crop rich in both
3 carbohydrate and protein. PPN are primary pests in all groundnut producing areas with
4 estimated losses of 12%.² Important nematode parasites of groundnuts in Africa include
5 RKN, although the crop is also host to the lesion nematode *Pratylenchus brachyurus*
6 (Godfrey, 1929) Filipjev & Schuurmans Stekhoven, 1941, the spiral nematode *Scutellonema*
7 *cavenessi* Sher, 1964, and the stem nematode *Ditylenchus africanus* Wendt, Swart, Vrain &
8 Webster, 1995, which affect productivity and may reduce marketability (Table 2).
9 *Aphelenchoides arachidis* Bos, 1977, originally reported and described damaging groundnut
10 pods and seeds in Nigeria, is now also known from South Africa and, if not already present in
11 other countries of ESA, certainly represents a quarantine risk.

12

13 **2.4 Root and Tuber Crops**

14

15 *2.4.1 Cassava and yam*

16

17 Cassava (*Manihot esculenta* Crantz) and yam (*Dioscorea* spp.) are very important staple food
18 crops and provide food security and income for millions of smallholders⁵⁴ – in some African
19 countries, cassava constitutes 80% of the *per capita* consumption of starchy staples.⁶

20

21 Cassava is often viewed as being resistant to, or uninfected by, nematode pests. However,
22 although it is host to a wide range of nematodes, RKN and *P. brachyurus* are the most
23 important pests, causing consistent losses.^{27,55} Damage appears to be more severe when
24 infection occurs on young plants.⁵⁶ The damage caused to cassava depends on nematode
25 isolate (race) and cassava cultivar.^{10,57} For example, high yielding elite varieties were

1 introduced to Mozambique from Nigeria (where they had fared well) yet became heavily
2 damaged by RKN and were consequently unsuitable for local production (Coyne DL, pers.
3 obs.).

4
5 Yams (*Dioscorea* spp.) are a starch staple crop across Africa. The major nematode pest, *S.*
6 *bradys*, which occurs mostly in West Africa, affects the tubers and causes ‘dry rot disease’.
7 Infected tissues discolor, initially becoming light brown before eventually turning black, the
8 most severe symptoms occurring during storage.²⁷ Dry rot causes a marked reduction in
9 quality and marketable value of the tubers while infected tubers suffer major deterioration in
10 storage. In Uganda, *P. sudanensis* has been observed causing similar symptoms to *S. bradys*,
11 while RKN are particularly prevalent and damaging, causing rotting and necrosis, especially
12 during storage.^{4,28} Nematode-damaged tubers used as planting material also re-infect the
13 subsequent crop and so perpetuate the problem.^{27,30}

15 2.2.2 Potato

16
17 Potato (*Solanum tuberosum* L.) is traditionally grown in cool environments, although in
18 recent years it has expanded its range in Africa to relatively warmer, more humid zones, the
19 latter being optimal for many pests, including nematodes. In cooler areas, where the potato
20 cyst nematode (*Globodera* spp.; PCN) has been introduced, it is a major problem and is
21 currently acknowledged by EPPO (European and Mediterranean Plant Protection
22 Organization) and NAPPO (North American Plant Protection Organization) as a potentially
23 important quarantine pest (www.eppo.int/QUARANTINE/listA2.htm;
24 http://www.pestalert.org/opr_search.cfm). RKN attacks the roots and tubers and is a major
25 constraint to both productivity and marketability of potatoes.⁵⁸ It also reduces seed quality

1 and, where infected material is used for seed, is easily spread from one field or country to
2 another.

3

4 Another quarantine species, *Ditylenchus destructor* Thorne, 1945, also known as the Potato
5 Rot Nematode, has been associated with potatoes in Kenya.¹⁰ It causes internal rotting to the
6 tubers and thereby reduces yield and marketability of the crop. It is easily dispersed by using
7 infested planting material.

8

9 Given the increasing importance of the crop, and with the exception of South Africa, there is
10 a serious lack of knowledge on nematode pests in many of the African potato cropping areas.

11

12 *2.2.3 Sweet Potato*

13

14 Sweet potato (*Ipomoea batatas* (L.) Lam) is host to a wide range of PPN species, although
15 only a few are of economic importance. *Rotylenchulus reniformis*, RKN and *Pratylenchus*
16 spp. infect sweet potato in mixed cropping systems and can result in severely distorted and
17 cracked tubers.^{59,60} Little is known about the nematode pests of sweet potato in the countries
18 of ESA.

19

20 **2.5 Cash Crops**

21

22 *2.5.1 Coffee*

23

24 In many coffee-producing countries in ESA, PPN reduce productivity and increase
25 production costs. *Meloidogyne* spp. and *Pratylenchus* spp. are the most economically

1 important nematodes (Tables 1 and 2) causing estimated yield losses of at least 15% and
2 affecting many plantations. A number of *Meloidogyne* species occur on coffee in ESA
3 including *M. incognita* (Kofoid & White, 1919) Chitwood, 1949, *M. africana* Whitehead,
4 1959 from Kenya (also found in Zaire), *M. decalineata* Whitehead, 1968 from Tanzania (also
5 found in Sao Tomé),¹⁰ and recently *M. paranaensis* Carneiro, Carneiro, Abrantes, Santos &
6 Almeida, 1996 in Uganda (Carneiro R, pers. obs.), and *M. hispanica* Hirschmann, 1986 and a
7 non-described species from the Tanga Region of Tanzania.⁶¹ At least two endemic
8 *Meloidogyne* species occur in ESA: *M. africana* Whitehead, 1959 from Kenya (also found in
9 Zaire), and *M. decalineata* Whitehead, 1968 from Tanzania (also found in Sao Tomé).¹⁰
10 There are other non-described species of *Meloidogyne* from northern Tanzania which infest
11 trees in both monocropped plantations and in smallholder farms where coffee is grown as a
12 mixed crop.¹⁰ Little information is available on the biology and yield losses caused by
13 African coffee RKN and the taxonomy and identification of these pests is both complex and
14 difficult, often requiring relatively sophisticated molecular methodologies for reliable species
15 diagnostics. As a result, appropriate management options for coffee need to be developed,
16 necessitating interdisciplinary research to: *i*) evaluate sampling strategies for assessment of
17 field populations and epidemiology; *ii*) assess their effect on productivity and resistance of
18 coffee genotypes (arabica and robusta); *iii*) evaluate cultural and chemical management
19 strategies; and *iv*) investigate parasite physiology, behavior and practical diagnostics.

20

21 2.5.2 Cotton

22

23 Within semi-arid subsistence agricultural systems cotton is often planted to generate cash and
24 can be grown as an insurance crop in regions where unreliable rain may cause staple food
25 crops to fail. Cotton is Tanzania's larger export crop after coffee, contributing \$90 million to

1 export earnings⁶² (<http://www.worldbank.org/af/wps/wp42.pdf>; <http://www.cotton.or.tz/>).

2 RKN are among the most important nematode pests of cotton, *M. incognita* being the most

3 widely distributed species and causing severe yield losses. It has been found in Tanzania,

4 Uganda and Zimbabwe.⁶³ Infected plants are more prone to Fusarium wilt in adverse

5 conditions, such as drought and high temperature. The *M. incognita* cotton isolate that occurs

6 in Tanzania is a serious pest and is related to an increased incidence of Fusarium wilt caused

7 by *Fusarium oxysporum* f.sp. *vasinfectum* (G.F. Atk.) W.C. Snyder & H.N. Hansen; it has

8 also been shown to break Fusarium resistance in *F. oxysporum* resistant lines.^{5,64} In addition,

9 the nematode facilitates dissemination of the wilt fungus in infected seed.^{10,65} The ‘African

10 cotton root nematode’, *M. acronea*, originally described from Malawi, does not produce galls

11 but causes an increase in fine lateral root growth around the feeding site.¹⁰ In Tanzania, *R.*

12 *reniformis*, the ‘reniform nematode’, produces some field symptoms that resemble those of

13 RKN. Although not producing root galling, it reduces root growth, especially on

14 seedlings.^{63,66} There are many other species of PPN associated with cotton including

15 *Scutellonema aberrans* (Whitehead, 1960) Sher, 1961, *S. brachyurus*, *S. clathricaudatum*

16 Whitehead, 1959, *S. magniphasmum* Sher, 1964 and *S. unum* Sher, 1964 in Tanzania.^{10,63,65}

17

18 2.5.3 Sugar Cane

19

20 Sugar cane is a plantation crop but is also planted by resource-poor farmers. Because it is a

21 plantation crop and of global importance it has been the focus of much research.¹ The number

22 of species of nematode affecting sugarcane is greater than for any other crop.⁶⁷ This is in part

23 due to its continuous monoculture with no more than a few months break between removal of

24 the old ratoon crop and replanting, but is also related to its extensive mat of surface roots

25 which is conducive to the build-up of the root lesion nematode (*Pratylenchus* spp.) of which

1 *P. zae* is reported to be the most common in Kenya.^{68,69} RKN are also of importance on
2 sugar cane throughout ESA, *M. incognita* and *M. javanica* (Treub, 1885) Chitwood, 1949
3 being recorded as the most prevalent.

4

5 *2.5.4 Tobacco*

6

7 Tobacco, although not a food crop is an economically important crop in many parts of Africa,
8 including the countries of ESA, particularly in Zimbabwe. There is a large and substantial
9 literature on tobacco nematodes⁷⁰ as PPN occur wherever it is grown, although the extent of
10 the problem is influenced by climate and soil type. The most important PPN on tobacco in
11 Africa are RKN and lesion nematodes (*Pratylenchus* spp.), although there are also other
12 species which can be problematic. Tobacco also exhibits a considerable degree of resistance
13 to different populations of nematodes⁷⁰ and has recently become a transgenic model plant to
14 investigate the role of various genes that may play a role in nematode resistance.⁷¹

15

16 **2.6 Vegetable Crops**

17

18 Vegetables are an important component of human nutrition and with increasing urbanization
19 they therefore play an increasing role in periurban cropping systems, thereby requiring the
20 development of intensive cropping systems with a drive towards more efficient/commercial
21 production systems. The lack of knowledge of PPN and their behavior in these new systems
22 remains a major challenge to vegetable production in ESA. Such intensive cropping will
23 doubtless result in a buildup of PPN and consequent damage and reduced yield.^{72,73} However,
24 despite high losses, PPN associated with vegetable production are often overlooked and
25 misdiagnosed with many nematode pests remaining unknown.^{11,31} As a result, many farmers

1 are unaware of PPN as pests,¹¹ even though successful production of vegetables is reliant on
2 appropriate nematode management, this being especially necessary in intensified periurban
3 systems where RKN, if left unchecked, can devastate the crop.⁷⁴ Although not all
4 smallholders have recognized nematodes as a biological constraint, up to 20% of vegetable
5 producers use nematicides when growing tomatoes.^{73,75}

6

7 Root-knot nematodes are a concern to many smallholders and commercial producers.^{72,73} An
8 estimate of 20% loss has been attributed to RKN in Kenya, but losses may be up to 50% and
9 total crop failure, principally due to *M. incognita* and *M. javanica*, is not uncommon.⁷⁶ Other
10 high yield losses reported in the region include 50% in beans; 38% in spinach and 32% in
11 okra.⁷⁷

12

13 RKN have been identified among the major pests of cabbage (*Brassica oleracea* var. *capitata*
14 L.),⁷⁷ and are the main pests of tomato (*Solanum lycopersicum* L.), causing considerable yield
15 losses.^{31,78} *Meloidogyne incognita* is often the most commonly identified pest, although given
16 the complexity of RKN diagnostics, it seems likely that a range of morphologically similar
17 species is present.⁷⁴ Correct species identification is complex and with around 100 named
18 species to date, there are probably still many more waiting to be described^{79,80}. In a recent
19 survey carried out in vegetable fields (i.e., tomato, green pepper, cassava, sweet pepper, okra
20 and carrot) in Tanzania and Uganda, species such as *M. arenaria*, *M. enterolobii*, *M. hapla*,
21 *M. javanica*, and *M. incognita* were biochemically identified by esterase patterns and SCAR
22 markers.⁷⁹ Rapid and accurate diagnostics, such as those offered by appropriate molecular
23 tools, may be vitally important when determining which resistant cultivars to plant and also
24 when recommending crop rotations aimed at reducing nematode populations, many
25 indigenous leafy vegetables, for example, also being susceptible to attack.⁸¹

1

2 *Pratylenchus* spp. (lesion nematodes) are known for their impact on crop yields globally. In
3 Kenya, *Pratylenchus* has been identified as a threat to vegetable production.⁷⁷ This lesion
4 nematode occurs at very high population densities in the Central, Eastern and Rift Valley
5 Provinces of Kenya. Among the species that have been found associated with vegetables are
6 *P. brachyurus*, *P. loosi* Loof, 1960, *P. neglectus* (Rensch, 1924) Filipjev & Schuurmans
7 Stekhoven 1941, *P. scribneri* Steiner, 1943 and *P. zaeae*.⁷⁷ In Uganda, lesion nematodes have
8 been reported in carrot, cabbage, pepper, tomato, cucumber and okra.³¹ The presence of
9 lesion nematodes at high population densities is a matter for substantial concern considering
10 their migratory endoparasitic mode of feeding and ability to attack a wide variety of field
11 crops.

12

13 *Helicotylenchus* spp. (spiral nematodes) are also frequently present, often occurring in large
14 numbers and as several different species. These nematodes also have a wide host range and
15 have been found on various vegetables, including cabbage, pepper, tomato, carrot and
16 cabbage in both Kenya and Uganda.^{31,77} The nematode partially burrows into the roots of
17 vegetable crops and feeds within the cortex, leading to brown necrotic areas on attacked
18 roots.

19

20 The reniform nematode, *Rotylenchulus* spp., has consistently been associated with beans,
21 pepper, tomato, okra and cabbage in both Kenya and Uganda,^{31,77} although opinions vary as
22 to its overall importance in yield reduction.

23

24 Other relevant species in fruit and vegetable crops include *Pratylenchus vulnus* Allen &
25 Jensen, 1951 in strawberries in the Central province of Kenya.¹⁰ Other genera reported

1 associated with many of vegetable crops in Kenya include *Belonolaimus*, *Hemicycliophora*,
2 *Hoplolaimus*, *Longidorus*, *Paratrichodorus*, *Trichodorus*, *Tylenchorhynchus*, *Scutellonema*,
3 and *Xiphinema*. In Uganda, various nematodes have been found associated with vegetables,
4 among them *Criconemoides*, *Hemicycliophora*, *Hoplolaimus*, *Quinisulcius*, *Scutellonema* and
5 *Xiphinema*. *Tylenchus*, *Filenchus* and *Coslenchus* are also abundant, although they do not
6 appear to have any significant economic importance as they tend to be fungal feeders or
7 epidermal root browsers.³¹

8

9 **2.7 Ornamentals**

10

11 The flower industry in ESA countries utilizes land that previously grew agricultural crops.
12 Consequently, many nematode problems affecting these crops have adapted to ornamental
13 plants, often causing severe damage. In Uganda, the screen-houses for ornamental plants
14 were erected on land previously cropped with bananas and there have been several nematode
15 problems on carnations and roses. Because the industry is export driven and there are strict
16 limitations and international restrictions on which pesticides can be used (*i.e.*, GlobalGAP
17 guideline observation), nematodes are becoming an increasingly major problem. There are
18 few, if any, nematicides on the market that can be used and they are extremely expensive.
19 The foliar chrysanthemum nematodes *Aphelenchoides* spp., the bulb nematodes *Ditylenchus*
20 spp. and RKN have been reported from ornamental plants in ESA (Gowen SR,
21 unpublished).¹⁰ All three genera are commonly associated with perennial herbaceous
22 ornamentals, a rapidly expanding segment of the floriculture industry in the region.
23 Vegetative propagation of many of these plants may also result in increased spread and
24 distribution of PPN. Symptoms of nematode infection may range from virtually none to poor
25 growth, stunting, wilting, nutrient deficiency, or even death when in combination with

1 secondary organisms. For example, the fungus *Botryotinia fuckeliana* (de Bary) Whetzel
2 1945 (= *Botrytis cinerea*) commonly infects leaves previously infected with foliar nematodes,
3 thereby masking nematode induced symptoms and increasing plant mortality.

4
5 With the increasing importance of nematodes in the ornamental industry, diagnosis of
6 nematodes as the cause of disease is an important step. For example, in Kenya, carnation
7 (*Dianthus caryophyllus* L.) is estimated to be grown on more than 500 ha, mainly under
8 greenhouse conditions, due to expanding market demand.^{17,18} The very major challenge lies
9 in educating growers involved in the flower industry about the existence of nematode
10 problems and the need to do a nematological survey of land previously used for a different
11 crop prior to establishment of ornamentals. However, this situation also provides a great
12 opportunity to raise the awareness of nematodes as a constraint to crop production, and for
13 nematologists to be seen to provide strategies in PPN management that are both sustainable
14 and acceptable. There remains a great need to research nematode host status, soil and nursery
15 sanitation, heat treatment, solarization, biofumigation and alternative chemical control for
16 nematodes in ornamentals.

17

18 **2.8 Nematode risks to new crops and varieties introduced in Africa**

19

20 Scientists involved in ongoing international efforts to introduce new crops (some of them
21 biofortified staple crops) should be aware of the potential risks that nematodes can pose.
22 Sweet potato is considered a ‘lifesaver’ crop that rescued Uganda in the 1990s when a virus
23 ravaged the cassava crop, and also in Southern and Central Mozambique in 2009/2010 after a
24 severe drought caused a 32% loss of cassava. The release of orange-fleshed sweet potato
25 (OFSP) drought tolerant, virus resistant, varieties has helped to address both crop insecurity

1 and the widespread problem of vitamin A deficiency in SSA⁸² and it is intended that OFSP
2 will be adopted in Burkina Faso, Ghana, Nigeria and Tanzania. The Ahipa project (funded by
3 the Belgian Development Cooperation), which was launched by the International Potato
4 Center to enhance the nutrient-rich yam bean or ahipa (*Pachyrhizus tuberosus* (Lam.)
5 Spreng.), is an effort to improve human nutrition, food security and sustainability of farming
6 systems in Central and West Africa. Ahipa fixes nitrogen in the soil, making it highly suited
7 to the needs of small farmers as an integral component of sustainable land use. Sweet potato
8 and ahipa are both susceptible to PPN, including RKN, and the implementation of
9 international programs to introduce these crops into African countries must consider the
10 potential of nematodes becoming a serious pest.

11

12 **3 MANAGEMENT STRATEGIES AND CONTROL MEASURES**

13

14 The completion of the nematode life-cycle is dependent on three interacting components: the
15 host plant, the parasite species and the environment (both biotic and abiotic). All
16 management strategies (Fig. 1) are geared to break this ‘pest triangle’ by manipulating one or
17 more of these factors. However, the diversity of PPN makes any overall control strategy
18 problematic. Whatever the strategy, the intent is similar: to reduce the initial PPN in the soil
19 prior to planting and to reduce the subsequent rate of nematode increase on the crop. An
20 overview of technologies applicable to PPN management can be seen in Figure 1. We will
21 review only a few, which are practised in subsistence farming in the region.^{10,11}

22

23 **3.1 Healthy Planting Material**

24

1 A cost effective method of protecting crops is to ensure that planting material is free from
2 PPN. Clean and healthy planting material is the easiest management strategy to protect crops
3 from PPN. Seeds, including tubers and suckers, should not become a source of inoculum as
4 the management of PPN infested soil is much more difficult and costly^{72,73}. Ensuring nursery
5 beds are free from PPN and employing care over soil sanitation also adds to best practice.
6 The burning of trash on the seedbed prior to seeding is practised by some farmers to produce
7 healthy plants for transplanting;⁷² solarization,⁷⁴ incorporation of either crotalaria or Mexican
8 marigold (*Tagetes minuta*) seedlings, and biological control agents introduced in the seedbed
9 have been also tried against RKN.⁷² Other techniques include, hot water treatment of seed
10 stocks,⁸⁰ paring, and use of vitro (tissue culture) plants in banana.^{37,80}

11

12 **3.2 Cultural practices**

13

14 Following harvest, PPN on the roots can be killed by destruction of root material. Cultural
15 practices that expose nematodes to extreme environments (*e.g.*, temperature and/or water
16 stress) will all reduce numbers and are considered good practice.⁷⁴ Uprooting harvested crops
17 and burning infected roots, or even just exposing the roots to sunlight, can all reduce PPN.
18 Trap cropping,⁷⁴ where a nematode host is planted and then removed before the nematode
19 can reproduce, can also reduce nematode infestations, although it is labor intensive and costly
20 and is not usually the method of choice.

21

22 **3.3 Exploiting natural resistance and crop rotation**

23

24 Many PPN do not reproduce equally well on all crops, or even on different cultivars of the
25 same crop. Even RKN, generally regarded as polyphagous, do not reproduce equally well on

1 different plants or even crop cultivars^{80,83} and this knowledge can be exploited to develop
2 crop rotation strategies. Sources of resistance to RKN have been found in locally available
3 genotypes of various popular crops grown by resource-poor farmers, including Brassicaceae,
4 green pepper, maize and tomato.⁸⁴ However, there are few crops for which totally nematode
5 resistant varieties are available.

6
7 The use of crop rotation in a sequential cropping system is widely practised and is useful for
8 managing PPN both in traditional and modern agriculture.^{11,80} Crop rotation, either on its own
9 or in conjunction with other cultural control measures, is the most practical method of
10 nematode management for African agriculture. Farmer participation aimed at ensuring the
11 use of technologies to control RKN on tomato through the use of micro-organisms, cultural
12 techniques and plant resistance within the cropping system preferred by the farmers has been
13 conducted in Kenya.^{72,73} Farmer participatory experimentation on organic cotton has shown
14 that rotation systems can be profitable, sustainable and equitable^{85,86}. This experience can be
15 transferred to countries like Tanzania which is fostering organic cotton, a crop that can be
16 severely affected by RKN.

17

18 **3.4 Chemical Control**

19

20 The use of nematicides has been a preferred option to control nematodes.⁷²⁻⁷⁴ However, they
21 are expensive and toxic, and most are being phased out because of their environmental
22 impact and health hazard. Pesticide-related ill health in several African countries and
23 cropping systems has been documented mainly in smallholder production systems ([http://pan-
24 afrique.org](http://pan-afrique.org)). Data from Tanzania have identified smallholder vegetables as high risk with
25 73% farmers applying pesticides weekly.⁸⁶ There is a need to fund training for farmers' field

1 research and develop ways to increase consumer and market demand for alternative
2 approaches so that an efficient transition between chemical phase-out and phase-in of
3 alternatives can be ensured.

4

5 **3.5 Biological Control and Green Manures**

6

7 The field application of biological control agents (BCA) such as species of the rhizobacteria
8 *Bacillus* and species of the obligate parasite *Pasteuria*, as well as the nematophagous fungus
9 *Pochonia chlamydosporia* (Goddard) Zare & W. Gams, 2001, and others has shown promise
10 in highly managed crop production systems⁸⁷. Recent advances in the use of BCA based on
11 bacterial seed treatments in an IPM context are having major effects on production in large
12 scale farming. For example, *Pasteuria nishizawae* (*Clarica R*) is being applied to soybean
13 seed on millions of hectares for soybean cyst nematode control and *P. penetrans* (Thorne
14 1940) Sayre & Starr, 1986 for nematode control on other crops. Both bacteria can be
15 produced in large amounts under commercial fermentation and their use over time could lead
16 to suppressive soils. The use of *Bacillus firmus* (*Votivo R*) as a bacterial seed treatment on
17 maize, cotton and soya, in combination with fungicide or insecticide to reduce PPN and
18 simultaneously other pests and diseases, is planted on millions of hectares of land in North
19 America. These forms of BCA applications should be examined for use in African
20 agriculture. Seed treatment with selected BCA could have a major impact on nematode
21 management in subsistence farming. Scientists in Africa can easily isolate new local and
22 effective strains of bacteria and fungi that could help in nematode management.⁸⁸ Inoculation
23 of tissue culture seedlings of horticultural crops with fungal antagonists, especially
24 mutualistic fungal endophytes is also being tested in Uganda and Kenya for nematode
25 management in banana and other crops^{33,80} and the fungus *Purpureocillium lilacinum* (Thom)

1 Luangsa-ard, Houbraken, Hywel-Jones & Samson, 2011 (= *Paecilomyces lilacinus*) is also
2 marketed in southern Africa for use in vegetable production and is being investigated for
3 banana applications.⁸⁹ Isolates of *P. chlamydosporia* can be effective in controlling RKN
4 under glasshouse and periurban agriculture conditions.⁹⁰ The search and isolation for African
5 isolates of the fungus from RKN has been done in Ghana, Kenya, Malawi, South Africa and
6 Zimbabwe.⁹¹

7 Soil amendments improve organic matter levels, help with moisture retention, and stimulate
8 microbial activity, thereby contributing to nematode and pathogen control. *Pochonia*
9 *chlamydosporia* is usually added to soil in a colonized rice substrate which acts as an energy
10 source for the fungus. However, in some fungi, high nitrogen/carbon levels can repress
11 infection-related genes and may compromise parasitic ability.⁹² One study in Kenya to
12 determine the efficacy of *P. chlamydosporia* and *P. lilacinum* in combination with labor
13 practices, such as the addition of different organic amendments and crop rotation to control
14 *Meloidogyne* in tomato, resulted in increased tomato yield (by up to 64-65%).⁹³

15 *Purpureocillium lilacinum* strain 251 (PL Plus®), organic substrates, and one botanical
16 pesticide (azadirachtin 0.15%) used in an experiment to assess PPN control in carnations
17 (*Dianthus caryophyllus* L.) showed that all treatments reduced PPN; a significant reduction
18 also occurred in RKN egg mass production and root galling with the fungus treatment.¹⁷

19 Other nematophagous fungi such as *Trichoderma* spp. are starting to be researched alongside
20 resistant tomato varieties for root-knot nematode management in Kenya.⁹⁴

21

22 **3.6 Genetically Modified Crops/Organisms**

23

24 Genetically modified (GM) crops or organisms (GMO) have a role to improve yield per unit
25 of land by mitigating biotic constraints such as pest and disease damage and competition

1 from weeds. It is expected that GM plants will help protect against abiotic constraints such as
2 drought and salinity, and enhance food-quality, post-harvest and processing properties
3 without further expansion of the agricultural frontier.⁹⁵ An example of GM approaches for
4 PPN control are banana and plantain (*Musa* spp.) which often depend on nematicides to
5 increase yield. *Musa* is the third most important crop of SSA, a region that produces 35% of
6 world bananas.⁹⁶ Most edible bananas are sterile and therefore produce no seed, a feature that
7 hampers traditional cross-pollination techniques and thus opens the way to biotechnology
8 approaches that can provide partial resistance to plants against a wide range of nematodes.
9 Transgenic *Musa* have been generated by using East African Highland Banana (EAHB) and
10 plantain OcIΔD86 (a protein-engineered version of the rice cystatin OcI) transformed lines in
11 Uganda (International Institute of Tropical Agriculture). Another approach with transgenic
12 *Musa* includes the integration of synthetic peptides that disrupt the chemosensation of
13 nematodes in a non-lethal manner, thereby reducing the number of nematodes that can locate
14 and invade the roots.⁹⁶

15

16 Biotechnology risk assessment capacity building is required as biosafety assessments for
17 GMOs are usually conducted under National Biosafety Frameworks (NBFs). Many SSA
18 countries have draft NBFs but only few have passed full legislation for GMO release.
19 Regulatory costs imposed on biotechnology created varieties are also a significant hurdle for
20 the development of GM crops suited to developing country farmers. There are some institutes
21 and research groups that make use of biotechnological methods that do not involve high
22 research and registration costs, such as conservation and multiplication of germplasm and
23 phytosanitation.⁹⁵

24

25 **3.7 Integrated Pest Management**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

The two major constraints on controlling PPN in Africa are the lack of knowledge of the biology of PPN, and identifying control measures that are effective and applicable, particularly for small-scale farmers.¹¹ Nematode management should become an integrated program of practices to provide sustainable control systems. The pros and cons of the most important concepts of Integrated Pest Management (IPM) for nematode management for tropical and subtropical crops growing in Africa have been outlined and discussed.⁷⁶ Each farming situation requires a different set of production and protection management inputs for greatest profitability, and to complicate matters further, these requirements may change from year to year. Climate, soil type, cropping system, nematode pest history, cultural practices, cultivar and environmental factors may all affect the nematode problems of a given site. In some areas where resistant cultivars have been deployed, their effectiveness has been rapidly reduced by lack of crop rotation, the emergence of new host races of the target nematode species, or by the emergence of new pests to which the crop is not resistant.

4 NEMATODES AS BENEFICIAL INVERTEBRATES THROUGH ECOSYSTEM SERVICES

Not all nematodes are detrimental to crop productivity – they can also be a benefit to growers by helping to control insect pests and, perhaps more importantly, in the recycling of nutrients. Recently, their importance in food webs has become recognized and they are increasingly being used as indicators of environmental disturbance and reduced biodiversity.^{18,97,98}

4.1 Environmental Indicators

1 Due to the cosmopolitan distribution of nematodes, they inhabit most environments and play
2 a key role in the food webs of many microbial ecosystems, including agricultural soils.⁹⁹
3 Nematode community analysis is attracting increased attention^{100,101} as they are good
4 monitors of biodiversity and environmental indicators.⁹⁷ Bio-safety evaluation of the effect of
5 transgenic banana was performed in Uganda using ecological indices such as Trophic
6 Diversity, Shannon-Weiner Diversity, Simpson's Diversity and various other structure
7 indices of nematode community composition and diversity.¹⁰²

8

9 **4.2 Entomopathogenic Nematodes**

10

11 Entomopathogenic nematode (EPN) species occur in the genera *Steinernema* and
12 *Heterorhabditis*. They have great potential for the biological control of insect pests because
13 they are easy to mass culture and apply, have low host specificity, and are very safe to the
14 user and environment. They serve as an effective, naturally occurring, mechanism for
15 suppressing soil-dwelling insect pests, or insects with at least one life stage inhabiting the
16 soil, and their occurrence usually indicates a well-balanced and diverse soil ecosystem. In
17 East Africa, only a few studies have focused on EPN, these largely being aimed at
18 establishing the presence of indigenous EPN so as to obtain isolates for future use in IPM
19 systems, for example, in Ethiopia,^{103,104} Tanzania (Haukeland S, pers. comm.) and Uganda.

20

21 The immediate need is to improve the capacity of scientists to characterize native EPN using
22 modern molecular methods (essential as they are morphologically conserved and require
23 molecular sequencing techniques for accurate diagnostics), and to provide the basic data and
24 experience for the extensive use of EPN as microbial control agents, thus providing a
25 potentially viable alternative to chemical insecticides, at least in certain situations such as

1 shadehouses, nurseries, etc. This will also serve as stimulation for the development of novel
2 and innovative approaches to pest management and indirectly help open new markets for
3 agricultural products abroad where concern over chemical residues in agricultural products
4 may limit exports.

5

6 **5 CHALLENGES**

7

8 The association of PPN with other soil pathogens in disease complexes, as well as with
9 abiotic stress conditions in crop physiological syndromes, ranks them high on the list of
10 major economic constraints, even if many small holders are unaware of the nature and danger
11 of nematode infestations and overlook these damaging and economically important pests.
12 There are no reliable estimates of the magnitude of annual losses for many crops, but surveys
13 have shown that PPN are widespread in ESA and cause more damage to crops than
14 recognized.^{10,28,72,73,77,81,84} Their association with other soil pathogens in disease
15 complexes,^{65,105,106} as well as with abiotic stress conditions in crop physiological syndromes,
16 ranks them high on the list of major economic constraints. As most PPN are root parasites,
17 they greatly affect water uptake, causing significant losses in the production of principal
18 crops, especially in the context of predicted climate change where restricted and/or
19 increasingly unreliable water resources is expected to impact many African countries.

20

21 **5.1 Education and Capacity Building**

22

23 In ESA, nematology is a relatively new science within the plant protection disciplines.
24 Although there are regional research institutes established, only a few agricultural faculties at
25 Universities teach nematology, whilst only a few national agricultural research institutions

1 conduct research projects in nematology. The number of nematologists (Table 5) is
2 inadequate for the maintenance of a vibrant community. This shortage is exacerbated as most
3 trained nematologists leave the discipline due to lack of suitable positions, institutional and
4 peer support. Another factor is the low profile of nematology in some crop protection and
5 plant pathology programs. Consequently, few research and training institutions employ
6 nematologists who work full time in the discipline. The poor recognition of nematology at
7 management and decision making levels usually results in poor and inadequate resourcing for
8 nematologists. Networks and critical mass are key to harmonized efforts and to prevent
9 isolation while engendering team spirit for accessing support and research resources.

10

11 *5.1.1 University Sector*

12

13 Recent consolidated activities such as the Post-Graduate Nematology Course (PINC)
14 supported by the Belgian Government or the Nematology Initiative for East and Southern
15 Africa (NIESA), supported by the Gatsby Charitable Foundation (UK), for example, have
16 provided important, focused support to nematology. However, greater buy-in from national
17 programs, etc., is needed for its impact to be realized. Nematology needs to be incorporated
18 into the curricula of a greater number of agriculture colleges and universities in the region
19 and trained staff need to be available as instructors.

20

21 In Kenya there has been an increase in the number of permanent staff and postgraduate
22 students in nematology, although these numbers are still inadequate. In Kenyatta University
23 faculty members actively involved in nematology funded projects are increasing and a new
24 MSc course on plant nematology is available. Overall, the number of postgraduate students
25 working in NIESA-participating institutions on nematodes of legumes, cabbages, tomatoes

1 and management of RKN on indigenous vegetables has increased from zero to over 15 during
2 2008-2013, and more MSc students at Sokoine University of Agriculture in Morogoro
3 (Tanzania) have opted to do their research in plant nematology. Although promising progress
4 has been made, there is still a great need for sponsorship of postgraduate students to do
5 nematology research in ESA.

6

7 *5.1.2 Extension and knowledge transfer: links to farmers*

8

9 The absence of extension staff has been noted in nematode workshops and training courses
10 held in the region. This group needs to be specifically targeted since they are well positioned
11 to facilitate transfer of tested technologies to growers. Although direct interaction with
12 individual farmers has been limited, capacity building projects have had an impact on
13 smallholder farmers. Rotations involving maize have been used to reduce RKN infestations
14 in tomato during the second cropping season in Kenya¹⁰⁷. This is a technology that farmers
15 understand and can adopt. The effect of different cropping cycles on RKN infestations has
16 been recognized by smallholders in areas where nematode surveys were carried out in
17 Zimbabwe. Farmers saw the benefits of cropping sequences that included a poor host to the
18 nematode pest or a long fallow period. Some growers are now able to distinguish between a
19 root system that has been damaged by nematodes and one that is healthy. There has also been
20 an increase in the number of samples being submitted to laboratories for analysis, more
21 requests for field visits and an increased awareness of dissemination of nematodes through
22 contaminated soil and the potential exclusion of pests.

23 The delivery of extension services has met with limited success worldwide. The ‘going
24 public’ and mobile ‘Plant Health Clinic’ examples (www.plantwise.org) are a positive
25 innovation in the region in the two-way communication between researchers and farmers.¹⁰⁸

1 However, they depend upon the availability of trained diagnosticians (including
2 nematologists) to attend public gatherings and disseminate appropriate knowledge to the
3 farmers. Extension scientists should also be knowledgeable regarding the role of quarantine
4 agencies and the need for appropriate diagnostics in preventing the introduction of new pest
5 nematodes.

7 *5.1.3 Agricultural Research and Development*

8
9 Another constraint to nematology in the region is the importance of agricultural research and
10 development as a driver of innovative crop protection. Establishing collaborative,
11 interdisciplinary research (soil scientists, soil ecologists, breeders, economists, statisticians,
12 sociologists) within and among countries is necessary to raise the profile, and hence funding,
13 of nematology in the region, especially if linked to developing effective IPM programs for
14 the major crop-nematode combinations so as to improve crop health and yield.

15
16 Additionally, there is a critical need to quantify crop losses and develop related damage
17 thresholds. Gathering meaningful data on yield losses caused by PPN in the region is crucial
18 but has been hampered by several factors, amongst them:

- 19 • the area of land held by small/poor-resource farmers (up to 70% of farmers own less
20 than 2 ha).^{108,109}
- 21 • the multitudinous combinations of vegetables and other staple crops grown in such
22 plots, these crops being potential hosts to several different species of PPN, amongst
23 them the various species of RKN.

24 Lack of knowledge can be dealt with through concerted action protocols between
25 stakeholders and execution of carefully planned experiments to resolve problems for the

1 major nematode pests on a regional basis and to analyze and quantify the costs of nematode
2 damage to the grower. This should be combined with devising effective alternatives of
3 nematode control in smallholder farm landscapes, developing, for example: *i)* rotations
4 coupled with studies of nematode host-ranges; *ii)* evaluating solarization as an important
5 control measure; *iii)* giving priority to development and deployment of host resistance for
6 nematode control; and *iv)* using demonstration plots, preferably in collaboration with local
7 farmers, to show the potential of soil amendments, biological control agents, field sanitation,
8 destruction of residual infected roots and clean planting material as important components of
9 integrated nematode control strategies and integrated biology management through
10 manipulation of farming systems, the environment and the host to enhance/develop a soil
11 biological community capable of suppressing nematode pests.¹¹⁰

12

13 The diversity of unique crops grown in many areas and different climatic zones, in
14 combination with local cropping systems, are often poorly understood with regard to
15 nematodes and their impact on crops. Solutions to these problems cannot simply be imported
16 using experiences and data from other regions of the world as these are often inapplicable or
17 inappropriate for the climatic, cropping systems or socio-economic factors that prevail. A
18 knowledge bank needs to be produced in ESA if we are to address the increasing food
19 shortages that seem likely to develop.

20

21 *5.1.4 Plant Inspection and Quarantine Services*

22

23 One of the basic principles in crop protection is the exclusion of pathogens from uninfected
24 areas. Whereas opportunities for the introduction of new nematodes are numerous and
25 frequent, ordinary plant quarantine regulatory actions within ESA are not well adapted to the

1 detection of PPN, most quarantine actions still relying upon simple visual inspections
2 supplemented by a hand lens or low power microscope, rather than molecular diagnostic
3 tools.

4
5 Although some endemic (e.g. *Aphelenchoides arachidis*, *Ditylenchus africanus*, *Globodera*
6 *capensis*, *Meloidogyne ethiopica*, *Scutellonema bradys*) and non-endemic African PPN
7 species (i.e. *Globodera rostochiensis*) currently have a limited distribution, they represent
8 potential threats to ESA countries and national quarantine regulations need to be
9 implemented. African nematodes such as *M. ethiopica* are included in the alert quarantine list
10 of international quarantine bodies such as EPPO (European and Mediterranean Plant
11 Protection Organization) https://www.eppo.int/QUARANTINE/Alert_List/alert_list.htm)
12 because of their potential economic impact.

13

14 *5.1.5 Nematode diagnostic services*

15

16 Careful sampling and accurate identification of nematodes present in planting material or in
17 established crops are essential to determine whether the species found are potentially
18 injurious, are the cause of existing disease problems, or can be classified as non-injurious.
19 Examination for PPN is usually valueless unless precise laboratory extraction protocols are
20 employed and identification is done by experienced nematologists. The use of image
21 databases and other expert identification systems, in conjunction with molecular techniques
22 with robust and standardized protocols in accredited official laboratories, are required for
23 analysis of plant materials so as to assure reliability and consistency. How this infrastructure
24 should be funded is a moot point, but contributions from national governments, overseas
25 agencies, grower's organizations or crop levies are all possible.

1
2 Currently, classical methods of nematode identification based on morphological and
3 morphometric characteristics, supplemented with host range tests, are used. The procedures
4 involved can be subjective, time-consuming and cannot always provide prompt answers and
5 timely advice to farmers. In addition, the specialized skills required for nematode
6 identification at the species level are not usually available and are becoming less so as
7 experienced taxonomists retire. Problems in this area could be alleviated through the use of
8 molecular, immunological and biochemical methods of nematode diagnosis, possibly in the
9 form of kits, which are more rapid, require less specialized skills and enable the identification
10 of cryptic species and juveniles. There is presently no centre within ESA that has the facilities
11 and expertise to do this. The laboratories established in ESA countries are equipped for
12 morphological work but would need to be upgraded in order to support appropriate molecular
13 and biochemical diagnostics. Molecular diagnostics of nematodes is an area that will need to
14 be developed as a matter of priority but needs to overcome the challenges due to limited
15 economic resources, infrastructure and trained personnel.¹¹¹ Quality Plant Pest Diagnostic
16 Services can be costly since they require “quality science, training, infrastructure” and in the
17 system of informal, low value food chains of staple crops prevalent in many African nations
18 it is not clear how such costs can be supported. A key issue is the positioning of plant health
19 standards to stimulate private sector engagement and a greater volume of formal local and
20 regional trade.¹¹¹

21
22 Examples of initiatives to build up pest diagnostic capacity include training projects,
23 networks and regional centers of excellence as exemplified, among others, by Bioscience
24 East and Central Africa (BeCa) in Kenya, the International Plant Diagnostic Network (IPDN)
25 a USAID-support initiative with two regional networks in West and East Africa, and NIESA.

1 The PlantWise worldwide diagnostic program (www.plantwise.org) of plant clinics
2 developed by CABI is a valuable resource for plant protection scientists in Africa and around
3 the globe, although nematology remains an area requiring further development.¹¹¹
4

5 **5.2 Knowledge Transfer: links to farmers**

6

7 The absence of extension staff has been noted in nematode workshops and training courses
8 held in the region. This group needs to be specifically targeted since they are well positioned
9 to facilitate transfer of tested technologies to growers. Although direct interaction with
10 individual farmers has been limited, capacity building projects have had an impact on
11 smallholder farmers. Rotations involving maize have been used to reduce RKN infestations
12 in tomato during the second cropping season in Kenya.¹⁰⁷ This is a technology that farmers
13 understand and can adopt. The effect of different cropping cycles on RKN infestations has
14 been recognized by smallholders in areas where nematode surveys were carried out in
15 Zimbabwe. Farmers saw the benefits of cropping sequences that included a poor host to the
16 nematode pest or a long fallow period. Some growers are now able to distinguish between a
17 root system that has been damaged by nematodes and one that is healthy. There has also been
18 an increase in the number of samples being submitted to laboratories for analysis, more
19 requests for field visits and an increased awareness of dissemination of nematodes through
20 contaminated soil and the potential exclusion of pests.

21

22 **5.3 Partnerships with stakeholders**

23

24 Countries such as Kenya and Uganda have been considered to “support the highest number of
25 national agriculture-related PhD and have some of the best-equipped laboratories”¹¹¹ but

1 despite professional expertise and infrastructure, they remain vulnerable to new and emerging
2 pests, prompting partnerships with Non-Governmental Organizations (NGOs) and the private
3 sector. Capacity building is a key area that connects commonalities within the discipline of
4 nematology in ESA and can develop synergy in activity. PPN control requires the creation
5 and maintenance of a network of nematologists that contribute to the pipeline between
6 research, development and the transfer of this knowledge to the grower. A policy that is
7 supportive of organizations and stakeholders, such as governments, research institutes,
8 universities, businesses, NGO, growers (large and small scale), extension workers, retailers,
9 consumers, educators and trainers engaged in delivering increased crop security through the
10 control of PPN, is needed.

11

12 *5.3.1 Call to action and delivery pipeline*

13

14 There is an on-going need to establish a set of wider scientific links with other nematologists
15 and/or institutions in the region for collaborative research projects targeted at local needs.
16 Such a campaign requires the establishment of nematological networks fostered by the use of
17 the internet to increase communications among regional nematologists and to provide access
18 to information resources. This will maintain and increase the critical mass of skilled people
19 via short term training aimed at the regional extension and plant inspection services. Implicit
20 in this is the requirement for career structures to allow ‘new-blood’ nematologists (*i.e.*, PhD
21 trained) to develop their careers as full-time nematologists, and require the provision of the
22 necessary infrastructure, such as laboratory space and essential capital equipment, to establish
23 and maintain research laboratories within the discipline. A key component will be the
24 engagement of numerous stakeholders that straddle the traditional boundaries between the

1 public sector, private sector and academia, not to mention gaining the support of other crop
2 protection scientists in other disciplines.

3

4 *5.3.2 Growers*

5

6 Throughout the whole area the systems of agricultural production practised by growers is
7 hugely diverse (Table 3) and nematology has to fit within this context. From poorly resourced
8 subsistence farmers undertaking low intensity mixed cropping for local consumption, to
9 highly intensive crop production systems that are mechanized to a large degree to produce
10 commodities for national and international markets, nematology needs to fit within, and adapt
11 to, an overall crop protection context. Between these two extremes there are smallholder
12 farmers that primarily employ subsistence production techniques but also plant commodity
13 crops to bring in necessary cash for basic requirements and also act as an insurance in the
14 event of crop failure. Although described here as polar opposites, there is actually a
15 continuum from low intensity agriculture to high intensity agriculture that is dependent on
16 infrastructure parameters such as proximity to markets and communication systems. Within
17 this crop protection context there are various stakeholders and organizations with which
18 growers need to interact.

19

20 *5.3.4 Non-Governmental Organizations*

21

22 The number and influence of agricultural NGOs working in Africa has grown exponentially
23 over the past 30 years,¹¹² and their contribution to the development and delivery of
24 technology to farmers is evident through increased advocacy at local and national level
25 related to promoting policy and research to benefit specific sub-groups of a population, and

1 influencing the strategy through which funds are allocated or projects designed. Seeking
2 complementarity and synergy with NGOs would improve community access to nematode
3 management technologies, especially when considering scale and impact, and the means by
4 which local knowledge and ideas can be leveraged to promote appropriate adoption of the
5 technologies. Clearly, the dissemination of knowledge through the development of online
6 data bases such as PlantWise (www.plantwise.org) as mentioned above should be built upon
7 and exploited through a coordinated approach.

8

9 *5.3.5 Private Sector*

10

11 This group includes commercial companies commonly associated with input supply or output
12 markets, and farmers' associations on crop or area basis. An input supplier provides an
13 avenue for collaborating with nematologists who would undertake extension activities of the
14 supplier as a part of marketing of the inputs. Extension and information channels may include
15 written information, posters, farmer meetings, radio, on-farm demonstrations, exhibits at farm
16 shows, and the use of mobile phones and the internet. In Zimbabwe, for instance, a mobile
17 information system has been set up that provides regular market information and agronomic
18 tips to farmers. This system could easily be exploited to provide basic information on
19 available nematology services. These new information channels will also provide an
20 opportunity to the nematologist to build and maintain good relations with farmers as they will
21 be providing information regarding access to, and use of, inputs.

22

23 Engaging commercial companies associated with output markets could be achieved through
24 providing extension services to their contract growing schemes where extension activities
25 allow close and frequent contact with their farmers. Engaging farmers' associations in

1 nematode management technology delivery might be more difficult as the association must
2 have revenue to provide services to their members.

3

4 **5.4 Raising Awareness of Damage Caused by Nematode Pests Among Farmers,** 5 **Scientists and Policy Makers.**

6

7 Fundamentally, raising awareness is an educational issue that can be achieved through
8 training programs at various levels: training of trainers, training workshops, farmer schools,
9 postgraduate training, conference presentations, publications, development of interactive
10 websites. Although there is better appreciation of nematodes as crop pests among scientists
11 and some farmers, it remains essential to raise awareness among policy-makers. International,
12 regional and national training workshops can be designed and presented for different
13 audiences. This is achievable, as shown by a nematode awareness presentation in Arusha
14 during the Tanzania Horticultural Association (TAHA) annual meeting in 2008. The fact that
15 the meeting was for all stakeholders throughout the horticulture field, including farmers,
16 extension staff, researchers, people from micro-finances and politicians, provided a unique
17 opportunity to discuss the economic importance of nematodes in crops involving policy-
18 makers, although addressing such a diverse audience is challenging.

19

20 **5.5 Climate Change**

21

22 Future climate projections indicate Africa as one of the most vulnerable continents to climate
23 change and climate variability, a situation aggravated by the interaction of ‘multiple stresses’,
24 occurring at various levels and low adaptive capacity.^{12,113} The projections to 2100 for Africa
25 consist of increased warming throughout the continent, and in all seasons, of up to 1.5 times

1 higher than the global average; decrease in annual rainfall in Mediterranean Africa and
2 Southern Africa but an increase in East Africa, and an increase in frequency and intensity of
3 droughts and floods¹¹⁴. Agricultural production will be severely compromised due to land
4 loss, shorter growing seasons, and greater uncertainty about what and when to plant¹¹⁵⁻¹¹⁷.
5 Higher soil temperatures may increase reproductive fitness and pathogenicity of soil-borne
6 diseases and pests, including PPN,¹¹⁸ as well as leading to the evolution of more pathogenic
7 populations of diseases that are currently benign.

8

9 **5.6 Capacity Building and Retooling**

10

11 There is a need to increase nematology research, extension service and training capacity, both
12 through funding of projects by institutional grants and support of advanced training
13 programmers. National agricultural research systems must be improved and strengthened
14 financially and technically to act as centers of excellence for the advancement of regional
15 agricultural nematology. This will facilitate research, development and implementation of
16 national nematode pest management programs; engage in training and delivery of
17 information to ensure increased nematode recognition by the farming community; develop
18 databases that quantify the relationship between nematode populations and crop-yield losses
19 under the prevailing biotic and abiotic factors; and reduce losses caused by nematodes by
20 implementing sustainable, environmentally friendly and more productive agricultural
21 systems.

22

23 It is advisable that efforts aimed at technology dissemination use demonstration plots and
24 participatory approaches. Initially, awareness must be created amongst the farmers, followed
25 by facilitated access and promotion of technologies such as the use of improved varieties or

1 good crop management practices so as to reduce pests and disease and improve post-harvest
2 handling. Capacity building and facilitating access to information is urgently needed by all
3 participants in the value chain.

4

5 **6 CONCLUSIONS**

6

7 There is a serious lack of knowledge on nematode pests of staple food crops in ESA cropping
8 areas and consequently only minimal awareness among farmers, extension workers and
9 politicians of the damage they cause. This lack poses numerous problems in securing
10 improved crop security through PPN control, including the procurement of essential funding.
11 Whilst preparing this review it became clear that research is most developed with respect to
12 cash crops grown for export, with little or no research on staple food crops, particularly those
13 grown by smallholders in multi-crop systems. Familiarization of smallholders with
14 nematodes has only been a small part of internationally funded projects aimed at developing
15 and encouraging control strategies for pests and diseases, the main targets of funding
16 initiatives being insects and fungal pathogens that attack the plant above ground. Whilst this
17 may be understandable from a macro-economic point of view, attention needs to be re-
18 focused further down the supply chain, particular objectives being food security,
19 sustainability and poverty alleviation. Farmer/researcher linkages must be developed so that
20 the latter are aware of nematode-related problems developing in the field and the former can
21 be informed of appropriate management programs. It is important that the damage caused by
22 nematodes and the benefits attained from their control are adequately demonstrated to the
23 farmer so that uptake success can be enhanced. More support and facilities will result from
24 increased awareness among growers and administrators. Future priorities include:

25 • Capacity building in nematology, including appropriate diagnostic services

- 1 • Extension as a key factor in knowledge dissemination between farmers, academia, and
2 local phytosanitation services (*i.e.*, health inspectors)
- 3 • Focus on major nematode pests include species of root-knot nematodes
4 (*Meloidogyne*), *Pratylenchus*, *Ditylenchus* and *Helicotylenchus*
- 5 • Channel efforts towards creating information on the economic importance of these
6 nematodes on staple crops such as cassava, maize and vegetables, but also major
7 commodities (*e.g.*, banana, coffee, cotton, peanut, ornamentals)
- 8 • Attracting appropriate funding to deliver priorities

9

10 In the context of predicted climate change and the possibility of diminishing water resources,
11 as well as the need to intensify crop production systems, PPN are likely to become
12 increasingly problematic in Africa. The opening and maintenance of communication channels
13 throughout the crop protection pipeline, from appropriate basic research and its development
14 to the delivery of nematode control, is therefore essential.

15

16 **ACKNOWLEDGEMENT**

17

18 The authors thank The Gatsby Charitable Foundation for financial support for the NIESA
19 (Nematology Initiative for East and Southern Africa) capacity building initiative, a project
20 that was envisaged and led by the late Professor Brian R Kerry.

21

22 **REFERENCES**

23

24 1 Luc M, Bridge J and Sikora RA, Reflections on nematology in subtropical and tropical
25 agriculture, in *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, ed.

- 1 by Luc M, Sikora RA and Bridge J. CAB Publishing, Wallingford, UK, pp. 1-10
2 (2005).
- 3 2 Chitwood DJ, Research on plant-parasitic nematode biology conducted by the United States
4 Department of Agriculture-Agriculture Research Service *Pest Manag Sci* **59** :748-753
5 (2003).
- 6 3 Smiley R, Plant parasitic nematodes affecting wheat yield in the Pacific West. Oregon State
7 University Extension Services EM 8887 (2005).
- 8 4 Mudiope J, Coyne DL, Adipala E and Talwana HAL, Damage to yam (*Dioscorea* spp.) by
9 root-knot nematode (*Meloidogyne* spp.) under field and storage conditions in Uganda.
10 *Nematropica* **42** :137-145 (2012).
- 11 5 Coyne DL, Akpheokhai LI and Adeniran AF, The yam nematode (*Scutellonema bradys*), a
12 potential threat to potato (*Solanum tuberosum*) production in West Africa. *Plant Pathol*
13 **60** :992-997 (2011).
- 14 6 Sharma SB, Price N and Bridge J, The past, present and future of plant nematology in
15 International Agricultural Research Centres. *Nematol Abstracts* **66** :119-142 (1997).
- 16 7 Duponnois R, Neyra M, Senghor K, Bâ AM, Effects of the root-knot nematode
17 *Meloidogyne javanica* on the symbiotic relationships between different strains of
18 *Rhizobium* and *Acacia holosericea* (A Cunn. ex G. Don). *Eur J Soil Biol* **35** :99-105
19 (1999).
- 20 8 Kimenju JW, Karanja NK and Macharia I, Plant parasitic nematodes associated with
21 common bean in Kenya and the effect of *Meloidogyne* infection on bean nodulation.
22 *Afr Crop Sci J* **7** :503-510 (1999).
- 23 9 Coyne DL and Oyekanmi EO, Symbiotic nitrogen fixation of two soybean genotypes as
24 affected by root-knot nematodes and microsymbionts. *J Biol Sci* **7** :122-226 (2007).

- 1 10 Bridge J, Plant nematodes of different crops and cropping systems in Africa. *South*
2 *African Nematology Symposium*, March (1995).
- 3 11 Bridge J, Nematode management in sustainable and subsistence agriculture. *Ann Rev*
4 *Phytopathol* **34** :201-255 (1996).
- 5 12 von Braun J, Swaminathan MS and Rosegrant MW, Agriculture, Food Security, Nutrition
6 and the Millennium Development Goals, 2003-2004 IFPRI Annual Report (2004).
- 7 13 World Bank, World Development Report 2008: Agriculture for Development. The World
8 Bank, Washington DC, USA (2008).
- 9 14 The Montpellier Panel Report, *Sustainable intensification: A new paradigm for African*
10 *Agriculture*. Agriculture for Impact, Imperial College, London, UK
11 www.ag4impact.org (2013).
- 12 15 Dixon J, Gulliver A and Gibbon D, Farming Systems and Poverty: Improving Farmers
13 Livelihoods in a Changing World. Rome, Italy, and Washington, D.C.: FAO and The
14 World Bank (2001).
- 15 16 Achterbosch T, Allbritton A, Quang DV, de Jager A, Njue E, Sonko R, Stallen M,
16 Wertheim-Heck S, van Wijk S (2005) Pro-Poor Horticultural Growth in East Africa and
17 South East Asia.
18 r4d.dfid.gov.uk/PDF/Outputs/EC.../ProPoorHorticultureFinalReport.pdf
- 19 17 Kimenju JW, Wachira PM, Lang'at JK, Otieno W and Mutua GK, Evaluation of selected
20 methods in the control of plant parasitic nematodes infecting carnation. *J Agric Sci* **6**
21 :2014. Doi: 10.5539/jas.v6n3p31 (2014).
- 22 18 Lnagat JK, Kimenju JW, Mutua GK, Muiro WM and Otieno W, Response of free-living
23 nematodes to treatments targeting plant parasitic nematodes in carnation. *Asian J Plant*
24 *Sci* **7** :467-472 (2008).

- 1 19 Viaene N, Coyne DL and Kerry BR, Biological and cultural management. In: *Plant*
2 *Nematology*, ed. by Perry RN and Moens M, pp. 346-369, CABI, UK (2006).
- 3 20 Kashaija IN, Speijer PR, Gold CS and Gowen S, Occurrence, distribution and abundance
4 of plant parasitic nematodes of bananas in Uganda. *Afr Crop Sci J* **2** :99-104 (1994).
- 5 21 Speijer PR, Gold CS, Karamura EB and Kashaija IN, Banana weevil and nematode
6 distribution patterns in highland banana systems in Uganda: preliminary results from
7 diagnostic survey, in *Proceedings of the First International Crop Science Conference*
8 *for Eastern and Southern Africa*, ed. by Adipala E, Bekunda MA, Tenywa JS, Ogenga-
9 Latigo MW and Mugah JO. Kampala, 14-18 June 1993, Makerere University,
10 Kampala, Uganda, pp. 285-289 (1994).
- 11 22 Seshu Reddy KV, Prasad JS, Speijer PR, Sikora RA and Coyne D, Distribution of plant-
12 parasitic nematodes on *Musa* in Kenya. *InfoMusa* **16** :18-23 (2007).
- 13 23 Bridge J, Nematodes of bananas and plantains in Africa: research trends and management
14 strategies relating to the small-scale farmer. Key note. International Conference on
15 Banana and Plantain for Africa. Kampala, 14-18 October, 1996. Leaflet 23 (1996).
- 16 24 Frison E and Sharrock S, The economic, social and nutritional importance of banana in the
17 world, in *Bananas and Food Security*, ed. by Picq C, Fouré E and Frison EA.
18 Proceedings of an International Symposium held in Douala, Cameroon, 10-14
19 November, 1998, INIBAP, Montpellier, France, pp. 21-35 (1999).
- 20 25 Kimenju JW, Waudu SW, Mwangombe AW, Sikora RA and Schuster RP, Distribution of
21 lesion nematodes associated with maize in Kenya and susceptibility of maize cultivars
22 to *Pratylenchus zaeae*. *Afr Crop Sci J* **6** :367-375 (1998).
- 23 26 Talwana HL, Butseyia MM and Tusiime G, Occurrence of plant parasitic nematodes and
24 factors that enhance population build-up in cereal-based cropping systems in Uganda.
25 *Afr Crop Sci J* **16** :119-131 (2008).

- 1 27 Bridge J, Coyne D and Kwoseh CK, Nematode parasites of tropical root and tuber crops,
2 in *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc M,
3 Sikora RA and Bridge J, CAB International, Wallingford, UK, pp. 221-258 (2005).
- 4 28 Coyne DL, Talwana LAH and Maslen NR, Plant parasitic nematodes associated with root
5 and tuber crops in Uganda. *African Crop Protection* **9** :87-98 (2003).
- 6 29 Coyne DL, Tchabi A, Baimey H, Labushagne N and Rotifa I, Distribution and prevalence
7 of nematodes (*Scutellonema bradys* and *Meloidogyne* spp.) on marketed yam
8 (*Dioscorea* spp.) in West Africa. *Field Crops Res* **96** :142-150 (2006).
- 9 30 Mudiope J, Speijer PR, Coyne D, Maslen RN and Adipala E, Nematode distribution and
10 damage to yam in central and eastern Uganda. *Afr Crop Sci J* **15** :93-99 (2007).
- 11 31 Bafokuzara N, Incidence of different nematodes on vegetable and fruits crops and
12 preliminary assessment of yield loss due to *Meloidogyne* species in Uganda. *Nematol*
13 *Bras* **20** :32-43 (1996).
- 14 32 Kimenju JW, Kagundu AM, Nderitu JH, Mambala F, Mutua GK and Kariuki GM,
15 Incorporation of green manure plants into bean cropping systems contributes to root
16 knot nematode suppression. *Asian J Plant Pathol* **4** :404-408 (2008).
- 17 33 Gheysen G, Kyndt T, Soraya de Carvalho F, Höfte M, Bert W, Janssen T, Mibey RK and
18 Njira PN, Analysis of endophytic fungi and plant-parasitic nematodes from irrigated
19 and upland rice ecosystems in Kenya. In *Proceedings 6th International Congress of*
20 *Nematology, 4-9 May 2014, Cape Sun, Cape Town South Africa*, p. 112 (2014).
21 [Abstract]
- 22 34 Kagoda F, *Genetic studies and recurrent selection for nematode resistance in maize*. A
23 thesis submitted in partial fulfillment of the requirements for the degree of Doctor of
24 Philosophy (PhD) in Plant Breeding. African Centre for Crop Improvement School of

- 1 Agricultural Sciences and Agribusiness Faculty of Science and Agriculture University
2 of KwaZulu Natal, Pietermaritzburg, South Africa (2010).
- 3 35 Mc Donald AH and Nicol JM, Nematode parasites of cereals, in *Plant Parasitic*
4 *Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc M, Sikora RA and
5 Bridge J, CAB International, Wallingford, UK, pp. 131-192 (2005).
- 6 36 Nyombi K, Towards sustainable highland banana production in Uganda: opportunities and
7 challenges *AJFAND* **13** :7544-7561 (2013).
- 8 37 Gowen SR, Quenéhervé P and Fogain R, Nematode parasites of bananas and plantains, in
9 *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc M,
10 Sikora RA and Bridge J, CABI, Wallingford, UK, pp. 611-643 (2005).
- 11 38 Speijer PR and Sikora RA, Influence of a complex disease involving *Pratylenchus*
12 *goodeyi* and a non-pathogenic strain of *Fusarium oxysporum* on banana root health, in
13 *Biological and Integrated Control of Highland Banana and Plantain Pests and*
14 *Diseases*, ed. by Gold CS and Gemmill B, International Institute of Tropical
15 Agriculture, Ibadan, Nigeria, pp. 231-239 (1993).
- 16 39 Price NS and Bridge J, *Pratylenchus goodeyi* (Nematoda, Pratylenchidae), a plant-
17 parasitic nematode of the montane highlands of Africa. *J Afr Zool* **109** :435-442 (1995).
- 18 40 Wehunt EJ, Hutshinson DJ and Edwards DI, Reactions of banana cultivars to the
19 burrowing nematode. *J Nematol* **10** :368-370 (1978).
- 20 41 Pinochet J and Rowe PR, Progress in breeding for resistance to *Radopholus similis* on
21 bananas. *Nematropica* **9** :76-78 (1979).
- 22 42 Sarah JL, Blavignac F, Sabatini C and Boisseau M, Une méthode de laboratoire pour le
23 criblage variétal des bananiers vis-à-vis la résistance aux nématodes. *Fruits* **47** :559-
24 564 (1992).

- 1 43 Price NS, Field trial evaluation of nematode susceptibility within *Musa*. *Fund Appl*
2 *Nematol* **17** :391-396 (1994).
- 3 44 Viaene N, Duenas J and De Waele, D, Screening for resistance and tolerance to
4 *Radopholus similis* and *Pratylenchus coffeae* in banana and plantain, in *Programme*
5 *and Abstracts of the 24th International Symposium of the European Society of*
6 *Nematologists, 5-8 Aug. 1998*, ed. by Brown D, Dundee, UK, p. 125 (1998).
- 7 45 Fogain R and Gowen SR, Yangambi km5 (*Musa* AAA, Ibota subgroup) a possible source
8 of resistance to *Radopholus similis* and *Pratylenchus coffeae*. *Fund Appl Nematol* **21**
9 :75-80 (1998).
- 10 46 Stoffelen R, Verlinden R, Pinochet J, Swennen RL and De Waele D, Host plant response
11 of *Fusarium* wilt resistant *Musa* genotypes to *Radopholus similis* and *Pratylenchus*
12 *coffeae*. *Int J Pest Manag* **46** :289-293 (2000).
- 13 47 Stoffelen R, Verlinden R, Xuyen NT, Swennen RL and De Waele D, Host plant response
14 of *Eumusa* and *Australimusa* bananas (*Musa* spp.) to migratory and root-knot
15 nematodes. *Nematology* **2** :907-916 (2000).
- 16 48 Okech SHO, Gold CS, Speijer P, Ssali H and Karamura E, Relationships between soil
17 fertility, banana weevil and nematodes in the East Africa highland cooking banana in
18 Ntungamo, south western Uganda. In *Proceedings of the First International*
19 *Conference on Banana and Plantain for Africa*, ed. by Craenen K, Ortiz R,
20 Karamura EB and DR Vuylsteke. *ACTA Horticulturae* **540** :505-511 (2000).
- 21 49 McIntyre BD, Speijer PR, Riha SJ and Kizito F, Effects of mulching on biomass,
22 nutrients and soil water in banana inoculated with nematodes. *Agronomy Journal* **92**
23 :1081-1085 (2000).

- 1 50 Ssali H, McIntyre BD, Gold CS, Kishaija IN and Kizito F, Effects of mulch and
2 mineral fertilizer on crop, weevil and soil quality parameters in highland banana,
3 *Nutr Cycl Agroecosys* **65** :141-150 (2003).
- 4 51 Gold CS, Okech SH, McIntyre BD, Kagezi G, Ragama PE and Night G, Effects of
5 mulch on banana weevil *Cosmopolites sordidus* (Germar) populations and damage in
6 Uganda, *Crop Protection* **25** :1153-1160 (2006).
- 7 52 Blair MW, Gonzalez LF, Kimani PM and Butare L, Genetic diversity, inter-gene pool
8 introgression and nutritional quality of common beans (*Phaseolus vulgaris* L.) from
9 Central Africa, *Theor Appl Genet* **121** :237-248 (2010).
- 10 53 Ngundo BW and Taylor DP, Effect of *Meloidogyne* spp. on bean yields in Kenya. *Plant*
11 *Dis Rep* **58** :1020-1023 (1998).
- 12 54 Gedil M and Sartie A, Perspectives on molecular breeding on Africa's main staple food
13 crops cassava and yam, in *Aspects of Applied Biology 96, Agriculture: Africa's "engine*
14 *for growth" – plant science & biotechnology hold the key*, ed. by Bruce T, Foyer C,
15 Halford N, Keys A *et al.* Association of Applied Biologists, Warwick, pp. 123-135
16 (2010).
- 17 55 Coyne DL, Nematode parasites of cassava. *Afr Crop Sci J* **2** :355-359 (1995).
- 18 56 Makumbi-Kidza NN, Speijer PR and Sikora RA, The influence of *Meloidogyne incognita*
19 on growth and storage-root formation of young cassava, *Manihot esculenta* Crantz,
20 plants. *Suppl to J Nematol* **32**(4S) :475-477 (2000).
- 21 57 Bridge J, Otim-Nape GW and Namaganda JM, The root-knot nematode, *Meloidogyne*
22 *incognita* causing damage to cassava in Uganda. *Afro-Asian J Nematol* **1** :116-117
23 (1991).

- 1 58 Scurrah MI, Niere B and Bridge J, Nematode parasites of *Solanum* and sweet potatoes, in
2 *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc M,
3 Sikora RA and Bridge J, CAB International, Wallingford, UK, pp. 193-219 (2005).
- 4 59 Akoroda MO, Aikpokpodion P, Aliyu TO, Fabunmi TO, Fatunbi AO and Olofinji EB,
5 Holistic sweet potato breeding and selection schemes: clonal trials in southwest
6 Nigeria. In: *Proc. 5th African Potato Association Conference*, pp. 61-67 (2000).
- 7 60 Coyne D, Nematodes - Section 4.8.4 within Chapter 4 Pests, disease and the agro-
8 ecosystem, in *Manual for Sweetpotato Integrated Production and Pest Management*
9 *Farmer Field Schools in sub-Saharan Africa*, ed. by Stathers T, Namanda S, Mwanga
10 ROM, Khisa G and Kapinga R. International Potato Centre, Kampala, Uganda, pp. 64-
11 65 (2005).
- 12 61 Verhaeven M, Root-knot nematodes in Tanzania: biocontrol and species characterization
13 based on isozyme phenotypes and mitochondrial sequences, MSc Thesis, Ghent
14 University, Belgium (2014).
- 15 62 Baffes J, *Tanzania's cotton sector: constraints and challenges in a global environment*.
16 Africa Region Working Paper Series No. 42. World Bank Group. December 2002, 49
17 pp. (2002).
- 18 63 Bridge J, Nematodes, in *Cotton Diseases*, ed. by Hillocks RJ, CAB International,
19 Wallingford, UK, pp. 331-353 (1992).
- 20 64 Sidhu G and Webster JM, Genetics of resistance in the tomato root-knot nematode-wilt-
21 fungus complex. *J Hered* **65** :153-156 (1974).
- 22 65 Hillocks RJ and Bridge J, The role of nematodes in Fusarium wilt of cotton. *Afro-Asian J*
23 *Nematol* **2** :35-40 (1992).
- 24 66 Birchfield W, Host-parasite relations of *Rotylenchulus reniformis* on *Gossypium hirsutum*.
25 *Phytopathology* **52** :862-865 (1962).

- 1 67 Cadet P and Spaull VW, Nematode parasites of sugarcane, in *Plant Parasitic Nematodes*
2 *in Subtropical and Tropical Agriculture*, ed. by Luc M, Sikora RA and Bridge J, CAB
3 International, Wallingford, UK, pp. 645-674 (2005).
- 4 68 Chirchir AK, Kimenju JW, Olubayo FM and Mutua GK, Abundance and distribution of
5 plant parasitic nematodes associated with sugarcane in Western Kenya. *Asian J Plant*
6 *Pathol* **2** :48-53 (2008).
- 7 69 Chirchir AK, Kimenju JW, Olubayo F and Mutua G, Cultivar resistance of sugarcane and
8 effects of heat application on nematodes in Kenya. *International Journal of*
9 *Agricultural Research* **6** :93-100 (2011).
- 10 70 Johnson CS, Way J and Barker KR, Nematode parasites of tobacco, in *Plant Parasitic*
11 *Nematodes in Subtropical and Tropical Agriculture*, ed. Luc M, Sikora RA and Bridge
12 J, CAB International, Wallingford, UK, pp. 675-708 (2005).
- 13 71 Priya, DP, Somasekhar N, Prasad JS, Kirti PB, Transgenic tobacco plants constitutively
14 expressing *Arabidopsis* NPR1 show enhanced resistance to root-knot nematode
15 *Meloidogyne incognita*. *BMC Res Notes*; [http://www.biomedcentral.com/1756-](http://www.biomedcentral.com/1756-0500/4/231)
16 [0500/4/231](http://www.biomedcentral.com/1756-0500/4/231); 231 (2011). *Theor Appl Genet* **121** :237-248 (2010).
- 17 72 Gowen SR, Integrated management of root-knot nematodes on vegetables in Kenya, in
18 Final Technical Report (1 October 1999-30 September 2002), Reading, UK, DFID
19 R7472 Crop Protection Programme, pp. 52 (2002).
- 20 73 Gowen SR, Promotion of sustainable approaches for the management of root-knot
21 nematodes of vegetables in Kenya, in Final Report, Crop Protection Programme,
22 Reading, UK, DFID, pp. 39 (2005).
- 23 74 Sikora, RA and Fernández E, Nematode parasites of vegetables, in *Plant Parasitic*
24 *Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc, M, Sikora RA and
25 Bridge J. CAB International, Wallingford, UK, pp. 319-392 (2005).

- 1 75 Oruko L and Ndun'gu B, in *Final socio-economic report for the Peri-Urban Vegetable*
2 *IPM thematic cluster*, CABI/KARI/HRI/NRI/University of Reading/IACR Rothamsted
3 Collaborative Project, pp. 49 (2001).
- 4 76 Kanyagia ST, *A survey of vegetable nematodes in Kenya*. Horticulture Research and
5 Development Project, Field Document No. 8, Thika Horticultural Research Station,
6 Kenya (1980).
- 7 77 Maina MJ, Waceke JW and Kariuki JM, Plant parasitic nematodes associated with
8 cabbages in Nyandarua and Embu Districts, *12th KARI Biennial Scientific Conference*
9 *Proceedings*. Kenya Agricultural Research Institute, Nairobi, Kenya, pp. 613-619
10 (2010).
- 11 78 Otipa MJ, Kimenju JW, Mureithi JG and Kyalo G, Potential of rotation crops in managing
12 root knot (*Meloidogyne* spp.) nematodes in tomato. *African Journal of Horticultural*
13 *Science* 2 :111-123 (2009).
- 14 79 Carneiro RMDG, Almeida MRA Aldemiro Junior J, Mattos VS, Correa VR and Coyne
15 DL, Characterization of *Meloidogyne* spp. from Uganda and Tanzania. In *Proceedings*
16 *6th International Congress of Nematology, 4-9 May 2014, Cape Sun, Cape Town South*
17 *Africa*, p. 178 (2014). [Abstract]
- 18 80 Coyne DL, Fourie D and Moens M, Current and future management strategies in resource-
19 poor regions, in *Root-knot Nematodes*, ed. by Perry RN, Moens M and Starr J, CAB
20 International, Wallingford, UK, pp. 444-475 (2009).
- 21 81 Nchore SB, Waceke JW and Kariuki JM, Incidence and prevalence of root-knot nematode
22 *Meloidogyne* species in selected indigenous leafy vegetables in Kisii and Trans-mara
23 Counties of Kenya. *12th KARI Biennial Scientific Conference Proceedings*, Kenya
24 Agricultural Research Institute, Nairobi, Kenya, pp. 675-681 (2010).

- 1 82 Menete Z, Andrade M, Ricardo J, Naico A, Alvaro A, Munhaua B, Harahagazwe D,
2 Grineberg W and Low J, Mitigating Disaster and fighting Vitamin A deficiency with
3 new drought-tolerant, orange fleshed sweetpotato. Leaflet October 2011. International
4 Potato Center, Maputo, Mozambique (2011).
- 5 83 Sasser JN, Root-knot nematodes – a global menace to crop production. *Plant Disease* **64**
6 :35-41 (1980).
- 7 84 Mc Donald AH, De Waele D, Fourie, D, Daneel MS and Mahela PW, Evaluation of
8 various practices approaches and technologies for sustainable nematode management in
9 resource poor production sectors. *Programme and Abstracts of papers, 10th African*
10 *Crop Science Society Conference*, 10-13 October 2011, Maputo, Mozambique, p. 262
11 (2011).
- 12 85 John J, Better food security for Senegal's organic farmers. *Pesticides News* **84** :21-23
13 (2009).
- 14 86 Williamson S, Addressing agrochemical externalities and food security in food & fibre
15 supply chains: Pesticide action network UK and Africa experience from the field, in
16 *Aspects of Applied Biology 102*. Delivering food security with supply chain led
17 innovations: Understanding supply chains, providing food security, delivering choice,
18 ed. by Martindale, W. Association of Applied Biologists, Warwick, UK, pp. 65-73
19 (2010).
- 20 87 Davies KG and Spiegel Y, Root patho-systems nematology and biological control, in
21 *Biological Control of Plant-parasitic Nematodes: Building Coherence between*
22 *Microbial Ecology and Molecular Mechanisms*, ed. by Davies K and Spiegel Y,
23 Springer, pp. 291-303 (2011).
- 24 88 Luambano-Nyoni, N, *Potential of biocontrol agents and compatible cultural practices for*
25 *root-knot nematode management in tomato*, Thesis, Doctor of Philosophy in Plant

- 1 Nematology Department of Plant Science and Crop Protection, Faculty of Agriculture
2 University of Nairobi, Kenya, pp. 113-124 (2009).
- 3 89 Kilama P, Dubois T, Coyne D, Niere B, Gold CS and Adipala E, Antagonism of
4 *Paecilomyces* spp. Isolated from banana (*Musa* spp.) roots and rhizosphere against
5 *Radopholus similis*. *Nematropica* **37** :215-225 (2007).
- 6 90 Manzanilla-López RH, Esteves I, Finetti-Sialer MM, Hirsch PR, Ward E, Devonshire J,
7 Hidalgo L and Kerry BR, *Pochonia chlamydosporia*: advances and challenges to
8 improve its performance as a biological control agent of sedentary endo-parasitic
9 nematodes. *J Nematol* **45** :1-7 (2013).
- 10 91 Bourne JM, Karanja PK, Kalisz H, Karanja DK, Mauchline TH and Kerry BR, Incidence
11 and severity of damage caused by *Meloidogyne* spp. and isolation and screening of the
12 nematophagous fungus *Pochonia chlamydosporia* from some of the main vegetable
13 growing areas in Kenya. *International Journal of Nematology* **14** :111-122 (2004).
- 14 92 Ward E, Kerry BR, Manzanilla-López RH, Mutua G, Devonshire J and Hirsch PR, The
15 *Pochonia chlamydosporia* serine protease Gene *vcpI* is subject to regulation by carbon,
16 nitrogen and pH: implications for nematode biocontrol. *PLoS ONE* **7(4)** :e35657
17 (2012).
- 18 93 Luambano N. Kimenju, JW and Waceke JW, Colonisation of the biological control agent
19 *Pochonia chlamydosporia* on the rhizosphere of plants which are poor host to root-knot
20 nematodes. *10th African Crop Science Society Conference* 10-13 October 2011,
21 Maputo, Mozambique (2011).
- 22 94 Kibunja JW, Birgen JK, Kariuki GM and Coyne DL, The use of *Trichoderma* spp.
23 alongside resistant tomato varieties for root-knot nematode management under field
24 conditions in coastal Kenya. In *Proceedings 6th International Congress of Nematology*,
25 4-9 May 2014, Cape Sun, Cape Town South Africa, p. 267 (2014). [Abstract]

- 1 95 Hull R, Bosse M and Tzotzos G, Training for implementing risk assessment regulations
2 for the release of GM crops, in: *Aspects of Applied Biology 96. Agriculture: Africa's*
3 "engine for growth" – Plant Science & biotechnology hold the key, ed. by Bruce T,
4 Foyer C, Halford, N, Keys A, *et al.* Association of Applied Biologists, Warwick, UK,
5 pp. 1-8 (2010).
- 6 96 Atkinson HJ, Arinaitwe G, Kiggundu A and Tripathi L, Designing nematode resistant
7 crops from Africa: progress and constraint, in *Aspects of Applied Biology 96.*
8 *Agriculture: Africa's "engine for growth" – Plant Science & biotechnology hold the*
9 *key*, ed. by Bruce T, Foyer C, Halford N, Keys A, *et al.* Association of Applied
10 Biologists Warwick, pp. 119-122 (2010).
- 11 97 Wilson M and Kakouli-Duarte T, *Nematodes as Environmental Indicators*, CABI,
12 Wallingford, UK, (2009).
- 13 98 Thuo AK, Kimenju JW, Kariuki GM, Karuku GN, Wendot PK and Melakeberhan H,
14 Seasonal variations of nematode assemblages and diversity in vertisols, cambisols and
15 arenosols soil groups in Kenya. In *Proceedings 6th International Congress of*
16 *Nematology, 4-9 May 2014, Cape Sun, Cape Town South Africa*, p. 228 (2014).
17 [Abstract]
- 18 99 Costa SR, van der Putten, WH and Kerry BR, Microbial ecology and nematode control in
19 natural ecosystems, in *Biological Control of Plant-parasitic Nematodes: Building*
20 *Coherence between Microbial Ecology and Molecular Mechanisms*, ed. by Davies KG
21 and Spiegel Y, Springer, pp. 61-67, (2011).
- 22 100 Ferris H and Bongers T, Indices developed specifically for analysis of nematode
23 assemblages, in *Nematodes as Environmental Indicators*, ed. by Wilson M and
24 Kakouli-Duarte T, CABI, Wallingford, UK, pp. 124-145 (2009).

- 1 101 Neher DA and Darby BJ, General community indices that can be used for analysis of
2 nematode assemblage, in *Nematodes as Environmental Indicators*, ed. by Wilson M
3 and Kakouli-Duarte T, CABI, Wallingford, UK, pp. 107-123 (2009).
- 4 102 Nakacwa R, *Characterization of soil nematode diversity and community composition in a*
5 *field planted to transgenic bananas as a basis for biosafety assessment in Uganda*. MSc
6 Thesis submitted to the Directorate of Research and Graduate Training Makerere
7 University, Tanzania (2011).
- 8 103 Nguyen KB, Tesfamariam M, Gozel U, Gaugler R and Adams BJ, *Steinernema*
9 *yirgalemense* n. sp (Rhabditida: Steinernematidae) from Ethiopia *Nematology* **6** :839-
10 856 (2004).
- 11 104 Mweke AN, Kimenju JW, Seif AA, Mutitu EW and Mutua GK, Potential of sequential
12 cropping in the management of root knot nematodes in okra. *Asian J Plant Sci* **4** :399-
13 403 (2008).
- 14 105 Kariuki PM, Kariuki F, Kariuri GM and Coyne DL, The effect of different tomato
15 genotype on root-knot nematode-fusarium wilt complex in coastal Kenya. In
16 *Proceedings 6th International Congress of Nematology, 4-9 May 2014, Cape Sun, Cape*
17 *Town South Africa*, p. 266-267 (2014). [Abstract]
- 18 106 Muriuki LK, Kariuki GJ, Kinyua ZM, Gathu RK and Coyne DL, Management of root-
19 knot nematodes-bacterial wilt complex using resistant tomato varieties in coastal
20 Kenya. In *Proceedings 6th International Congress of Nematology, 4-9 May 2014, Cape*
21 *Sun, Cape Town South Africa*, p. 268 (2014). [Abstract]
- 22 107 Waceke JW, Minimizing pest damage on crops in SSA, *Falling Walls Conference*, 8-9th
23 November 2011, Berlin, Germany [Abstract].
- 24 108 Duncan A and Howell J, *Structural Adjustment and the African Farmer*. Overseas
25 Development Institute, London, UK (1992).

- 1 109 Jayne TS, Yamano T, Weber M, Tschirley D, Benfica R, Chapoto A and Zulu B,
2 Smallholder income and land distribution in Africa: Implications for poverty reduction
3 strategies. *Food Policy* **28** :253-275 (2003).
- 4 110 Stirling GR, *Biological Control of Plant-parasitic nematodes*, 2nd Edition, CABI,
5 Wallingford, UK (2014).
- 6 111 Smith J, Waage J, Woodhal JW, Bishop SJ and Spenece N, The challenge of providing
7 plant diagnostics services for Africa. *Eur J Plant Pathol* **121** :365-375 (2008).
- 8 112 Walsh S, *Leveraging Non-Governmental Organizations for Scale and Impact: Lessons*
9 *Learned from the Crop Crisis Control Project*. Catholic Relief Services, Bujumbura,
10 Burundi, 6 pp. (2008).
- 11 113 Oyekale AS, Gedion KE. Rural households' vulnerability to climate-related income
12 shocks and adaption options in central Malawi. *J Food Agric Environ* **10** :1505-1510
13 (2012).
- 14 114 UNFCCC (United Nations Framework Convention on Climate Change). *Climate*
15 *change: impacts, vulnerabilities and adaptation in developing countries* (2007).
- 16 115 Boko M, Niang I, Nyong A, Vogel C, Githeko A, Medany M, Osman-Elasha B, Tabo R
17 and Yanda P, Africa, in *Climate Change 2007: Impacts, Adaptation and Vulnerability.*
18 *Contribution of Working Group II to the Fourth Assessment Report of the*
19 *Intergovernmental Panel on Climate Change*, ed. Parry ML, Canziani OF, Palutikof JP,
20 van der Linden PJ and Hanson, CE. Cambridge University Press, Cambridge, UK, pp.
21 433-467 (2007).
- 22 116 Christensen JH, Hewitson B, Busuioc A, Chen A, Gao X, Held I, Jones R, Kolli RK,
23 Kwon W-T, Laprise R, Magaña Rueda V, Mearns L, Menéndez CG, Räisänen J, Rinke
24 A, Sarr A and Whetton P, Regional Climate Projections, in *Climate Change 2007: The*
25 *Physical Science Basis. Contribution of Working Group I to the Fourth Assessment*

- 1 *Report of the Intergovernmental Panel on Climate Change*, ed. by Solomon S, Qin D,
2 Manning M, Chen Z, Marquis M, Averyt KB, Tignor M and Miller HL, Cambridge
3 University Press. Cambridge, UK and New York, NY, USA, pp. 847-940 (2007).
- 4 117 Roberts J, Solving Kenya's food crisis, one indigenous crop at a time. *Solve Climate News* (2
5 Sep 2009), [http://solveclimateneeds.com/news/20090902/solving-kenya%E2%80%99s-](http://solveclimateneeds.com/news/20090902/solving-kenya%E2%80%99s-food-crisis-one-indigenous-crop-time?page=2)
6 [food-crisis-one-indigenous-crop-time?page=2](http://solveclimateneeds.com/news/20090902/solving-kenya%E2%80%99s-food-crisis-one-indigenous-crop-time?page=2) (2009).
- 7 118 Tubiello FN, Soussana J and Howden SM, Crop and pasture response to climate change.
8 *PNAS* **104** :19686-19690 (2007).
- 9 119 De Waele D and Elsen A, Challenges in tropical plant nematology. *Annu Rev*
10 *Phytopathol* **45** :457-485 (2007).
- 11 120 Kerry BR and De Leij FAAM, Key factors in the development of fungal agents for the
12 control of cyst and root-knot nematodes, in *Biological Control of Plant Diseases*, ed. by
13 Tjamos EC, Papavizas GC and Cook RJ, Plenum, London, pp. 139-144 (1992).
- 14 121 Sikora RA, Niere B and Kimenju JW, Endophytic microbial biodiversity and plant
15 nematode management in African agriculture, in *Biological Control in IPM systems in*
16 *Africa*, ed. by Neuenschwander P, Borgemeister C and Langewald J, CAB
17 International, Wallingford, UK, pp. 179-192 (2003).
- 18 122 Arim OJ, Waceke JW and Kimenju JW, Effects of Canavalia and Mucuna intercrops on
19 *P. zaeae* damage and yield of maize in subsistence agriculture. *Plant Soil* **284** :243-251
20 (2006).
- 21 123 Kimenju JW, Karanja NK and Nyongesa MW Diversity and abundance of nematodes in
22 agroecosystems of Kenya. *Journal of Tropical Microbiology* **3** :24-33 (2004).
- 23 124 Bridge J, Plant nematode pests of banana in East Africa with particular reference to
24 Tanzania, in *Les nematodes et le charancon du bananier: Situation et perspectives de*

- 1 *la recherche, Actes dun seminaire, Bujumbura (BDI) 1987/12/07-11. INIBAP:*
2 Montpellier (FRA) (1988).
- 3 125 Nsemwa LTH, Problems of banana weevil and nematodes in the Southern Highlands of
4 Tanzania. *Fruits* **46** :541-142 (1991).
- 5 126 Mbwana ASS, Waudu SW and Seshu-Reddy KV, Host range of the lesion nematode,
6 *Pratylenchus goodeyi*, highly destructive on highland bananas in East Africa.
7 *Proceedings of the 22nd International Symposium of the European Society of*
8 *Nematologists, Gent, Belgium* (abstract), p. 88 (1994).
- 9 127 Speijer PR and Fogain R, Musa and Esente nematode pest status in selected African
10 countries. Mobilizing IPM for sustainable banana production in Africa. *Proceedings of*
11 *a workshop on banana IPM held in Nelspriuit 1998/11/23-28. INIBAP, Montpellier,*
12 France, pp. 99-108 (1999).
- 13 128 Page SLJ, Mguni C and Sithole S, *Pests and diseases of crops in communal areas of*
14 *Zimbabwe*, ODA Technical Report, 1985, London, UK (1990).
- 15 129 Mai WF, Nematodes. *American Phytopathological Society*, pp. 93-101 (1980).
- 16 130 Franco J, Potato cyst nematodes: *Globodera* spp. *Technical Information Bulletin* **9** :21
17 International Potato Center, Lima, Peru, (1981).
- 18 131 Dobson H, Cooper J, Manyangarirwa W, Karuma J and Chiimba W, *Integrated*
19 *Vegetable Pest Management*. Natural Resources Institute, University of Greenwich, UK
20 (2002).
- 21 132 Muturi, J, Waceke W, Gichuki C and Runo SM, Use of isoenzymes phenotypes to
22 characterize the major root knot nematodes (*Meloidogyne* spp.) parasitizing indigenous
23 leafy vegetables in Kisii, *12th KARI Biennial Scientific Conference, Kenya, Kenya*
24 Agricultural Research Institute, Kenya, pp. 605-612 (2010).

- 1 133 Kavuluko JM, Waceke JW, Gichuki C. and Runo SM, Characterization of RKN
2 (*Meloidogyne* spp. from selected legumes in Mbeere District using isoenzymes
3 phenotypes, 12th KARI Biennial Scientific Conference, Kenya, Kenya Agricultural
4 Research Institute, Kenya, pp. 92-97 (2010).
- 5 134 Caveness FE, Root knot nematodes. *Annual Report IITA*, Ibadan, Nigeria, pp. 64-65
6 (1981).
- 7 135 McSorley R, Ohair SK and Parrado JL, Nematodes of cassava. *Nematropica* **13** :261-287
8 (1983).
- 9 136 Coyne DL and Namaganda JM, Root knot nematodes, *Meloidogyne* spp. incidence on
10 cassava in two areas of Uganda. *Roots Newsletter* **1** :2-3 (1994).
- 11 137 Campos VP and Villain L, Nematode parasites of coffee and cocoa, in *Plant Parasitic*
12 *Nematodes in Subtropical and Tropical Agriculture*, ed. by Luc M, Sikora RA and
13 Bridge J. CABI Publishing, Wallingford, UK, pp. 529-580 (2005).
- 14 138 Kagoda F, Dereraa J, Pangirayi, T and Coyne DL, Awareness of plant-parasitic
15 nematodes, and preferred maize varieties, among smallholder farmers in East and
16 Southern Uganda: implications for assessing nematode resistance breeding needs in
17 African maize. *International Journal of Pest Management* **56** :217-222 (2009).
- 18 139 Rashid MH, Yasmin L, Kibria MG, Millik AKMSR and Hossain SMM, Screening of
19 okra germplasm for resistance to yellow vein mosaic virus under field conditions. *Plant*
20 *Pathol J* **1** :61-62 (2002).
- 21 140 Kimani WNE, *Studies on fungal disease agents and their interactions with nematodes,*
22 *nutrition and genotypes on pyrethrum wilt development.* PhD thesis, University of
23 Nairobi, Kenya (2001).
- 24 141 Bond JP, McGawley EC and Hoy JW, Distribution of plant-parasitic nematodes on
25 sugarcane in Louisiana and efficacy of nematicides. *J Nematol* **32** :493-501 (2000).

- 1 142 Edgerton CW, Tims TC and Mills PJ, Relation of species of *Pythium* to the root-rot
2 disease of sugarcane. *Phytopathology* **19** :549-564 (1969).
- 3 143 Spaull VW, Nematodes associated with sugarcane in South Africa. *Phytophylactica* **13**
4 :175-179 (1981).
- 5 144 Nzioki HS, Survey on genera, distribution and abundance of plant-parasitic nematodes in
6 the South Nyanza sugarcane zone. *Kenya Sugar Research Foundation Technical*
7 *Bulletin* **1** :14-24 (2007).
- 8 145 Njuguna LK and Bridge J, Plant parasitic nematodes of Irish potatoes (*Solanum*
9 *tuberosum*) in Central Province and sweet potatoes (*Ipomoea batatas*) in Central,
10 Nyanza and Coast Provinces of Kenya. *International Journal of Nematology* **8** :21-26
11 (1998).
- 12 146 Kalele DN, Affokpon A, Coosemans J and Kimenju JW, Suppression of root knot
13 nematodes in tomato and cucumber using biological control agents *African Journal of*
14 *Horticulture* **3** :72-80 (2010).
- 15 147 Nono-Womdim R, Swai IS, Mroso LK, Chadha ML and Opena RT, Identification of
16 root-knot nematode species occurring on tomatoes in Tanzania and resistant lines for
17 their control. *Plant Dis* **86** :127-130 (2002).
- 18 148 Beije CM, Kanyagia ST, Muriuki SJN, Otieno EA, Seif AA and Whittle AM,
19 *Horticultural Crops Protection Handbook*, National Horticultural Research Station,
20 Thika, Kenya (1984).
- 21 149 Sutherland JA and Kibata G, Technical Report II KARI/ODA Crop Protection Project,
22 National Agricultural Research Laboratories (1993).
- 23