

Citation for published version:

Christopher R. Gelino, et al, 'WISEP J061135.13-041024.0 AB: A J-BAND FLUX REVERSAL BINARY AT THE L/T TRANSITION', *The Astronomical Journal*, Vol. 148 (6), July 2014.

DOI:

<http://dx.doi.org/10.1088/0004-6256/148/1/6>

Document Version:

This is the Published version.

Copyright and Reuse:

© 2014 The American Astronomical Society.

Content in the UH Research Archive is made available for personal research, educational, and non-commercial purposes only. Unless otherwise stated, all content is protected by copyright, and in the absence of an open license, permissions for further re-use should be sought from the publisher, the author, or other copyright holder.

Enquiries

If you believe this document infringes copyright, please contact Research & Scholarly Communications at rsc@herts.ac.uk

WISEP J061135.13–041024.0 AB: A *J*-BAND FLUX REVERSAL BINARY AT THE L/T TRANSITION

CHRISTOPHER R. GELINO^{1,2}, R. L. SMART³, FEDERICO MAROCCO⁴, J. DAVY KIRKPATRICK², MICHAEL C. CUSHING⁵,
GREGORY MACE⁶, RENE A. MENDEZ⁷, C. G. TINNEY^{8,9}, AND HUGH R. A. JONES⁴

¹ NASA Exoplanet Science Institute, Mail Code 100-22, California Institute of Technology, 770 South Wilson Avenue, Pasadena, CA 91125, USA

² Infrared Processing and Analysis Center, Mail Code 100-22, California Institute of Technology, 1200 E. California Boulevard, Pasadena, CA 91125, USA

³ Istituto Nazionale di Astrofisica, Osservatorio Astrofisico di Torino, Strada Osservatorio 20, I-10025 Pino Torinese, Italy

⁴ Centre for Astrophysics Research, University of Hertfordshire, Hatfield AL10 9AB, UK

⁵ Department of Physics and Astronomy, MS 111, University of Toledo, 2801 W. Bancroft Street, Toledo, OH 43606-3328, USA

⁶ Department of Physics and Astronomy, UCLA, 430 Portola Plaza, Box 951547, Los Angeles, CA 90095-1547, USA

⁷ Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile

⁸ School of Physics, University of New South Wales, Sydney, NSW 2052, Australia

⁹ Australian Centre for Astrobiology, University of New South Wales, Sydney, NSW 2052, Australia

Received 2013 July 26; accepted 2014 April 20; published 2014 May 28

ABSTRACT

We present Keck II laser guide star adaptive optics observations of the brown dwarf WISEP J061135.13–041024.0 showing it is a binary with a component separation of $0''.4$. This system is one of the six known resolved binaries in which the magnitude differences between the components show a reversal in sign between the *Y/J* band and the *H/K* bands. Deconvolution of the composite spectrum results in a best-fit binary solution with L9 and T1.5 components. We also present a preliminary parallax placing the system at a distance of 21.2 ± 1.3 pc. Using the distance and resolved magnitudes we are able to place WISEP J061135.13–041024.0 AB on a color–absolute magnitude diagram, showing that this system contributes to the well-known “*J*-band bump” and the components’ properties appear similar to other late-type L and early-type T dwarfs. Fitting our data to a set of cloudy atmosphere models suggests the system has an age >1 Gyr with WISE 0611–0410 A having an effective temperature (T_{eff}) of 1275–1325 K and mass of 64–65 M_{Jup} , and WISE 0611–0410 B having $T_{\text{eff}} = 1075$ –1115 K and mass 40–65 M_{Jup} .

Key words: binaries: general – brown dwarfs – stars: fundamental parameters – stars: individual (WISEP J061135.13–041024.0) – stars: low-mass

Online-only material: color figure

1. INTRODUCTION

The *H*-band flux of stellar objects changes by nearly 12 orders of magnitude from O stars down to the Y dwarfs (Kirkpatrick et al. 2012). The progression of absolute *H* magnitude (M_H) along this sequence is fairly smooth except for two major kinks.¹⁰ The first occurs in the early-type M dwarfs, where the formation of H_2 at these temperatures results in a short plateau in M_H (Mould 1976; Mould & Hyland 1976). The other kink lies at the transition between the L and T dwarfs. It is here that the absolute *H* magnitude shows a distinct flattening before turning downward at the early-mid type T dwarfs (Tinney et al. 2003). The source of this flattening, and the slight increase in absolute *J* at these same spectral types, is not understood, but could be related to clouds, as discussed below.

Clouds are required in the atmospheric models of L dwarfs in order to replicate the observed spectral and photometric trends. Conversely, models to match the observed properties of T dwarfs require the clouds to be completely dissipated, leaving the atmospheres clear of condensates. The transition from the cloudy L dwarfs to the clear T dwarfs is not well understood and has been notoriously difficult to model. A gradual lowering of a cloud deck through the L to T cooling sequence using a single sedimentation efficiency (f_{sed} ; Ackerman & Marley 2001) produces a shift in the near-IR magnitudes that is too slow to account for the observed colors and absolute magnitudes of L and T dwarfs (Knapp et al. 2004). However, if f_{sed} increases

from ~ 2 for the L dwarfs to infinity for the T dwarfs, giving the T dwarfs cloud-free atmospheres, then the bump in absolute *J* magnitude can be modeled. Conceptually, the change in f_{sed} means that condensates in the L dwarfs would precipitate out of the clouds like a light rain shower. As the object cools to later spectral types, the clouds fall deeper in the atmosphere and the condensates rain out progressively harder and harder, until the clouds drop below the photosphere and the atmosphere appears cloud-free.

The rapid clearing of the clouds and transition from L dwarf to T dwarf can also be explained by holes in the clouds (Ackerman & Marley 2001; Burgasser et al. 2002; Marley et al. 2010; Skemer et al. 2013). Small holes in the cloud deck allow hotter flux from deeper layers to emerge, analogous to Jupiter’s 5 μm “hot-spots” (Westphal et al. 1974). The presence of the holes results in a slightly brighter luminosity and bluer *J–K* color. The coverage area of the holes would rapidly progress from 0% for the cloudy L dwarfs to 100% for the clear T dwarfs.

In addition to the bump in absolute *J* magnitude, the L-T transition region is also characterized by a resurgence in the strength of FeH absorption at 9896 Å (Burgasser et al. 2002). This feature becomes weaker through the L dwarf sequence, nearly disappearing in L8 dwarfs, but then reappears in the mid-T dwarfs. Both the “sudden downpour” and the “cloud disruption” models can account for this resurgence in the FeH feature. In the sudden downpour model, the increase in f_{sed} means that the Fe cloud becomes thinner and drops deeper into the atmosphere. Consequently, the amount of FeH that is visible increases, thereby increasing the strength of the absorption feature (Cushing et al. 2008). Conversely, the holes in the cloud

¹⁰ There could be a third kink in this sequence in the Y dwarfs (Kirkpatrick et al. 2012).

Table 1
1.0–1.3 μm Flux Reversal Binaries

Binary	SpT _A	SpT _B	Reference
WISEP J061135.13–041024.0 AB	L9	T1.5	1
SDSS J102109.69–030420.1AB	T1	T4	2
WISE J104915.57–531906.1AB	L7.5	T0.5	3
2MASS J14044941–3159329AB	T1 \pm 1	T5 \pm 1	4
SDSS J153417.05+161546.1AB	T1.5	T5.5	5
2MASS J17281150+3948593AB	L5	L6.5	6, 7

References. (1) This work; (2) Burgasser et al. 2006b; (3) Burgasser et al. 2013; (4) Looper et al. 2008; (5) Liu et al. 2006; (6) Gizis et al. 2003; (7) Burgasser et al. 2010.

disruption model allows gaseous FeH below the cloud deck to become visible (Burgasser et al. 2002), which results in stronger FeH features in the observed spectra of early-T dwarfs.

Although some objects on the bright end of the bump may be unresolved binaries (Burrows et al. 2006), these “crypto-binaries” cannot account for all objects in the bump. Several binary systems, typically consisting of a late-L to early-T dwarf primary and an early- to mid-T dwarf secondary, have been discovered in which the secondary component is brighter than the primary in the 1.0–1.3 μm region (see Table 1). These systems indicate that the brightening in *J*-band from late-type L to early-type T is an intrinsic property of these objects and not the result of some selection effect. For example, the theory that the apparent brightening in the color–magnitude diagram is caused by objects with widely varying gravities (Tsuji & Nakajima 2003) can be discounted because such a scenario requires objects at vastly different ages, which is not likely in these presumably coeval binary systems.

In this paper we present the sixth binary showing a flux reversal in the 1.0–1.3 μm region. WISEP J061135.13–041024.0 (WISE 0611–0410, hereafter) was first presented in Kirkpatrick et al. (2011) as a T0 with a spectro-photometric distance of 17.7 pc. It was discovered in a *Wide-field Infrared Survey Explorer* (WISE; Wright et al. 2010) search for bright sources with brown dwarf-like colors and no counterpart in the Two Micron All Sky Survey catalog (2MASS; Skrutskie et al. 2006). Such properties are indicative of a source that either has a large proper motion or one with a very red *J* – *W2* color. In the case of WISE 0611–0410 the source *is* present in the 2MASS catalog (see Table 2 for a summary of the photometry of this object), but its position has moved by more than 3'' between the two surveys. Follow-up observations discussed here have accurately measured the proper motion and provide a first determination of the parallax. We also show resolved photometry and spectral deconvolution that clearly demonstrate the flux reversal in the 1.0–1.3 μm region.

2. OBSERVATIONS

2.1. NTT

Parallax observations were carried out using the European Southern Observatory 3.5 m New Technology Telescope (NTT) and its infrared spectrograph and imaging camera SofI (Son of ISAAC; Moorwood et al. 1998) as part of program 186.C-0756: *NTT Parallaxes of Southern Extremely Cool objects*. The observing, analysis, and reduction procedures are described in detail in Smart et al. (2013) and are only briefly summarized here. The observations were obtained in SofI’s large field mode, providing a 4.9×4.9 field with 0.288 pixel^{-1} in

Table 2
WISE 0611–0410 Photometry

Property	Value	Reference
Spectral type (near-IR)	T0	4
2MASS <i>J</i>	15.489 ± 0.055	1
2MASS <i>H</i>	14.645 ± 0.048	1
2MASS <i>K_s</i>	14.221 ± 0.070	1
MKO <i>J</i>	15.398 ± 0.006	3
MKO <i>H</i>	14.743 ± 0.005	3
MKO <i>K</i>	14.292 ± 0.005	3
IRAC 3.6 μm	13.069 ± 0.017	4
IRAC 4.5 μm	12.924 ± 0.017	4
WISE <i>W1</i> ^a	13.559 ± 0.026	5
WISE <i>W2</i> ^a	12.920 ± 0.027	5
WISE <i>W3</i> ^a	11.926 ± 0.282	5
WISE <i>W4</i> ^a	> 8.619	5

Note.

^a Its designation in the AllWISE Source Catalog is WISEA J061135.13–041024.1.

References. (1) 2MASS; (2) this work; (3) UKIDSS; (4) Kirkpatrick et al. 2011; (5) AllWISE Source Catalog (Cutri et al. 2013).

Table 3
NIRC2 Observing Log

Date (UT)	Filter ^a	Exposure (s)	FWHM (mas)	Strehl (%)
2012 Jan 13	<i>H</i>	1800	147	0.3
2012 Apr 15	<i>J</i>	180	75	2.5
	<i>H</i>	180	70	6.3
	<i>K_s</i>	180	67	14.4
2012 Sep 6 ^b	<i>H</i>	30	100	2.0
2012 Nov 29	<i>H</i>	360	69	5.3
2013 Sep 21	<i>Y</i>	360	129	0.7
	<i>K</i>	360	98	9.2

Notes.

^a All filters are on the MKO system (Tokunaga et al. 2002).

^b Only a single image was obtained on this date.

the *J* band. Each epoch consists of 18 dithered observations of 4×30 s at each pointing for a total exposure time of 36 minutes. The observations were combined using the *jitter* routine of the *Eclipse* (Devillard 1997, version 5.0) package. Objects in the field were found and centroided using the Cambridge Astrometry Survey Units *imcore* maximum likelihood barycenter software (CASUTOOLS,¹¹ v. 1.0.21).

During nights of exceptional seeing, one of us (R.A.M.), noted that the point-spread function of WISE 0611–0410 was slightly elliptical compared to other objects in the field. This prompted the high-resolution follow-up and spectral analysis discussed here.

2.2. Keck II/NIRC2

High resolution images of WISE 0611–0410 were obtained using the Keck II laser guide star adaptive optics (LGS-AO) system (Wizinowich et al. 2006; van Dam et al. 2006) with NIRC2 on the nights of 2012 January 13, 2012 April 15, 2012 September 6, 2012 November 29, and 2013 September 21 (see Table 3). Our target was too faint in *R* band to serve as a tip-tilt reference star for the wavefront corrector, so we used the

¹¹ <http://apm49.ast.cam.ac.uk/surveys-projects/software-release>




Figure 1. Keck LGS-AO NIRC2 images of WISE 0611–0410 AB obtained on 2012 April 15 (bottom row) and on 2013 September 21 (top row). The images are shown with north up and east to the left.

USNO-B star 0858–0074221 (Monet et al. 2003) with $R = 16.5$ and located $35''$ from our target. Our interest was focused on the target and its immediate vicinity, so we used the narrow camera mode with a nominal pixel scale of $10 \text{ mas pixel}^{-1}$ and single-image field of view of $10''$. We used a three-point dither pattern to avoid the high noise levels in the lower-left quadrant of the array. This pattern was repeated with different offsets to build up a longer exposure. The Mauna Kea Observatories (MKO; Tokunaga et al. 2002) Y , J , H , K , and K_s filters were used on one or more nights (Table 3).

The images were reduced using scripts in the IDL environment. A dark frame was first subtracted from each science frame. Then, a sky frame, created from the median average of all images acquired for WISE 0611–0410 exclusive of the frame being reduced, was subtracted. The sky-subtracted frames were then divided by a dome flat. Finally, the images were shifted to align the target to a common location and a deep mosaic was created from the median average of the stack. For the single image obtained on 2012 November 29, we could only dark subtract and flat-field the science frame. Fortunately, since our interest in that data was to follow the relative motions of the objects, the lack of a sky frame did not impede our goals. Table 3 lists the total exposure times for each of the mosaics, as well as the FWHM and Strehl ratios for the objects in frames.

The Y , J , H , K , and K_s mosaics from the 2012 April 15 and 2013 September 21 nights are shown in Figure 1. As was suspected from the NTT data, this source is easily resolved into two components. Visual inspection of the images shows that the relative brightness changes sign between the Y & J and H , K , & K_s images. We discuss the implications of this below.

3. ANALYSIS

3.1. Parallax and Proper Motion

We have 20 observations of WISE 0611–0410 from 2010 December 23 through 2013 February 8. An observation in the middle is chosen as the master frame and the measured

Table 4
Parallax and Proper Motion of WISE 0611–0410

Parameter	Value
α, δ (J2000)	6:11:35.0, $-4:10:24.9$
Epoch (yr)	2013.1629
Absolute parallax (mas)	47.25 ± 3.22
Distance (pc)	21.2 ± 1.3
μ_α (mas yr^{-1})	83.70 ± 2.50
μ_δ (mas yr^{-1})	-279.13 ± 2.02
Relative to absolute correction (mas)	0.84
Duration of observations (yr)	2.18
Reference stars, no. of observations	291, 19

coordinates transformed to a standard system using 2MASS objects in the field. All other observations are transformed to this standard system using all common objects and a simple linear transformation. Once all frames are on the same system, the motion of the target is fit for the astrometric parameters: position, relative proper motion, and relative parallax. This procedure is iterated while removing reference stars, which have brightnesses straddling that of WISE 0611–0410, with high scatter about the transformation. One epoch with an anomalously large residual ($\sim 3\sigma$) compared to the best fit was also dropped. After this iteration a correction of the relative parallax to an absolute value is calculated using the galaxy model of Mendez & van Altena (1996). The resulting astrometric motion of WISE 0611–0410 across the sky from these 19 epochs, along with our fit, is shown in Figure 2 and the astrometric parameters are produced in Table 4.

3.2. Resolved Photometry

Our Keck imaging easily resolves WISE 0611–0410 into its component sources, allowing for simple calculation of the magnitude differences in the five filters. To derive the MKO-based photometry of the two components, we need MKO magnitudes for the composite source in those filters. MKO J , H ,




Figure 2. Apparent motion of WISE 0611–0410 AB on the sky over the 19 NTT epochs. The offsets are relative to the middle observation. Also drawn is the best-fit curve to the parallax and proper motion.

Table 5
Properties of WISE 0611–0410 AB

Parameter ^a	A	B	Δ
MKO Y (mag)	-0.40 ± 0.01
MKO J (mag)	16.31 ± 0.01	16.01 ± 0.01	-0.30 ± 0.01
MKO H (mag)	15.43 ± 0.01	15.57 ± 0.01	$+0.14 \pm 0.01$
MKO K_s (mag)	$+0.66 \pm 0.01$
MKO K (mag)	14.72 ± 0.01	15.50 ± 0.01	$+0.78 \pm 0.01$
MKO $J-K$ (mag)	1.59 ± 0.01	0.51 ± 0.01	...
Est. Sp. Type	L9	T1.5	...

Note. ^a All magnitudes are apparent magnitudes.

and K images of WISE 0611–0410 were obtained by the UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007) Galactic Clusters Survey. The H and K data were obtained from the public Data Release 9; J -band data comes from unreleased images observed on 2012 December 9. These images show an elliptical source, similar to what is seen in the NTT data. No MKO Y or K_s images of this field have been obtained. This UKIDSS photometry, as well as the 2MASS photometry, is given in Table 2.

The separation between the two components is large enough that basic aperture photometry can be used to measure the magnitude differences. We use the data from 2012 April 15 and 2013 September 21 for the photometry since they are of the highest quality and include the largest set of filters. The photometry is performed on the mosaicked images (Figure 1) and is presented in Table 5. We denote the source that is fainter at Y and J as component A in this system.

3.3. Resolved Astrometry and Companionship

We have observed WISE 0611–0410 AB using high-resolution imaging on five different epochs with the purposes of confirming a common proper motion between the sources and searching for any orbital motion. For a given epoch, we measure




Figure 3. Motion of WISE 0611–0410 B with respect to WISE 0611–0410 A from the NIRC2 data. The inset shows the five epochs in more detail. We do not detect any significant motion of the companion, relative to component A, throughout our NIRC2 observations.

Table 6
NIRC2 Astrometry

Date (UT)	Measured		Predicted if BG Object	
	$\Delta\alpha \cos(\delta)$ (mas)	$\Delta\delta$ (mas)	$\Delta\alpha \cos(\delta)$ (mas)	$\Delta\delta$ (mas)
2012 Jan 13	-177.2 ± 15.8	346.8 ± 5.8	-177.2	346.8
2012 Apr 15	-184.4 ± 6.5	353.4 ± 3.2	-205.7	424.3
2012 Sep 6	-180 ± 10	349 ± 10	-234.3	530.1
2012 Nov 29	-182.5 ± 3.7	346.6 ± 4.9	-256.1	591.9
2013 Sep 21	-181.8 ± 3.4	338.6 ± 1.9	-322.6	808.6

the separation in R.A. and decl. from source A to source B on all of the individual images. We then take the average separations from the measurements and assign the standard deviation as the error in those quantities. Since the 2012 September 6 epoch only contained a single, good quality image, we conservatively set the separation error in each axis to be 10 mas (≈ 1 pixel).

The final measurements are shown in Table 6 and graphically in Figure 3. If source B was a background object, then its position would have changed by 480 mas over the course of the NIRC2 observations. A positional offset that large is easily detectable in these high quality observations. However, there is no significant change in the relative position of source B over the five epochs. Therefore, we can confidently confirm that these sources are physically bound and, additionally, no orbital motion is yet detected. The parallax and proper motion fit also do not show any evidence for binary motion over more than 2 yr of NTT observations.

3.4. Composite Spectrum Deconvolution

To determine the spectral types of the individual components of WISE 0611–0410 AB we used a three-step deconvolution procedure, similar to that described in Day-Jones et al. (2013). First we fit the entire spectrum ($0.85\text{--}2.45 \mu\text{m}$) of our target




Figure 4. Near-IR spectrum of WISE 0611–0410 AB from Kirkpatrick et al. (2011; black line). Also shown are the spectra of the template L9 dwarf SDSS J085234.90+472035.0 (blue line; Burgasser et al. 2010) and the T1.5 dwarf SDSS J175024.01+422237.8 (green line; Burgasser et al. 2010), whose convolution (red line) provides the best fit to the observed WISE 0611–0410 AB spectrum. The spectral regions used in the binary fitting are denoted by the dotted lines. The flux reversal between 1.0 and 1.3 μm observed in the NIRC2 photometry is also evident in the comparison of the template spectra.




Figure 6. Same as Figure 5 except for $\log g = 5$ (cm s^{-2}).




Figure 7. Same as Figure 5 except for $\log g = 5.477$ (cm s^{-2}).




Figure 5. Absolute MKO J magnitude as a function of MKO $J-K$ for a suite of models with $\log g = 4.477$ (cm s^{-2}). The atmosphere models include $f_{\text{sed}} = 1, 2, 3, 4$ with equilibrium chemistry and three levels of non-equilibrium chemistry mixing ($\log K_{zz} = 2, 4, 6$). Effective temperatures vary from model to model but generally range from 500–2400 K for the equilibrium chemistry models and 900–1800 K for the $\log K_{zz} = 6$ models. For reference, the locations of the $T_{\text{eff}} = 1500, 1300,$ and 1000 K models are plotted with diamonds (top to bottom).

(Kirkpatrick et al. 2011) with the near-infrared spectroscopic standards defined in Kirkpatrick et al. (2010) and Burgasser et al. (2006a) to determine the best-fit standard via χ^2 minimization. The best-fit standard selected this way is SDSS

J120747.17+024424.8 with spectral type T0, identical to the spectral type found by Kirkpatrick et al. (2011).

Then we repeated the fit using a set of synthetic binaries. Synthetic binaries were created combining the template spectra taken from the SpeX-Prism library,¹² scaling them to the appropriate relative flux level using the spectral type–absolute magnitude calibration presented in Marocco et al. (2010). We selected only those binaries giving a lower χ^2 compared to the standard template. Among those binary pairs, the best-fit binary was found by only fitting three regions of the spectrum: 1.10–1.25 μm (containing CH_4 and H_2O absorption bands), 1.55–1.75 μm (containing the CH_4 absorption band), and 2.10–2.35 μm (containing the CO absorption band). The features within these intervals change significantly at the transition between L and T types, and are therefore the most suitable to identify and deconvolve unresolved L/T pairs. The best-fit template identified with this procedure (see Figure 4) consists of a L9 dwarf (SDSS J085234.90+472035.0) and a T1.5 dwarf (SDSS J175024.01+422237.8).

¹² <http://pono.ucsd.edu/~adam/browndwarfs/speXprism/>




Figure 8. Contours of model fitting $\log \chi^2$ as a function of T_{eff} and $\log g$ for the 16 combinations of f_{sed} and $\log K_{\text{zz}}$ for WISE 0611–0410 A. The dot-dashed lines denote $f_{\text{sed}} = 2$, $\log K_{\text{zz}} = 0$ objects with ages 0.1, 1, and 6 Gyr (left to right). The upturn in T_{eff} from ≈ 1250 K to ≈ 1450 K for the 0.1 Gyr model is caused by deuterium fusion in the cores of these objects. The contour levels and associated colors are the same in all panels. The $\log K_{\text{zz}} = 0$ model represents the equilibrium chemistry case. Note that the gravities for the $f_{\text{sed}} = 3$ and 4, $\log K_{\text{zz}} = 6$ do not span as wide of a range as the other model cases.

We tested the significance of our deconvolution using an F -test to compare the result of the fit with synthetic binaries against the fit with the standard template alone (Burgasser et al. 2010). We use an F distribution threshold of 99.5%, which translates to a critical value (η_{crit}) of 1.41. If the ratio of the two χ^2 (defined as η) is larger than η_{crit} , then the combined template provides a better fit at the 99.5% confidence level. We obtained $\eta = 1.73$ for the binary fit using L9 and T1.5 templates and, therefore, assign those spectral types to the two components.

3.5. Discussion

3.5.1. Comparison to Cloudy Models

A true test to firmly pin down the atmospheric properties of these brown dwarfs requires high-resolution, high signal-to-

noise ratio spectroscopy. With such data it would be possible to distinguish the effects of gravity and non-equilibrium chemistry that are discernible in the models. With no resolved spectroscopy we will attempt to estimate the physical properties of WISE 0611–0410 AB using our measured photometry and distance.

We use the atmosphere models of Saumon & Marley (2008) and Stephens et al. (2009) for comparison to our binary. These models incorporate a variety of cloud sedimentation efficiencies (f_{sed} ; Ackerman & Marley 2001) that dictate the size of the condensate particles and the thickness of the cloud deck. Small values of f_{sed} produce thick clouds comprised of small particles, whereas large values of f_{sed} produce thin clouds with large particles. These models also include a vertical mixing component through the eddy diffusion coefficient, K_{zz} . This vertical mixing can throw the observable abundances




Figure 9. Same as Figure 8 except for WISE 0611–0410 B.

of molecules such as CO, CH₄, H₂O, NH₃, and N₂ out of chemical equilibrium by dredging up molecules favored at high temperatures and pressures (e.g., N₂ and CO) and mixing them higher in the atmosphere, where cooler temperature molecules (e.g., CH₄, H₂O, NH₃) are favored. If the replenishment of the “hot” molecules is faster than the reaction to convert them to the “cold” molecules, then there will be an over-abundance of CO compared to CH₄, for example. Values of K_{zz} are typically 10^4 – 10^6 cm² s⁻¹; in our model suite $\log K_{zz}$ is “0” (representing the chemical equilibrium case and *not* $K_{zz} = 1$), 2, 4, and 6.

It is important to note that the models we are utilizing are atmosphere models computed at a fixed grid of $\log g$ and T_{eff} . The evolutionary models of Saumon & Marley (2008) are then used to compute the mass. These evolutionary models are computed for clear atmospheres (not used in our analysis) and cloudy models with $f_{\text{sed}} = 2$. The masses and ages for other values of f_{sed} are approximate and based on $f_{\text{sed}} = 2$ models with

similar T_{eff} and $\log g$. Furthermore, although we show models with $\log g$ as high as 5.477 (cm s⁻²), these models produce objects that are physically impossible (Saumon & Marley 2008). In fact, the maximum value of $\log g$ for a cloudy set of models is 5.38 (cm s⁻²).

Figures 5, 6, and 7 show M_J as a function of J – K for models at a variety of effective temperatures, $\log K_{zz}$, and f_{sed} with $\log g = 4.477$, 5.0, and 5.477 (cm s⁻²). As shown in these figures, increasing the model gravity tends to shift absolute magnitudes to fainter values and the J – K color to the blue. Likewise, models with a higher sedimentation efficiency are bluer in J – K than lower f_{sed} models since the high f_{sed} models more closely resemble cloud-free atmospheres. Finally, increasing K_{zz} tends to produce redder J – K colors while leaving M_J relatively unchanged.

Since the grid of T_{eff} and gravities can be somewhat coarse for a given value of f_{sed} and $\log K_{zz}$ we have constructed

Table 7
Best-fit Models

	WISE 0611–0410 A	WISE 0611–0410 B
χ^2	17.7 ± 4.8	9.8 ± 5.5
Mass (M_{Jup})	64.8 ± 0.4	52.1 ± 12.0
T_{eff} (K)	1300 ± 26	1096 ± 20
$\log g$ (cm s^{-2})	5.3 ± 0.1	5.1 ± 0.1
f_{sed}	2	4
$\log K_{zz}$	0	6
Approx. age (Gyr)	1.5–4.0	1.0–2.0

a two-dimensional interpolation using steps of 10 K and 0.1 (cm s^{-2}) for the T_{eff} and $\log g$ axes, respectively. We acknowledge that this interpolated grid of models may not reproduce a model computed at each T_{eff} and $\log g$, but feel that the interpolation does a sufficient job of estimating the actual model properties.

To find the best-fitting model, we minimize χ^2 as given in the following equation:

$$\chi^2 = \sum_i^{J,H,K} \frac{(M_{i,\text{obs}} - M_{i,\text{model}})^2}{\sigma^2(M_{i,\text{obs}})}, \quad (1)$$

where $M_{i,\text{obs}}$ and $\sigma^2(M_{i,\text{obs}})$ are the absolute magnitude and the standard deviation of the observed absolute magnitude, respectively, and $M_{i,\text{model}}$ is the absolute magnitude from the model. The χ^2 minimization is performed for each discrete set of f_{sed} and K_{zz} models. Errors in the best-fit parameters (mass, T_{eff} , and $\log g$) are computed by randomly and independently varying the J , H , and K absolute magnitudes over the range expected from their errors.

The results of the χ^2 minimization calculations as a function of T_{eff} and $\log g$ are shown in Figures 8 and 9 for each set of K_{zz} and f_{sed} model. Also shown on these plots are the loci of $f_{\text{sed}} = 2$ models having ages of 0.1, 1, and 6 Gyr. In general, WISE 0611–0410 A (Figure 8) is fit well by $f_{\text{sed}} = 2$ models having relatively high gravity ($\log g \gtrsim 5$ [cm s^{-2}]) and $T_{\text{eff}} \approx 1200$ –1300 K. The corresponding mass for these models is 50–65 M_{Jup} with ages > 1 Gyr.

Conversely, the ranges of $\log g$ and T_{eff} of the best-fitting models for WISE 0611–0410 B greatly depend on the cloud and mixing properties (Figure 9). Strictly speaking, the best-fitting models are with $f_{\text{sed}} = 3$ and very low gravity. The ages inferred from these models (< 1 Gyr) are inconsistent with the ages for the WISE 0611–0410 A models. The $f_{\text{sed}} = 4$ models that provide a good fit to WISE 0611–0410 B have larger gravities and older ages, but do not fit the data quite as well as the $f_{\text{sed}} = 3$ models. To determine which set of models are a better physical fit to WISE 0611–0410 AB, we need to look at other observables that constrain the age of the system.

The space motion of WISE 0611–0410 AB effectively rules out membership of any known young moving group. Using the method from Marocco et al. (2010), and assuming a radial velocity range of ± 100 km s^{-1} , we find that WISE 0611–0410 AB has a low probability of $\sim 12\%$ of being younger than 500 Myr and $\sim 7\%$ of being younger than 100 Myr. The composite spectrum does not show any indicators that this system is very young, so we can assume that the age of the system is more than a few hundred million years old (Kirkpatrick 2005). This effectively rules out the low-gravity, $f_{\text{sed}} = 3$ models for WISE 0611–0410 B, so we adopt the higher gravity, older $f_{\text{sed}} = 4$ models.




Figure 10. Absolute MKO J magnitude as a function of MKO $J-K$ for the partly cloudy models of Marley et al. (2010) with cloud hole fractions of 0%, 25%, 50%, 75%, and 100%. The atmosphere models have effective temperatures 1000–2000 K and use equilibrium chemistry with $f_{\text{sed}} = 2$ and $\log g = 5$. WISE 0611–0410 A has properties consistent with a fully cloudy atmosphere. WISE 0611–0410 B is most closely matched to a model with $T_{\text{eff}} = 1150$ K and 25% fractional hole coverage.

(A color version of this figure is available in the online journal.)

The parameters for the best-fitting models with age > 1 Gyr are shown in Table 7. WISE 0611–0410 A is best fit by a model with $T_{\text{eff}} \approx 1300$ K, whereas WISE 0611–0410 B is best fit by a model with $T_{\text{eff}} \approx 1100$ K. The effective temperatures for both objects are largely consistent with those expected for late-type L dwarfs and early-type T dwarfs (Stephens et al. 2009). Likewise, the cloud properties are also not unusual for these types. Being a late-type L dwarf, WISE 0611–0410 A is expected to have thick clouds covering most of its atmosphere. Models with $f_{\text{sed}} = 2$ are generally the best types for fitting late-type L dwarfs (Saumon & Marley 2008). Early-mid T dwarfs require models that are less cloudy than the L dwarfs, making an $f_{\text{sed}} = 4$ model appropriate. However, an alternate model can also be used to characterize WISE 0611–0410 B (Section 3.5.2).

Finally, one interesting aspect of the model fitting is the very different levels of atmospheric mixing implied for the two objects. While this could represent a genuine physical difference between these objects, it is impossible to make this assessment based entirely on the broadband photometry in three filters. High resolution spectroscopy of both components is needed to truly ascertain if either object shows non-equilibrium chemistry.

3.5.2. Comparison to Partly Cloudy Models

WISE 0611–0410 B falls in the transition region between the late L dwarfs and the early T dwarfs, where models with partial cloud coverage can be used to describe the source. We use the models of Marley et al. (2010) to infer an alternate physical description of this source. These models, shown in Figure 10, compute the emitted flux for brown dwarfs with fixed masses and




Figure 11. Absolute MKO J (top), H (middle), and K (bottom) magnitudes as a function of near-IR spectral type. WISE 0611–0410 is denoted by the blue triangles. Other 1.0–1.3 μm flux reversal binaries are shown with red circles; binary components are connected by a red line. Components of other binaries are shown with magenta, inverted triangles. Brown dwarfs not known to be binaries are given by black stars. Data from the other L and T dwarfs are taken from the compilation of Dupuy & Liu (2012), and references therein) and Burgasser et al. (2013), from which young and metal poor objects have been excluded.

temperatures, but having different fractions of cloud coverage. All models have $f_{\text{sed}} = 2$, $\log g = 5.0$, and equilibrium chemistry. As expected, WISE 0611–0410 A is consistent with having a completely cloudy atmosphere (no holes in the cloud decks). WISE 0611–0410 B has $T_{\text{eff}} \approx 1200$ K and hole fraction = 25% for this set of models. If a different set of base models was used, then the best-fitting T_{eff} and hole fraction would be different (Marley et al. 2010), so caution should be used when implying a definite atmospheric structure based on this one case. Nonetheless, it is important to point out that the T_{eff} inferred by the partly cloudy models is consistent with the T_{eff} inferred from the cloudy models with varying degrees of f_{sed} and atmospheric mixing.




Figure 12. Absolute MKO J magnitude as a function of MKO $J-K$. Data and point types are the same as in Figure 11.

3.5.3. Comparison to Other Brown Dwarfs

The resolved photometry and parallax measured here allow us to compare WISE 0611–0410 AB to other brown dwarfs. In Figure 11 we use the absolute magnitudes and spectral types estimated from the spectrum deconvolution to compare WISE 0611–0410 AB to other L and T dwarfs compiled by Dupuy & Liu (2012, and references therein) and Burgasser et al. (2013). Both components appear to be normal with respect to similarly-typed brown dwarfs. However, since both spectral types are determined from the deconvolution of the composite spectrum, they should only be regarded as estimates. The color–magnitude diagram shown in Figure 12 bypasses the spectral type uncertainty by making the comparison based entirely on the well-measured photometry and parallax. The photometry of WISE 0611–0410 AB is well matched to the photometry of objects with similar spectral types and colors, indicating that WISE 0611–0410 AB does not have significantly different metallicity or cloud properties compared to most other brown dwarfs.

3.5.4. Orbit Estimations

In order to estimate the orbital period of this system we first need to estimate the orbital semimajor axis and the masses of the components. Since we have not detected any significant orbital motion of the secondary, we need to estimate the semimajor axis using a conversion factor from the projected separation to semimajor axis. We use the conversion given in Dupuy & Liu (2011) rather than the more commonly used conversion in Fischer & Marcy (1992) because the former is based on very low-mass star and brown dwarf binaries, whereas the latter is based on higher mass, stellar binaries. Dupuy & Liu (2011) compute several conversion factors based on the discovery method and binary type. For these observations, we use the “moderate discovery bias,” for which the inner working angle

of our observations is approximately half of the true semimajor axis. Using the factor for all very low-mass binaries ($a/\rho = 1.08$) and the projected separation from the most recent epoch (8.1 AU), we estimate the true semi-major axis to be 8.8 ± 0.5 AU. The mass estimates are done using the model fitting described in the previous sections and are presented in Table 7. The orbital period for the best-fitting model parameters is then 78 ± 8 yr.

4. CONCLUSIONS

We have discovered a new brown dwarf binary system that straddles the L/T dwarf transition. Resolved, differential photometry shows that the earlier-type primary is *fainter* than the cooler secondary in the *Y* and *J* bands, but brighter in *H* and *K_s*, making this one of six known systems that shows a flux reversal in the 1.0–1.3 μm region. Although we do not have resolved spectroscopy of the components, deconvolving the composite spectrum into two spectra indicates the best-fit spectral types are L9 and T1.5. We also present a new parallax of the system placing it at 21.2 ± 1.3 pc. Comparison to other L and T dwarfs shows that this system does not appear to have any unusual metallicity or age properties compared to the bulk of the field brown dwarfs, making it a good system to study for characterizing solivagant¹³ brown dwarfs. However, with an orbital period estimated at 78 yr, it will be some time before a good determination of the component masses can be made.

The authors acknowledge telescope operators Heather Hershey, Carolyn Jordan, Julie Renaud-Kim, and Cynthia Wilburn and instrument specialists Scott Dahm, Marc Kassis, and Luca Rizzi for their assistance during the Keck observations. We would also like to thank Chas Beichman for obtaining images of WISE 0611–0410 on 2012 November 29 and 2013 September 21, Mark Marley and Didier Saumon for providing the models, Mike Read for help in mining the UKIDSS database and useful discussions, and the anonymous referee for useful comments. Based partially on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere, Chile program 186.C-0756. R.L.S., H.R.A.J., and F.M. would like to acknowledge the Marie Curie 7th European Community Framework Programme grant no. 247593 Interpretation and Parameterization of Extremely Red COOL dwarfs (IPERCOOL) International Research Staff Exchange Scheme. R.A.M. acknowledges partial support from project PFB-06 CATA and from project IC120009 “Millennium Institute of Astrophysics (MAS)” of the Iniciativa Científica Milenio del Ministerio de Economía, Fomento y Turismo de Chile. C.G.T. acknowledges the support of ARC grants DP0774000 and DP130102695.

This publication makes use of data products from 2MASS, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology (Caltech), funded by the National Aeronautics and Space Administration (NASA) and the National Science Foundation. This publication makes use of data products from the *Wide-field Infrared Survey Explorer (WISE)*, which is a joint project of the University of California, Los Angeles, and

the Jet Propulsion Laboratory (JPL)/Caltech, and *NEOWISE*, which is a project of JPL/Caltech. *WISE* and *NEOWISE* are funded by NASA. This work was supported by a NASA Keck PI Data Award, administered by the NASA Exoplanet Science Institute. Some of the data presented herein were obtained at the W. M. Keck Observatory from telescope time allocated to NASA through the agency’s scientific partnership with the Caltech and the University of California. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

Facilities: Keck:II (NIRC2, LGS-AO), NTT (SofI)

REFERENCES

- Ackerman, A. S., & Marley, M. S. 2001, *ApJ*, **556**, 872
 Burgasser, A. J., Cruz, K. L., Cushing, M., et al. 2010, *ApJ*, **710**, 1142
 Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006a, *ApJ*, **637**, 1067
 Burgasser, A. J., Kirkpatrick, J. D., Cruz, K. L., et al. 2006b, *ApJS*, **166**, 585
 Burgasser, A. J., Marley, M. S., Ackerman, A. S., et al. 2002, *ApJL*, **571**, L151
 Burgasser, A. J., Sheppard, S. S., & Luhman, K. L. 2013, *ApJ*, **772**, 129
 Burrows, A., Sudarsky, D., & Hubeny, I. 2006, *ApJ*, **640**, 1063
 Cushing, M. C., Marley, M. S., Saumon, D., et al. 2008, *ApJ*, **678**, 1372
 Cutri, R. M., Wright, E. L., Conrow, T., et al. 2013, Explanatory Supplement to the AllWISE Data Release Products, <http://wise2.ipac.caltech.edu/docs/release/allwise/expsup/>
 Day-Jones, A. C., Marocco, F., Pinfield, D. J., et al. 2013, *MNRAS*, **430**, 1171
 Devillard, N. 1997, *Msngr*, **87**, 19
 Dupuy, T. J., & Liu, M. C. 2011, *ApJ*, **733**, 122
 Dupuy, T. J., & Liu, M. C. 2012, *ApJS*, **201**, 19
 Fischer, D. A., & Marcy, G. W. 1992, *ApJ*, **396**, 178
 Gizis, J. E., Reid, I. N., Knapp, G. R., et al. 2003, *AJ*, **125**, 3302
 Kirkpatrick, J. D. 2005, *ARA&A*, **43**, 195
 Kirkpatrick, J. D., Cushing, M. C., Gelino, C. R., et al. 2011, *ApJS*, **197**, 19
 Kirkpatrick, J. D., Gelino, C. R., Cushing, M. C., et al. 2012, *ApJ*, **753**, 156
 Kirkpatrick, J. D., Looper, D. L., Burgasser, A. J., et al. 2010, *ApJS*, **190**, 100
 Knapp, G. R., Leggett, S. K., Fan, X., et al. 2004, *AJ*, **127**, 3553
 Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, **379**, 1599
 Liu, M. C., Leggett, S. K., Golimowski, D. A., et al. 2006, *ApJ*, **647**, 1393
 Looper, D. L., Gelino, C. R., Burgasser, A. J., & Kirkpatrick, J. D. 2008, *ApJ*, **685**, 1183
 Marley, M. S., Saumon, D., & Goldblatt, C. 2010, *ApJL*, **723**, L117
 Marocco, F., Smart, R. L., Jones, H. R. A., et al. 2010, *A&A*, **524**, A38
 Mendez, R. A., & van Altena, W. F. 1996, *AJ*, **112**, 655
 Monet, D. G., Levine, S. E., Canzian, B., et al. 2003, *AJ*, **125**, 984
 Moorwood, A., Cuby, J.-G., & Lidman, C. 1998, *Msngr*, **91**, 9
 Mould, J. R. 1976, *A&A*, **48**, 443
 Mould, J. R., & Hyland, A. R. 1976, *ApJ*, **208**, 399
 Saumon, D., & Marley, M. S. 2008, *ApJ*, **689**, 1327
 Skemer, A. J., Marley, M. S., Hinz, P. M., et al. 2013, arXiv:1311.2085
 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, **131**, 1163
 Smart, R. L., Tinney, C. G., Bucciarelli, B., et al. 2013, *MNRAS*, **433**, 2054
 Stephens, D. C., Leggett, S. K., Cushing, M. C., et al. 2009, *ApJ*, **702**, 154
 Tinney, C. G., Burgasser, A. J., & Kirkpatrick, J. D. 2003, *AJ*, **126**, 975
 Tokunaga, A. T., Simons, D. A., & Vacca, W. D. 2002, *PASP*, **114**, 180
 Tsuji, T., & Nakajima, T. 2003, *ApJL*, **585**, L151
 van Dam, M. A., Bouchez, A. H., Le Mignant, D., et al. 2006, *PASP*, **118**, 310
 Westphal, J. A., Matthews, K., & Terrile, R. J. 1974, *ApJL*, **188**, L111
 Wizinowich, P. L., Le Mignant, D., Bouchez, A. H., et al. 2006, *PASP*, **118**, 297
 Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, **140**, 1868

¹³ Solivagant means “wandering alone” and is used to denote brown dwarfs that are isolated from other stars or brown dwarfs.