

# The electronic bands of CrD, CrH, MgD and MgH: application to the “deuterium test”

Ya. V. Pavlenko<sup>1,2,\*</sup>, G. J. Harris<sup>3</sup>, J. Tennyson<sup>3</sup>, H. R. A. Jones<sup>1</sup>,  
J. M. Brown<sup>4</sup>, C. Hill<sup>4</sup> and L.A.Yakovina<sup>1</sup>

<sup>1</sup>*Main Astronomical Observatory, Academy of Sciences of Ukraine, Golosiiv Woods, Kyiv-127, 03680 Ukraine*

<sup>2</sup>*Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, Hertfordshire AL10 9AB, UK*

<sup>3</sup>*Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT UK*

<sup>4</sup>*Oxford University, Physical & Theoretical Chemistry Laboratory, South Parks Rd, Oxford OX1 3QZ, UK*

## ABSTRACT

We compute opacities for the electronic molecular band systems  $A\ ^6\Sigma^+ - X\ ^6\Sigma^+$  of CrH and CrD, and  $A\ ^2\Pi - X\ ^2\Sigma^+$  of MgH and MgD. The opacities are computed by making use of existing spectroscopic constants for MgH and CrH. These constants are adjusted for the different reduced masses of MgD and CrD. Frank-Condon factors are used to provide intensities for the individual vibronic bands. These results are used in the computation of synthetic spectra between  $T_{\text{eff}} = 1800$  and 1200 K with an emphasis on the realisation of “deuterium test”, first proposed by Bejar et al. (1999) to distinguish brown dwarfs from planetary mass objects. We discuss the possible use of CrD and MgD electronic bands for the “deuterium test”. We find CrD to be the more promising of the two deuterides, potentially, the most useful bands of CrH/CrD are the  $\Delta v = +1$  and  $\Delta v = -1$  at 0.795 and 0.968  $\mu\text{m}$ .

**Key words:** stars: molecular spectra deuterium abundance – stars: fundamental parameters – stars: late-type – stars: brown dwarfs – stars: evolution – planets: evolution

\* E-mail: yp@mao.kiev.ua

## 1 INTRODUCTION

The “deuterium test” was suggested as a method of identifying planetary mass objects among cool objects (Béjar et al. 1999, Chabrier et al. 2000). In practise, it was proposed to search for absorption lines of molecules containing deuterium (HDO, CrD, FeD, etc.). Deuterium is burnt in stellar interiors via the fusion reaction  ${}^2\text{D}(p,\gamma){}^3\text{He}$  at temperatures ( $T > 8 \times 10^5$  K). The interiors of substellar objects with  $M < 13 M_J$ , where  $M_J$  is a Jovian mass ( $0.001 M_\odot$ ), do not reach temperatures high enough for deuterium to ignite (Saumon et al. 1996). As a result the deuterium abundance in the atmospheres of these objects is unchanged from the formation of these objects. This gives rise to the definition of a brown dwarf as an object which has insufficient mass to fuse  ${}^1\text{H}$  to  ${}^4\text{He}$ , but has sufficient mass to fuse  ${}^2\text{H}$  to  ${}^3\text{He}$ . By comparison, a planetary mass object has insufficient mass to ignite fusion of any sort (Saumon et al. 1996).

In higher-mass objects such as stars, deuterium burning is completed comparatively quickly ( $t = 1 - 3$  million years) during the evolution prior to the star’s period on the main sequence (D’Antona & Mazitelli 1998). The deuterium depletion rate depends on the mass of the star or brown dwarf and thus the “deuterium test” can be used in a number of ways.

- Discern planets from brown dwarfs in a population of low-mass objects.
- Determine the evolutionary status of young objects in open clusters with ages of several million years.
- Study the evolution of the abundance of deuterium in the atmospheres of low-mass substellar objects, a phenomenon which is poorly understood. The rate of depletion of deuterium depends upon rotation, magnetic field strength, and other parameters which affect the efficiency of convection in low mass objects.

Such investigations can usefully be combined with the “lithium test” proposed by Rebolo et al. (1992), Magazzú et al. (1993). For stars ( $M > 75 M_J$ ), the burning of lithium,  $\text{Li}(p,\alpha){}^4\text{He}$ , becomes efficient at evolutionary stages preceding the main sequence at interior temperatures of  $T > 2.5 \times 10^6$  K (D’Antona & Mazitelli 1998). The “lithium test” has been successfully applied to identify brown dwarfs in a population of ultracool dwarf stars (Rebolo et al. 1996, Basri 2000). The lithium test is relatively easily applied to M-dwarfs. The reason for this is that the resonance lines of neutral atomic lithium lie in the optical region of the spectrum, at  $0.6708 \mu\text{m}$ . One blemish with the “lithium test” is the severe blending of lithium lines with the background of TiO lines, nonetheless Li lines break through the

molecular background. In the spectra of cooler M and L dwarfs Ti atoms are depleted onto dust particles (Tsuji et al. 1996, Jones & Tsuji 1997, Pavlenko 1998) so TiO absorption is weakened. In the L dwarf regime the appearance of lithium lines contends with dust opacity and the wings of K I and Na I resonance lines (see Pavlenko et al. 2000 and references therein).

The realisation of the “deuterium test” is considerably more difficult. Conventionally, the deuterium abundance of hot objects is determined from analysis of multicomponent features on the background of the  $L_\alpha$   $0.1215 \mu\text{m}$  H I line seen in emission. Unfortunately, this method cannot be used in the case of ultracool dwarfs which are covered by a thick envelope of neutral hydrogen. Observations also yield H I emission lines ( $H_\alpha$   $0.6563 \mu\text{m}$ ) formed in the outermost layers of hot chromospheres or accretion disks around young stars. However, such a plasma is variable and may be polluted by interstellar material.

Taking into account these circumstances, it is logical to analyse the spectral lines of deuterated molecules formed in the photospheric layers of ultracool dwarfs. The first investigation and analysis of the combined spectra of  $\text{H}_2\text{O}$  /HDO were carried out by Chabrier et al. (2000) and Pavlenko (2002). Due to the change in mass and the breakdown of molecular symmetry the vibration-rotation bands of HDO in the mid-infrared spectrum shift with respect to the  $\text{H}_2\text{O}$  bands. There are a few spectral regions which can be used for detection of HDO lines in the IR spectra of ultracool dwarfs:  $3.5 - 4$ ,  $\sim 5$ ,  $6-7 \mu\text{m}$  (Pavlenko 2002). The main problem is that despite the shift in wavelength such HDO lines will be on a background of far stronger  $\text{H}_2\text{O}$  lines.

A possible alternative to HDO are the diatomic hydrides. Strong molecular bands of diatomic metal hydrides such as MgH and CrH can be observed in the optical spectrum of ultracool dwarf stars. The MgH band system  $A^2\Pi - X^2\Sigma^+$  can be observed at  $0.47-0.6 \mu\text{m}$ , and the CrH band system  $A^6\Sigma^+ - X^6\Sigma^+$  show absorption features at  $0.6-1.5 \mu\text{m}$ .

The molecules MgH and CrH have been known in astrophysics for a long time. MgH has been more extensively studied than CrH, because it can be observed in the spectra of G to M type stars. The dissociation energy of MgH is very low (1.285 eV) so lines of this molecule are very sensitive to the temperature and gravity variations in stellar atmospheres. MgH lines were used to determine temperatures in the atmospheres of cool giants (Wyller 1961) and the Sun (Sinha et al. 1979, Sinha & Joshi 1982), and for the determination of the surface gravity of stars (Bell & Gustafsson 1981, Bell et al. 1985, Berdyugina & Savanov 1992, Bonnel & Bell 1993).

The pure rotational spectrum of MgH and MgD radicals ( $X^2\Sigma^+$ ) in their ground state  $v=0$  and  $v=1$  vibrational modes has been studied by Ziurys et al. (1993). The first MgH linelist was computed by Kurucz (1993). Recently, more extensive studies of MgH transitions were performed by Weck et al. (2003, 2003a, 2003b) and Skory et al. (2003).

Although CrH has been known since Gaydon & Pearse (1937), its electronic spectrum remained relatively unstudied for many years. Engvold et al. (1980) identified lines of CrH in a spectrum of sunspots as formed by  $X^6\Sigma^+ - X^6\Sigma^+$  transitions. They used the results of studies of multiplicity of  $\Sigma$ -terms of CrH by Klehman & Uhler (1959) and O'Connor (1967). Later Ram, Jarman & Bernath (1993) performed a rotational analysis of 0–0 band of the  $A^6\Sigma^+ - X^6\Sigma^+$  electronic transition and obtained improved rotational constants for the  $v'=0$  vibrational state. Combining these results with those of Bauschlicher et al. (2001) and Lipus et al. (1991) for the vibrationally excited transitions, Burrows et al. (2002) computed an extended linelist for CrH.

Recently Shin et al. (2005) have measured radiative lifetimes of the  $v = 0,1$  levels of  $A^6\Sigma^+$  state of CrH. These measured lifetimes are about 16% – 45 % longer than those obtained by Burrows et al. (2002). These results provide evidence that the oscillator strengths of Burrows et al. (2002) should be corrected by a factor of 0.8 for at least the transitions to the  $v' = 0$ .

The submillimeter spectra of CrH and CrD formed by pure rotational transitions in the ground electronic state, have been observed in the laboratory by Halfen & Ziurys (2004). Electronic bands of MgD and CrD are likely to be located in the same spectral regions as the corresponding bands of MgH and CrH. In this paper we model the bands of these molecules to analyse the possibility of their use for the determination of the D/H ratio in the atmospheres of ultracool dwarfs.

In section 2 we present a description of the procedures used to compute the molecular bands of CrH, CrD, MgH and MgD. In section 3 we present the vibrational-rotational constants of MgH, MgD, CrH and CrD. In section 4 we present the results of the computation of molecular bands. In section 5 we discuss the possibility of using the electronic bands of diatomic molecules for a realisation of the “deuterium test”.

## 2 PROCEDURE OF COMPUTATIONS

### 2.1 Profile of electron bands of diatomic molecule

In this paper we compute the profile of the electronic band averaged over its rotational structure (Naersisian et al. 1989):

$$\int k_{\omega} d\omega = \frac{A q_{v'v''} S_e S_{j'j''} \omega_{j'j''}}{Q(T)} \times \exp\left(-\frac{hc(E_e'' + E_v'' + E_j'')}{kT}\right) \times \left[1 - \exp\left(-\frac{h\omega_{j'j''}}{kT}\right)\right], \quad (1)$$

where as symbols " label lower levels of transitions,  $A = 8\pi^3(3hc)^{-1}$ ,  $k$  is Boltzmann's constant and  $T$  is the temperature,  $\omega$  is a frequency and  $q_{v'v''}$  are Franck-Condon factors for the corresponding transition, the strength of transition  $S_e = S_e(0,0) \times \omega_{v'v''}/\omega_{0,0}$  (Schadee 1968),  $Q(T)$  is the partition function of the molecule.

For a given frequency the sum of the contribution from the  $P$ ,  $Q$  and  $R$  branches is:

$$k_{\omega} = \frac{A S_e F_e}{n\lambda Q(T)} \sum q_{v'v''} C_{vj} \sum |\lambda_j^v|, \quad (2)$$

where  $F_e = \exp(-hcE_e''/(kT))/\delta E_e$ ,

$C_{vj} = (\delta E_e + \delta E_v)(2j'' + 1) \exp(-hc(E_j'' + E_v''))$ ,

$\lambda = 1/\delta\omega$ .

This method of the computation of the profiles of molecular bands was successfully applied to the modelling of the bands of TiO in the spectra of ultracool dwarfs (Pavlenko 1997a, 1998a, Pavlenko 2000), and CN and C<sub>2</sub> bands in the spectra of evolved stars (Pavlenko et al. 2000a,c). It is worth noting that direct comparison of our JOLA synthetic spectra with more sophisticated “line by line” computations for linelists of a few molecules show good agreement (Pavlenko 1998a) both in the profiles of bands and positions of the band heads.

The JOLA approach has been used for the modelling of the CrH bands  $\Delta v = 0$  band system located at 0.8640  $\mu\text{m}$  in the spectra of ultracool dwarfs (Pavlenko 1999, Pavlenko et al. 2000). These bands are not considered for the deuterium test here, because the wavelengths of corresponding lines of CrH and CrD are nearly identical.

## 2.2 Computations of molecular constants of isotopic molecules

For a given electronic state, the energy levels of a diatomic molecule can, in general, be fitted by:

$$\begin{aligned}
 E_{e,v,N} = & T_e + \omega_e(v + 1/2) - \omega_e x_e(v + 1/2)^2 \\
 & + B_e N(N + 1) - \alpha_e N(N + 1)(v + 1/2) \\
 & - D_e(N(N + 1))^2 \\
 & + \beta_e(N(N + 1))^2(v + 1/2),
 \end{aligned} \tag{3}$$

where  $T_e$ ,  $\omega_e$ ,  $\omega_e x_e$ ,  $B_e$ ,  $\alpha_e$ ,  $D_e$  and  $\beta_e$  are the various spectroscopic constants,  $v$  and  $N$  are vibrational excitation and rotational quantum numbers, respectively. To a good approximation the electronic structure of the various isotopic analogues of a given molecule are the same. However, as the nuclei have different masses, the reduced mass and moment of inertia of the molecule are different, resulting in different rotation-vibration frequencies. For the case of an anharmonic oscillator and non-rigid rotator model of the molecule, the rotational-vibrational constants depend upon the ratio of reduced mass  $\mu_{ab} = m_a \times m_b / (m_a + m_b)$  of two isotopic molecules (see Wang & Xia 1996, Craybeal 1988).

The ratios of spectroscopic constants of two isotopic species, labelled as “o” and “i”, of reduced masses  $\mu_o$  and  $\mu_i$ , depend on the ratio  $\rho = (\mu_o/\mu_i)^{(1/2)}$ . So that the following approximations can be used:

$$(\omega_e)_i / (\omega_e)_o = \rho$$

$$(\omega_e x_e)_i / (\omega_e x_e)_o = \rho^2$$

$$(B_e)_i / (B_e)_o = \rho^2$$

$$(D_e)_i / (D_e)_o = \rho^4$$

$$(\alpha_e)_i / (\alpha_e)_o = \rho^3$$

**Table 1.** Reduced masses and ratios of reduced masses for molecules of interest.

Molecule	$\mu/u$	$\rho$
MgH	0.967481	0.721116
MgD	1.86051	
CrH	0.988438	0.713873
CrD	1.93958	

$$(\beta_e)_i/(\beta_e)_o = \rho^5$$

The reduced masses of the molecules of interest here are given in Table 1 using the dominant species  $^{24}\text{Mg}$  and  $^{52}\text{Cr}$ . The spectroscopic constants of MgH and CrH have been taken from Balfour & Cartwright (1976), Lemoine et al (1988) and Bauschlicher et al. (2001). The computed spectroscopic constants for MgD and CrD are given in Table 2. We adopted oscillator strengths  $f_e = 0.059$  (Kuznezova et al. 1980) and 0.001 (Pavlenko 1999) for the MgH ( $A^2\Pi - X^2\Sigma^+$ ) and the CrH ( $A^6\Sigma^+ - X^6\Sigma^+$ ) band system, respectively.

We calculated the Franck-Condon factors  $q_{v''v'}$  for the MgH and CrH band profile computations. The RADEN program of Kuzmenko et al. (1984), and the FRANKQ program (Tsymbol, 1984) were used, respectively. In the FRANKQ program a Morse potential is used to represent the true potential, and is solved for rotation-vibration motion. We compared the computed  $q_{v''v'}$  values with the results of Nicholls (1981) and found only small differences. It is worth noting that the adopted system of  $q_{v''v'}$  determines the relative strength of the  $(v'',v')$  bands, but not their location in the spectrum. In this work we are interested in the computation of the main bands of molecules of interest, therefore the choice of any  $q_{v''v'}$  does not critically affect our main results.

### 2.3 Partition function and equilibrium constants

The internal partition function is given by a sum over all states:

$$Q_{int} = \sum_{e,v,N} (2N+1)(2S+1) \exp\left(\frac{-(E_{e,v,N} - E_0)}{kT}\right), \quad (4)$$

where,  $S$  is electronic spin,  $E_{e,v,N}$  is the energy of the state with electronic excitation  $e$ , vibrational excitation  $v$  and with rotational angular momentum  $N$ ,  $E_0$  is the zero point energy. When rotational angular momentum is coupled to the electronic spin a splitting of

**Table 2.** Spectroscopic constants of MgH, MgD, CrH and CrD in  $\text{cm}^{-1}$ .

Molecule, data source		$T_e$	$\omega_e$	$\omega_e x_e$	$B_e$	$\alpha_e$	$D_e$	$\beta_e$
MgH,								
Balfour & Cartwright (1976)	$A \ ^2\Pi$	19216.8	1599.50	32.536	6.1913	0.1931	$3.60 \times 10^{-4}$	$6.1 \times 10^{-6}$
Lemoine et al. (1988)	$X \ ^2\Sigma^+$	0.00	1495.25	31.637	5.6443	0.1845	$3.53 \times 10^{-4}$	$2.57 \times 10^{-6}$
MgD,								
This work	$A \ ^2\Pi$	19216.8	1153.43	16.918	3.2195	0.07241	$9.73 \times 10^{-5}$	$1.2 \times 10^{-6}$
Lemoine et al. (1988)	$X \ ^2\Sigma^+$	0.0	1078.14	16.147	2.9668	0.068294	$9.61 \times 10^{-5}$	$2.79 \times 10^{-7}$
CrH,								
Bauschlicher et al. (2001)	$A \ ^6\Sigma^+$	11552.7	1524.80	22.280	5.3427	0.141349	$3.01 \times 10^{-4}$	$9.18 \times 10^{-5}$
	$X \ ^6\Sigma^+$	0.0	1656.05	30.491	6.2222	0.180978	$3.52 \times 10^{-4}$	$5.21 \times 10^{-6}$
CrD,								
this work	$A \ ^6\Sigma^+$	11552.7	1088.32	11.35	2.7227	0.051395	$7.82 \times 10^{-5}$	$1.70 \times 10^{-5}$
	$X \ ^6\Sigma^+$	0.0	1182.00	15.533	3.1709	0.065804	$9.14 \times 10^{-5}$	$9.66 \times 10^{-7}$
Ram & Bernath (1995)	$A \ ^6\Sigma^+$	11559.7	1066.42	-	2.7627	0.049687	$7.03 \times 10^{-5}$	-
	$X \ ^6\Sigma^+$	11559.7	1183.19	15.60	3.1754	0.065908	$9.09 \times 10^{-5}$	-

the spin degeneracy occurs. For the case of the CrH  $X \ ^6\Sigma^+$  electronic states,  $S=5/2$ , this results in 6 separate states with angular momentum  $J = N \pm 5/2, N \pm 3/2, N \pm 1/2$ . Similarly for MgH,  $X \ ^2\Sigma^+$ ,  $J = N \pm 1/2$ . The internal partition function is then expressed as:


$$Q_{int} = \sum_{e,v,J,N} (2J+1) \exp\left(\frac{-(E_{e,v,J,N} - E_0)}{kT}\right) \quad (5)$$

Energy levels are computed using the rotation, vibration constants listed in Table 2. It is worth noting a good agreement of our data with the constants obtained by Ram & Bernath (1995) from the high resolution spectroscopy of the  $A \ ^6\Sigma^+ - X \ ^6\Sigma^+$  system of CrD molecule. The partition functions of MgH, MgD, CrD are computed via direct summation of these levels, see equation 4. For CrH, we have obtained accurate energy levels from the linelist of Burrows et al. (2002). This list of energy levels covers the  $X \ ^6\Sigma^+$  and  $A \ ^6\Sigma^+$  electronic states, extends from  $J = 0.5$  to 39.5 and includes the vibrational states  $v = 0,1,2,3$  in the ground electronic state and  $v = 0,1,2$  for the excited electronic state. The CrH partition function is calculated by the direct summation of the Burrows et al. (2002) energy levels. In addition, to cover the states omitted from the Burrows et al. (2002) linelist, we supplement energy levels computed with the spectroscopic constants. The partition functions are listed in Table 3.

Figures 1 and 2 show the partition functions and the respective fits of Sauval & Tatum (1984). For CrH there is excellent agreement with the partition function of Sauval & Tatum (1984), but for MgH our partition function is slightly larger than that of Sauval & Tatum (1984) at low temperature and smaller at temperatures above 3200 K. Sauval & Tatum

**Table 3.** Internal partition functions,  $Q$ , as a function of temperature (powers of ten are given in parenthesis).

$T / \text{K}$	$Q(\text{MgH})$	$Q(\text{MgD})$	$Q(\text{CrH})$	$Q(\text{CrD})$
50	1.322(1)	2.440(1)	3.608(1)	6.855(1)
100	2.575(1)	4.815(1)	7.011(1)	1.351(2)
150	3.832(1)	7.194(1)	1.042(2)	2.018(2)
200	5.091(1)	9.582(1)	1.384(2)	2.687(2)
250	6.353(1)	1.199(2)	1.726(2)	3.359(2)
300	7.623(1)	1.445(2)	2.069(2)	4.041(2)
350	8.905(1)	1.698(2)	2.415(2)	4.738(2)
400	1.021(2)	1.961(2)	2.764(2)	5.455(2)
450	1.154(2)	2.236(2)	3.118(2)	6.199(2)
500	1.290(2)	2.524(2)	3.479(2)	6.973(2)
550	1.431(2)	2.825(2)	3.849(2)	7.780(2)
600	1.577(2)	3.142(2)	4.228(2)	8.623(2)
650	1.727(2)	3.474(2)	4.618(2)	9.503(2)
700	1.883(2)	3.823(2)	5.020(2)	1.042(3)
750	2.045(2)	4.188(2)	5.435(2)	1.138(3)
800	2.212(2)	4.569(2)	5.864(2)	1.238(3)
850	2.386(2)	4.969(2)	6.306(2)	1.343(3)
900	2.566(2)	5.385(2)	6.763(2)	1.451(3)
950	2.753(2)	5.820(2)	7.235(2)	1.564(3)
1000	2.947(2)	6.272(2)	7.722(2)	1.682(3)
1100	3.355(2)	7.233(2)	8.744(2)	1.930(3)
1200	3.791(2)	8.267(2)	9.831(2)	2.197(3)
1300	4.256(2)	9.378(2)	1.098(3)	2.482(3)
1400	4.750(2)	1.057(3)	1.221(3)	2.787(3)
1500	5.274(2)	1.183(3)	1.350(3)	3.111(3)
1600	5.827(2)	1.318(3)	1.487(3)	3.455(3)
1700	6.410(2)	1.460(3)	1.630(3)	3.818(3)
1800	7.023(2)	1.610(3)	1.782(3)	4.203(3)
1900	7.664(2)	1.767(3)	1.941(3)	4.608(3)
2000	8.333(2)	1.932(3)	2.108(3)	5.035(3)
2100	9.029(2)	2.104(3)	2.283(3)	5.483(3)
2200	9.751(2)	2.283(3)	2.467(3)	5.953(3)
2300	1.050(3)	2.469(3)	2.659(3)	6.446(3)
2400	1.127(3)	2.661(3)	2.859(3)	6.962(3)
2500	1.206(3)	2.859(3)	3.069(3)	7.501(3)
2600	1.287(3)	3.063(3)	3.288(3)	8.063(3)
2700	1.370(3)	3.272(3)	3.517(3)	8.650(3)
2800	1.455(3)	3.486(3)	3.755(3)	9.261(3)
2900	1.542(3)	3.705(3)	4.003(3)	9.896(3)
3000	1.629(3)	3.928(3)	4.261(3)	1.056(4)
3100	1.719(3)	4.154(3)	4.530(3)	1.124(4)
3200	1.809(3)	4.385(3)	4.810(3)	1.195(4)
3300	1.901(3)	4.618(3)	5.101(3)	1.269(4)
3400	1.993(3)	4.854(3)	5.403(3)	1.345(4)
3500	2.086(3)	5.093(3)	5.716(3)	1.424(4)
3600	2.180(3)	5.334(3)	6.042(3)	1.506(4)
3700	2.275(3)	5.577(3)	6.379(3)	1.590(4)
3800	2.370(3)	5.821(3)	6.728(3)	1.676(4)
3900	2.465(3)	6.067(3)	7.089(3)	1.765(4)
4000	2.561(3)	6.314(3)	7.463(3)	1.857(4)
4100	2.657(3)	6.562(3)	7.849(3)	1.951(4)
4200	2.753(3)	6.811(3)	8.248(3)	2.048(4)
4300	2.849(3)	7.060(3)	8.659(3)	2.148(4)
4400	2.945(3)	7.310(3)	9.084(3)	2.250(4)
4500	3.041(3)	7.559(3)	9.521(3)	2.354(4)
4600	3.137(3)	7.809(3)	9.971(3)	2.461(4)
4700	3.233(3)	8.059(3)	1.043(4)	2.571(4)
4800	3.329(3)	8.308(3)	1.091(4)	2.683(4)
4900	3.424(3)	8.557(3)	1.140(4)	2.797(4)
5000	3.519(3)	8.805(3)	1.190(4)	2.913(4)




**Figure 1.** Partition functions of MgH and MgD computed in this work, and the fit of Sauval & Tatum (1984).

(1984) make use of a high temperature approximation to the rotational partition function. This uses the rigid rotor approximation and is determined by rewriting the partition function sum as an integral from zero to infinity. In contrast our partition function sum is truncated at the dissociation energy  $D_0$ , which for MgH is only 1.285 eV. Thus at high temperature Sauval & Tatum (1984) predict a higher partition function than would be expected by direct summation of bound states.

We have calculated equilibrium constants for the reactions:  $\text{MgH} + \text{D} \rightleftharpoons \text{MgD} + \text{H}$  and  $\text{CrH} + \text{D} \rightleftharpoons \text{CrD} + \text{H}$ . The dimensionless equilibrium constants are given by:

$$\begin{aligned}
 K &= \frac{n(\text{CrH})n(\text{D})}{n(\text{CrD})n(\text{H})} \\
 &= \frac{Q(\text{CrH})Q(\text{D})}{Q(\text{CrD})Q(\text{H})} \exp\left(-\frac{\Delta E}{kT}\right)
 \end{aligned} \tag{6}$$

where  $n$  are the number densities, and  $\Delta E$  is the energy difference between zero point energies of CrH and CrD;  $Q$  is the total partition functions which can be factorised into



**Figure 2.** Partition functions of CrH and CrD computed in this work, and the fit of of Sauval and Tatum (1984).

internal and translational components,  $Q = Q_{int}Q_{tran}$ , where  $Q_{tran} = (2\pi M_i kT/h^2)^{3/2}$ .  $g_j$  is the degeneracy of energy level  $j$ . By convention nuclear spin degeneracy is omitted but electronic spin degeneracy is included. Thus for a proton  $Q_{int} = 1$  and for an electron  $Q_{int} = 2$ . The equilibrium constants are listed in Table 4. The equilibrium constants indicate that at low temperatures the formation of the deuterides are strongly favoured over the hydrides, but at high temperatures the hydrides become weakly favoured.

It is worth noting that in cool dwarfs hydrogen and deuterium are primarily found in their molecular forms,  $H_2$ , HD and  $D_2$ . In a full chemical equilibrium calculation the partition functions and dissociation energies of  $H_2$ , HD and  $D_2$  will have an affect upon the abundances of the molecules of interest. So a more appropriate reaction may be  $MgH + HD \rightleftharpoons MgD + H_2$ . However, to calculate equilibrium constant for this reaction would require the partition functions of  $Q(HD)$  and  $Q(H_2)$  and their dissociation energies. Expressing the equilibrium constants relative to atomic D and H avoids any uncertainties arising from the

partition functions and dissociation energies of H<sub>2</sub>, HD and D<sub>2</sub>. A calculation of the partition functions and dissociation energies of H<sub>2</sub>, HD and D<sub>2</sub> is beyond the scope of this project.

In calculating the equilibrium constants we assumed that the electronic partition functions of H and D are identical. This is a good approximation as the main differences in the energy levels are proportional to the change in reduced mass and are thus small.

## 2.4 Ionisation-dissociation equilibrium

The equations of ionisation-dissociation equilibrium were solved for media consisting of atoms, ions and molecules. We took into account  $\sim 100$  components (Pavlenko 1998, 2000). The constants for equations of chemical balance were taken from Tsuji (1973) and Gurvitz et al. (1989). To find the densities of species we apply the technique applied by Kurucz (1975) in the ATLAS5 – ATLAS12 programs. In the framework of the local thermodynamic equilibrium (LTE) approach, the densities of a molecule consisting of atoms labelled 1, 2, ... $n$  in the  $k+$  ionisation stage can be described by the Gouldberg-Waage equation:

$$\frac{n_{1,2,\dots,l}^{p+}}{\prod_i^l n_i} = \frac{Q_{1,2,\dots,n} Q_e^p}{\prod_i^n Q_i} \exp\left(-\frac{E_{1,2,\dots,l}}{kT}\right), \quad (7)$$

where  $n_i$  is the number density of the  $i^{\text{th}}$  component,  $Q_i$  are the total partition functions for species  $i$ , and  $E_{1,2,\dots,l}$  is the dissociation energy ( $D_0$ ) of molecule which consists of  $l$  atoms and has  $p^+$  charge.

In the ATLAS program a computationally more convenient format is used:

$$\frac{\prod_i^l N_i}{N_{1,2,\dots,l}} = \exp[-E_{1,2,\dots,l}/kT + g(T)], \quad (8)$$

here the function  $g(T)$  is given by:

$$g(T) = b - c(T + d(T - e(T + fT))) + \frac{3}{2}(l - p - 1) \ln T, \quad (9)$$

values of  $b, c, d, e$  (Table 5) are determined by the fitting to the data (see Table 4) using the least square minimisation procedure for the temperature range 400 - 5000 K.

As was expected, at LTE the equilibrium constants of our hydrides and deuterides are very similar. Indeed, the equilibrium constants are  $\sim Q_{int,i}/M_i$ ,  $M_i$  is the mass of the  $i^{\text{th}}$  species. Partition functions of the deuterated molecules are larger by approximately a factor

**Table 4.** Equilibrium constants for deuterium exchange reactions (powers of ten are given in parenthesis).

$T/K$	$K(\text{MgH}+\text{D}\rightleftharpoons\text{MgD}+\text{H})$	$K(\text{CrH}+\text{D}\rightleftharpoons\text{CrD}+\text{H})$
50	3.998(3)	1.759(3)
100	7.504(2)	4.973(2)
150	1.994(1)	1.515(1)
200	3.250(1)	2.644(1)
250	4.350(1)	3.690(1)
300	5.270(1)	4.600(1)
350	6.028(1)	5.372(1)
400	6.647(1)	6.020(1)
450	7.152(1)	6.561(1)
500	7.567(1)	7.012(1)
550	7.908(1)	7.390(1)
600	8.191(1)	7.707(1)
650	8.427(1)	7.975(1)
700	8.625(1)	8.201(1)
750	8.793(1)	8.395(1)
800	8.935(1)	8.560(1)
850	9.057(1)	8.703(1)
900	9.162(1)	8.826(1)
950	9.254(1)	8.933(1)
1000	9.333(1)	9.027(1)
1100	9.464(1)	9.182(1)
1200	9.567(1)	9.304(1)
1300	9.648(1)	9.400(1)
1400	9.714(1)	9.479(1)
1500	9.768(1)	9.542(1)
1600	9.812(1)	9.595(1)
1700	9.849(1)	9.640(1)
1800	9.880(1)	9.677(1)
1900	9.906(1)	9.709(1)
2000	9.928(1)	9.737(1)
2100	9.947(1)	9.761(1)
2200	9.963(1)	9.783(1)
2300	9.977(1)	9.803(1)
2400	9.988(1)	9.822(1)
2500	9.998(1)	9.840(1)
2600	1.001	9.857(1)
2700	1.001	9.874(1)
2800	1.002	9.891(1)
2900	1.003	9.909(1)
3000	1.003	9.927(1)
3100	1.003	9.945(1)
3200	1.004	9.965(1)
3300	1.004	9.985(1)
3400	1.004	1.001
3500	1.004	1.003
3600	1.004	1.005
3700	1.005	1.008
3800	1.005	1.011
3900	1.005	1.013
4000	1.005	1.016
4100	1.005	1.019
4200	1.005	1.022
4300	1.005	1.026
4400	1.005	1.029
4500	1.005	1.032
4600	1.005	1.036
4700	1.005	1.039
4800	1.005	1.043
4900	1.005	1.047
5000	1.005	1.051

**Table 5.** Chemical equilibrium constants for MgH, MgD, CrH and CrD (powers of ten in parenthesis).

Molecule	$D_0$ /eV	$b$	$c$	$d$	$e$	$f$
CrH	2.863	3.690(1)	6.678(-3)	2.541(-6)	5.115(-10)	3.929(-14)
MgH	1.285	3.605(1)	6.834(-3)	2.551(-6)	5.049(-10)	3.834(-14)
CrD	2.888	3.731(1)	6.919(-3)	2.607(-6)	5.192(-10)	3.967(-14)
MgD	1.311	3.648(1)	7.111(-3)	2.647(-6)	5.214(-10)	3.942(-14)

<sup>a</sup> From Shayesteh et al. (in prep.)


2, which corresponds to the larger molecular weight of deuterium. Fitted chemical equilibrium constants computed for CrH, CrD, MgD and MgH are shown in Table 5.

## 2.5 Synthetic spectra

Synthetic spectra were computed using the program WITA6 (Pavlenko 2000) assuming LTE, hydrostatic equilibrium and a one-dimensional model atmosphere without sources and sinks of energy. Hereafter we use the term “synthetic spectra” instead of “spectral energy distributions” to simplify the text. Theoretical synthetic spectra were computed for the model atmospheres of dwarfs with effective temperatures  $T_{\text{eff}} = 1200 - 1800$  K from the COND grid of Allard et al. (2001) using solar metallicity (Anders & Grevesse 1989). Unless otherwise mentioned all models are for  $\log g = 5.0$ .

## 2.6 Atomic lines opacity

To compute the band profiles of the electronic bands in the spectra of late spectral class dwarfs one must account for absorption of the resonance lines of Na I (0.5891, 0.5897  $\mu\text{m}$ ) and K I (0.7667, 0.7701  $\mu\text{m}$ ), which are extremely strong. In fact, optical and near infrared spectra of L- and T-dwarfs are governed by them (Pavlenko 1997, 1998). Temperatures in the atmospheres of ultracool dwarfs are lower than 2000 K, therefore alkali atoms exist there mainly in the form of neutrals. Furthermore, due to the low opacity of the atmospheres of ultracool dwarfs their photospheres occur in high pressure layers. Strong resonance lines of the most abundant alkali metals, i.e. sodium and potassium, are formed in very dense and cool plasmas. Due to the high efficiency of pressure broadening, their formally computed equivalent widths may reach a few thousand Angstroms (Pavlenko 2000). Burrows & Volobuev (2003) show that we cannot use the conventional theory of collisional broadening.



**Figure 3.** Molecular bands of the  $A^2\Pi - X^2\Sigma^+$  system of MgH and MgD in the theoretical spectrum computed for a 1800/5.0/0 COND model atmosphere. Spectra are shown for  $D/H = 1$ . The lower line on the plot shows the pure spectrum of MgH computed in the framework of JOLA approximation. For comparison the observed spectra of two L dwarfs 2MASS1632+19 (Martín et al. 1999) and SDSS 0236+0048 (Leggett et al. 2001) are shown as solid lines.

A more sophisticated theory involving quasi-stationary broadening should be used for them (see also Allard et al. 2003, and Zhu et al. 2006).

In this work we use the potentials of quasi-stationary chemical interactions of K and Na with the most numerous species, atomic He and molecular  $H_2$  computed by GAMESS (Granovsky et al. 2004). Our procedure is described in more detail in Pavlenko et al. (2007). In calculations of K I profiles we used a combined profile: the cores of these lines were computed in the framework of the collisional approach and their wings ( $\delta\lambda > 0.004 \mu\text{m}$ ) were treated by quasi-stationary theory.

### 3 RESULTS

#### 3.1 MgH & MgD

Computed JOLA profiles of the electronic band system  $A\ ^2\Pi - X\ ^2\Sigma^+$  of MgH and MgD are shown in Fig. 3. It is worth noting that a simulation based on a sound quantum mechanical model, including spin-orbit coupling in the  $A\ ^2\Pi$  state, agreed reasonably well with the JOLA calculation (see Hill 2007). First, the synthetic spectra were computed for the case of a solar abundances ( $N(\text{H})/N(\text{total})=0.9$ ). Then all the hydrogen was replaced by deuterium ( $N(\text{D})/N(\text{Total})=0.9$ ). As stated above, the aim of this paper is a comparative analyses of the possibilities of detection of the deuterated molecules MgD and CrD which are formed on the background of the electronic bands of the more abundant conventional hydrides MgH and CrH. The detection of MgD bands is very challenging for several reasons:


- The bands of MgD lie at relatively short wavelengths. High precision observations of this spectral region are difficult in objects with low  $T_{\text{eff}}$ .
- The differences in the positions of the band heads of MgH and MgD are rather small. The MgD bands will be formed on a background of far stronger MgH bands.
- MgH and MgD bands in the spectra of cool dwarfs are close to a strong Na I resonance line and the lines of other molecules/atoms.

#### 3.2 CrH & CrD

The relative behaviour of CrH and CrD bands is shown in Fig. 4. In general, the bands of the  $A\ ^6\Sigma^+ - X\ ^6\Sigma^+$  system occupy a larger wavelength range. Due to the larger reduced mass, distances between heads of CrD bands of different  $\Delta v$  are smaller than those for the CrH. As a result, bands of negative and positive  $\Delta v$  are shifted bluewards and redwards with respect to the corresponding CrH bands.

The transitions with  $\Delta v = 1$  form a band head at  $0.76843\ \mu\text{m}$  which coincides with the core of the strong K I resonance doublet. Thus in L dwarfs with strong CrH absorption the contribution of CrH has to be accounted in the modelling of  $0.77\ \mu\text{m}$  K I feature.

For the  $\Delta v = 0$  case the bands of CrH and CrD are co-incident. For the  $\Delta v$  positive case the CrD band heads are displaced to longer wavelengths of the CrH band head, thus the strongest CrD lines will be blended with the lines from the tail of the corresponding CrH band. Bands with  $\Delta v = 2$  are blended with the K I resonance line. Bands with  $\Delta v = 3$  are located in the spectral region of the so-called “satellite rainbows” of the K I line (Burrows



**Figure 4.** Molecular bands of the  $A^6\Sigma^+ - X^6\Sigma^+$  system of CrH and CrD and a theoretical spectrum computed for a 1800/5.0/0 COND model atmosphere. Spectra are shown for  $D/H = 1$ . The lower line on the plot shows the pure spectrum of MgH computed in the framework of JOLA approximation. For comparison the observed spectra of two L dwarfs 2MASS1632+19 (Martín et al. 1999) and SDSS 0236+0048 (Leggett et al. 2001) are shown as solid lines.

& Volobuev 2003). For the  $\Delta v$  negative case the CrD band heads are displaced to shorterer wavelengths of the CrH band head, these bands offer the best possibility of detection.

#### 4 DISCUSSION

The detection of deuterated molecules in the spectra of ultracool dwarfs provide a challenge for both theoreticians and observers. Indeed, in the atmospheres of planetary mass objects ( $M < 13 M_J$ ) we cannot expect ratio  $D/H > 2 \times 10^{-5}$ . This means the lines of deuterated molecules should be about 5000 times weaker than those of the hydrides. The ideal case would be a spectral region where the molecular bands are not blended. So a crucial requirement is a large difference in the wavelengths of the band heads of the hydrides and deuterides.

CrH appears to be more useful than MgH in the search for deuterated species. The band

heads of CrD are displaced significantly from the band heads of CrH. Bands of CrH are observed in the spectra of the latest L dwarfs (Kirkpatrick et al. 1999). The CrD bands are located in the “near infrared” spectra, where fluxes are much higher than in the “optical” spectral regions.

In this paper we show that the most useful bands for the realisation of the deuterium test are  $\Delta v = +1$  and  $\Delta v = -1$  ( $\lambda = 0.795$  and  $0.968 \mu\text{m}$ , respectively). The  $\Delta v = -1$  band looks especially promising. It is located in the near infrared region with an absence of strong background absorption features. However, portions of this CrD band will be swamped by the 0-0 FeH Wing-Ford band at  $0.99\mu\text{m}$  and possibly by water bands. Still, the case for CrD looks better than for HDO lines which are formed on a background of strong H<sub>2</sub>O lines (Pavlenko 2002). High quality line lists are required to test these possibilities fully.

It is worth noting, that a potential problem lies in the possibility that Cr atoms are absorbed onto dust particles. The depletion of Cr will reduce the strength of both CrD and CrH bands. Fortunately as CrH bands are located in the same spectral region, we can “scale” the CrD depletion processes by adjusting the strength of CrH bands.

For more precise studies, more accurate and detailed linelists of CrH and CrD are required. The calculation of such linelists would require new improved computations supported by new laboratory measurements. One problem that concerns us is that even once we have a good agreement with the model and experimental data for the MgH or CrH molecule, there are still perturbations about which we know very little from experiments done so far. Nonetheless, MgH is a non-starter for the deuterium test and as we note above, the model adopted in our paper is more likely to be reliable for CrH and CrD. It is worth noting that the use of the pure rotational-vibrational bands located in the mid and far infrared spectral region may offer an alternative. Indeed, the displacement between CrH and CrD rotation-vibration bands is even larger, than for the case of electronic bands.

Nevertheless, we cannot be certain that we have identified the best candidate systems for the deuterium test. Future investigations of deuterated molecules in different spectral regions are important to determine which offers the best possibility for the realisation of this test. The ideal solution would be to detect lines of deuterated molecule(s) in different spectral regions. This presents a serious observational challenge which can only be met in combination with careful laboratory measurement and the computation of high quality molecular spectra.

## 5 ACKNOWLEDGEMENTS

We thank anonymous Referee for the helpful remarks. We thank Prof. Peter Bernath for a constructive discussion and for providing us with a copy of his paper prior to publication. The UK Particle Physics and Astronomy research council is thanked for visitor and PDRA support. The Royal Society is thanked for a visitor grant. YP is supported by a Leverhulme Trust grant. This work was partially supported by the Cosmomicrophysics program of National Academy of Sciences and National Space Agency of Ukraine. Our research has made use of the SIMBAD database operated at CDS, Strasbourg, France.

## REFERENCES

- Allard, F., Hauschildt, P. H., Alexander, D. R., Tamanai, A., Schweitzer, A. 2001, *ApJ*, 556, 357.
- Allard, M.F., Allard, F, Hauschildt, P.H., Kielkopf, J.F., and Machin, L. 2003, *A&A.*, 411, L473.
- Anders, E., Grevesse, N., 1989, *GeGoAA*, 53, 197.
- D’Antona, F., Mazzitelli, I. 1998. In *Brown Dwarfs and Extrasolar Planets*, Ed. by R. Rebolo, E. Marti n, and M. R. Zapatero Osorio, *Astron. Soc. Pac. Conf. Ser.* 134, 442.
- Balfour, W. J., Cartwright, H. M., 1976, *A&AS*, 26, 389
- Bauschlicher, C.W.Jr., Ram, R.S., Bernath, P.F., et al. 2001. *J. Chem. Phys.*, 115, 1312.
- Basri, G. 2000, *ARA&A*, 38, 485
- Bejar, V. J. S., Zapatero Osorio, M. R., and Rebolo, R. 1999. *ApJ*, 521, 671.
- Bell, R. A., Gustafsson, B., 1981, *BAAS*, 13, 829.
- Bell, R. A., Edvardsson, B., Gustafsson, B., 1985, *MNRAS*, 212, 497.
- Berdyugina, S. V., Savanov, I. S, 1992, *SvA*, 36, 425.
- Bonnell, J. T., Bell, R. A., 1993, *MNRAS*, 264, 334.
- Burrows, A., Ram, S.R., Bernath, P. et al. 2002, *ApJ.*, 577, 986.
- Burrows, A., Volobuyev, M. 2003, *ApJ*, 583, 985.
- Chabrier, G., Baraffe, I., Allard, F., and Hauschildt, P., 2000, *ApJ*, 542, L122.
- Craybeal J.D. 1988. *Molecular Specroscopy*, McGraw-Hill Book Company.
- Engvold, O., Wohl, H, Brault, J.W., 1980, *A&A Suppl. Ser.*, 42, 209.
- Gaydon, A.G., Pearse, R.V.B. 1937. *Nature*, 140, 110.
- Granovsky, A.A. 2004, <http://quantum-2.chem.msu.ru/gran/gamess/index.html>

- Gurvitz, L. V., Weitz, I. V., Medvedev, V. A. 1989, Thermodynamic properties of individual substances. Moscow. Science
- Halfen, D.T., Zirijs, L.M. 2004. 611, L65.
- Hill, C. 2007 <http://www.physchem.ox.ac.uk/~hill/research/MgH/index.html>
- Jones, H.R.A., Tsuji, T. 1997, ApJ, 480, 39.
- Kirkpatrick, J. D., Allard, F., Bida, T., Zuckerman, B., Becklin, E. E., Chabrier, G.; Baraffe, I. 1999, ApJ, 519, 834,
- Kleman, B., Uhler, U., 1959, Can. J. Phys. 37, 537.
- Kuz'menko, N., Kuznetzova, L.A., Kuz'yakov, Y.Y, 1984, Franc-Condon factors for two-atomic molecules, Moscow University Publ., 1.
- Kuznezova, L.A., Kuzmenko. N.E., Kuz'yakov, Y.Y., PLastinin, Y.A. 1989, Two atomic molecules transition probabilities, Moscow, Nauka, 315 p.
- Leggett, S. K., Allard, F., Geballe, T.R., et al., 2001, ApJ, 548, 908.
- Lemoine, B., Demuynck, C., Destombes, J.L., Davies, P. B., 1988, J. Chem. Phys, 89, 673
- Lipus, K., Bachem, E, Urban, W. 1991, Mol. Phys., 73,1041.
- Magazzù, A., Martín, E. L., Rebolo, R., 1993, ApJ, 404, L17
- Martín , E. L., Delfosse, X., Basri, G., Goldman, B., Forveille, T., Zapatero Osorio, M.R. 1999, AJ, 118, 2466.
- Naersisian, S. E., Shavrina, A. V., Yaremchuk, A. A., 1989, Astrofizika, 30, 147.
- Nicholls, R.W. 1981, ApJSS, 47, 299.
- Pavlenko Ya.V. 1997, Odessa Astron. Publ., 10, 76.
- Pavlenko, Ya. V. 1997a, Astron. Astrophys. Sci., 253, 43.
- Pavlenko Ya.V. 1998, Atronom. Report, 42, 787.
- Pavlenko Ya.V. 1998a, Atronom. Report, 42, 787.
- Pavlenko, Y. V. 1999, Astron. Rep., 43, 748.
- Pavlenko, Ya.V. 2000, Astron. Rept., 44, 219
- Pavlenko, Ya.V. 2002, Astron. Rept., 46, 567
- Pavlenko, Ya.V., Zhukovska, S., Volobuyev, M. 2007, Astron. Rept., 51, 282..
- Ram, R.S., Bernath, P.F., 1995, J. Molec. Specrosc., 172, 91.
- Ram, R.S., Jarman, C.N., Bernath, P.F., 1993, J. Molec. Specrosc., 161, 445.
- Rebolo, R., Martin, E. L., Basri, G., Marcy, G. W., Zapatero-Osorio, M. R. 1996, ApJ, 469, 53.
- Saumon, D., Hubbard, W.B., Burrows, A. et al. 1996, ApJ, 460, 993

- Schadee, A. 1968, *ApJ*, 151, 239.
- Shayesteh, A., Henderson, R. D. E., Le Roy, R. J., Bernath, P. F. (in preparation).
- Shin, S., Bruch, D.J., Morse, M.D. 2005, *ApJ*, 619, 407.
- Sinha, K., Shukla, D. S., Tripathi, M. B. 1979, *BASI*, 7, 38.
- Sinha, K., Joshi, G. C., 1982, *BASI*, 10, 329.
- Skory, S., Weck, P. F., Stancil, P. C., Kirby, K. 2003, *ApJS*, 148, 599.  
2003, *ApJ*, 584, 459
- Tsuji, T. 1973, *A&A*, 23, 411
- Tsuji, T., Ohnaka, K., Aoki, W., Nakajima, T. 1996, *A&A*, 308, L29.
- Tsymbal, V.V. 1977, Tables of Franck-Condon factors with account the rotational-vibrational interactions for astrophysically important molecules. I. Titanium Oxide, Dep. No. 246-77.
- Zu-Geng Wang, Hui-Rong Xia 1996, *Molecular and Laser Spectroscopy*, Springer Verlag.
- Weck, P. F., Schweitzer, A., Stancil, P. C., et al. 2003, *ApJ*, 584, 459.
- Weck, P. F., Stancil, P. C., Kirby, K, 2003a, *ApJ*, 582, 1263.
- Weck, P. F., Schweitzer, A., Stancil, P. C., et al. 2003b, *ApJ*, 582, 1059.
- Wyller, A.A., 1961, *ApJ*, 134, 805
- Ziurys, L. M., Barclay, W. L. Jr., Anderson, M. A. 1993, *ApJ*, 402L, 21.
- Zhu, C., Babb, J.F., and Dalgarno, A. 2006, *Phys. Rev. A*, 73, 012506.

