



Healthcare Systems and Covid-19: Lessons to be Learnt from Efficient Countries

Journal:	<i>International Journal of Health Planning and Management</i>
Manuscript ID	HPM-20-00549
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	20-Jul-2020
Complete List of Authors:	Ordu, Muhammed; Osmaniye Korkut Ata University, Department of Industrial Engineering Kirli Akin, Hediye; Osmaniye Korkut Ata University, Department of Industrial Engineering Demir, Eren; University of Hertfordshire Business School, Hertfordshire Business School
Keywords:	Pandemic Outbreak, Covid19, Region, Data Envelopment Analysis, Decision Support System

SCHOLARONE™
Manuscripts

1
2
3 **Muhammed Ordu** is an Assistant Professor in the Department of Industrial Engineering at Osmaniye
4 Korkut Ata University in Turkey. His research areas are healthcare modelling, forecasting, simulation
5 and optimization. He is also interested in engineering economy and financial topics. His recent paper
6 was appeared in Journal of the Operational Research Society about a novel healthcare resource
7 allocation decision support tool: A forecasting-simulation-optimization approach.
8
9

10
11 **Hediye Kirli Akin** is an Assistant Professor in the Department of Industrial Engineering at Osmaniye
12 Korkut Ata University in Turkey. She focuses on modelling and optimization, experimental design and
13 scheduling problems in Healthcare Systems.
14
15

16
17 **Eren Demir** is a Professor at the University of Hertfordshire in the UK. He has widely in the area of
18 healthcare analytics and decision support system. He is involved in the use of modelling techniques in
19 healthcare management and the development of tools that can aid healthcare decision-making.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Title:** Healthcare Systems and Covid-19: Lessons to be Learnt from Efficient Countries
5

6 **Short Running Title:** Healthcare Systems and Covid-19
7

8
9 **Author List:** Muhammed Ordu¹, Hediye Kirli Akin¹, Eren Demir²
10

11
12 **Corresponding Author:** Muhammed Ordu, Department of Industrial Engineering, Faculty of
13 Engineering, Osmaniye Korkut Ata University, 80010, Osmaniye, Turkey
14

15
16
17 **Institutional Affiliations:** ¹ Department of Industrial Engineering, Faculty of Engineering,
18 Osmaniye Korkut Ata University, 80010, Osmaniye, Turkey; ² Hertfordshire Business School,
19 University of Hertfordshire, Hatfield, United Kingdom
20
21
22
23
24
25
26

27 **Abstract**

28
29
30 **Background:** The novel coronavirus is rapidly spreading over the world and puts the health
31 systems of countries under intense pressure. High hospitalization levels due to the pandemic
32 outbreak have caused the intensive care units to work above capacity.
33
34
35

36
37 **Purpose:** A Data envelopment analysis (DEA) based modelling approach was developed to
38 evaluate the effectiveness of regions (i.e. city, country or clinical commissioning groups)
39 against the pandemic outbreak. The objective is to enable related authorities better manage the
40 struggle against the outbreak and put in place the emergency action plans immediately.
41
42
43
44
45
46

47 **Methodology/Approach:** DEA method was used to measure the efficiency scores of countries.
48 Super efficiency DEA method was also applied to countries based on the level of efficiencies
49 they have achieved. Sixteen countries were selected that have been facing with Covid19
50 pandemic outbreak for at least five consecutive weeks after their 100th confirmed case.
51
52
53
54
55

56
57 **Results:** A total of 80 DEA models were developed, i.e. 16 DEA models for each week. The
58 percentage of efficient countries decreased dramatically over time, from 43.75% in the first
59
60

1
2
3 week to 25% in the fifth week. Unlike most European countries, China and South Korea
4
5 increased their effectiveness after first week of implementing all the necessary measures.
6
7

8 **Conclusion:** This study sheds light into better understanding the effectiveness of policies
9
10 adopted by countries and their management strategy in dealing with Covid-19 pandemic. Our
11
12 model will enable political leaders to identify inadequate policies as quickly as possible and
13
14 learn from their peers for more effective decisions.
15
16

17
18 **Key Words:** Pandemic Outbreak, Covid19, Region, Data Envelopment Analysis, Decision
19
20 Support System
21
22

23 24 25 26 **1. Introduction**

27
28 Coronaviruses are a large family of viruses that can cause a variety of conditions, from common
29
30 cold to more severe diseases, such as acute respiratory syndromes.
31
32

33
34 The first case was reported to the World Health Organization on December 31, 2019, cases of
35
36 unknown pneumonia in Wuhan, China.¹ Since then, the city of Wuhan has taken unprecedented
37
38 countermeasures against the outbreak, including closure of schools and business.
39
40

41
42 Despite these measures, the spread of the disease was not contained, and the beast is spreading
43
44 very fast, so far effecting 213 countries worldwide. As of April 24 2020, there are 2,631,839
45
46 confirmed COVID-19 cases and 182,100 deaths.²
47
48

49
50 The world has experienced and struggled with numerous outbreaks and pandemics in the past,
51
52 such as cholera, black plague, typhoid and influenza, but nothing like COVID-19, disrupting
53
54 every aspect of our lives. A sudden influx of patients to hospitals within a very short period of
55
56 time, meant that even the most advanced healthcare systems around the world were unable to
57
58 cope with the demand, and resources were stretched to its limit (e.g. beds, ventilators, personal
59
60

1
2
3 protective equipment's, intensive care beds, doctors and nurses), forcing some countries to
4
5 construct new COVID-19 hospitals and departments within days and weeks.
6
7

8 COVID-19 has prompted the scientific community to research on a wide range of issues, the
9
10 most pressing been the search for therapeutics and vaccines; modelling the spread of the disease
11
12 and expected mortality; capacity planning of hospitals; the impact on supply chain logistics and
13
14 the economy, and association of meteorological factors and disease spread.³
15
16

17
18 However, no studies have been conducted to determine how efficient and effective countries
19
20 are in terms of using scarce hospital resources to treat COVID-19 patients, in the form of
21
22 benchmarking, thus the opportunity of healthcare senior decision makers and political leaders
23
24 to learn from best practices. They will be able to understand where their country and hospitals
25
26 are heading in such a global disaster in terms of demand-capacity. Political leaders can
27
28 determine the cities of their country which need to be considered more in terms of emergency
29
30 action plan, specify how to optimize their limited resources with minimal loss.
31
32

33
34 Advanced statistical modelling is typically carried out for benchmarking
35
36 organisations/institutions, e.g. multilevel modelling. However, the main reasons to use DEA
37
38 method are as follows: 1) "DEA method enables to compare variables with "the best value"
39
40 whereas statistical methods analyse the variables based on "average value" using central
41
42 tendency approach"⁴, 2) There are multiple inputs and outputs, where DEA is the only
43
44 methodology that is able to deal with such datasets. In this study, we developed a data
45
46 envelopment analysis (DEA) based modelling to determine the efficiency levels of countries
47
48 (out of the selected 16 nations) with respect to healthcare systems preparedness (in the form of
49
50 availability of resources, i.e. number of physicians and hospital beds), demography (i.e.
51
52 population, elderly people and age) and COVID-19 confirmed cases.
53
54
55

56
57 We collected weekly number of COVID-19 cases for each country after 100th confirmed case,
58
59 as significant increase is observed after 100th case, where intensive care units have reached
60

1
2
3 intolerable levels. Note that the date that any country reaches the 100th confirmed case is
4 different. A total of 16 countries (See Table 1) is selected that had the outbreak for at least five
5 consecutive weeks after their 100th confirmed case (as of April 11, 2020).
6
7

8
9
10 DEA is an extremely popular technique applied to a wide range of sectors, including health,
11 education, agriculture, finance and manufacturing services.⁵ DEA gives a more meaningful
12 index of comparative performance, establishing both the opportunities of improvement for
13 healthcare services and reliable rankings for countries. DEA replaces multiple efficiency ratios
14 by a single weighted sum of outputs over the weighted sum of inputs. The strength and ease of
15 use of this method has attracted many researchers to develop models that rank hospital
16 departments; compare healthcare services, and establish the efficiencies of clinics (i.e. surgery,
17 gynaecology) in hospitals between countries⁶⁻¹².
18
19

20
21
22 Using the DEA-based modelling approach, this study will enable healthcare organizations and
23 governments to develop an effective strategy around resource planning and use (now and in the
24 future) by comparing most effective nations in the fight against COVID-19, and the opportunity
25 to learn from those that are highly efficient. Furthermore, assessing weekly changes on
26 efficiency levels of countries and better understanding whether the measures taken and
27 implemented plans are sufficient or not, will enables us to assist organizations and governments
28 to reallocate effectively and efficiently related resources (i.e. staff, beds, budgets), bring the
29 regions struggling with the COVID-19 outbreak under control, and allow us to debate the
30 efficiencies of action plans against COVID-19.
31
32

33
34
35 The DEA and the super efficiency methodology are explained in Section 2; Section 3 presents
36 the DEA-based modelling framework, where results are presented in Section 4, and finally
37 Section 5 concludes the article.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. Methods

DEA is a method which is applied to determine the relative efficiencies of decision-making units (DMUs) (e.g. hospitals, universities, countries) and has been widely used for a few decades.¹³ DEA method with the CCR (Charnes-Cooper-Rodes) model was developed by Charnes et al.¹⁴ and the DEA model with the BCC (Banker- Charnes-Cooper) model by Banker et al.¹⁵. The DEA model is a fractional programming which maximizes a ratio dividing the virtual outputs by virtual inputs. The weights are calculated by means of mathematical programming technique.¹³ The fractional programming was converted to linear programming by Charnes et al.¹⁶. The fractional programming of our DEA model for DMU 1 (i.e. United States) is adapted from Cooper et al.¹⁷ by considering 16 DMUs, 7 inputs and 3 outputs shown in Table 2. The formulation of a typical DEA model is in Appendix. The objective function (1) maximizes the ratio dividing output by input. Constraint (Eq. 2) ensures each ratio for every DMU not to exceed 1. Constraints (3) and (4) are positive variables.¹⁷

The linear programming (adapted from Cooper et al.¹⁷) obtained from the fractional programming above is for DMU 1 in Appendix. The objective function (5) maximizes the weighted sum of outputs. Constraints (6) ensures the weighted sum of outputs is less than or equals to the weighted sum of inputs for every DMU. Constraints (7) and (8) are positive variables.¹⁷

DEA determines efficiencies in the range of [0,1] and maximum efficiency score does not exceed 1. This therefore causes an uncertainty on the relative efficiency scores of each efficient DMU. In other words, the DEA model does not allow to rank efficient DMUs between each other. Andersen and Petersen¹⁸ modified the DEA method by excluding the evaluated DMU in the comparison with all other DMUs. They provided the efficient DMU to have efficiency score greater than 1.

1
2
3 In our study, there exists a total of 16 constraints which compares the evaluated DMU with all
4 DMUs (see Eq. 2 and 6). Several DMUs (e.g. China) are determined as efficient DMU as can
5 be seen in the next section. To rank these DMUs with each other, we used super efficiency DEA
6 method and remove the related constraint from our DEA model. For example, the Constraint
7 (9) in Appendix was removed from the DEA model developed for China in the first week in the
8 study period.
9
10
11
12
13
14
15
16
17
18
19

20 **2.1. Inputs and Outputs**

21
22
23 Figure 1 illustrates the hierarchy of the developed DEA model. The inputs are categorized into
24 three groups: demography, cases and resources. *Number of populations* is the sum of people
25 living in the country. *Percentage of people who is 70 or over* is a ratio that a specific age group
26 including elderly and risky population in terms of Covid19 over the number of populations.
27 *Median age* is the age that divides the population sorted from the youngest person to the eldest
28 one into two equal parts. *Total confirmed cases of Covid19* is the cumulated number of cases
29 from the time 100th case occurred. *Weekly total confirmed cases of Covid19* is the number of
30 cases occurring in a specific time horizon. *Number of physicians* is the number of consultants
31 per 1000 people. *Hospital beds* is the number of available beds per 100,000 people. The outputs
32 are total confirmed deaths, periodical total confirmed deaths and non-mortality rate of Covid19.
33 *Total confirmed deaths* are the cumulated number of deaths due to Covid19 from the time 100th
34 case occurred. *Weekly total confirmed deaths* are the number of deaths due to Covid19
35 occurring in a specific time horizon. *Non-mortality rate of Covid19* is a ratio of people who is
36 infected with Covid19 and not died over the total number of confirmed cases.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2.2. Data

The data for a total of 222 regions (i.e. countries and continents) are recorded and the data we used in this study is from 31st December 2019 to 11th April 2020.

The data were provided in .csv format from (Roser et al. ¹⁹, World Population Review²⁰) and imported into Microsoft SQL Server version 12.0. for database programming and analysis purposes. The data period depends on the data of 100th confirmed case within each country. After initial checks, we decided to include 16 countries in our study according to the criterion explained in the next section. Therefore, the starting dates of the first week for each country differ due to the dates of 100th occurring case, i.e. 10th of March 2020 for USA and 29th of January for Japan. Table 1 shows the starting dates of the first week and the number of days since the 100th confirmed case.

Table 2 gives the descriptive statistics of the input and output variables used in this study. The percentage of elderly people as well as median age is very high in Japan, whereas in Iran the opposite. There is a huge difference regarding total confirmed cases between the countries having minimum and maximum values, i.e. the total confirmed cases in the United States is around six times the average value. The highest number of physicians per 1000 people in Norway and Japan uses around 13 beds per 100,000 people. The lowest values for these two inputs are observed in Iran. The highest number of deaths is observed to be in the United States, whereas UK with the highest mortality rate.

3. A DEA-Based Modelling Framework

The DEA-based modelling framework includes the 5 steps to determine weekly efficiencies of regions struggling with the pandemic outbreak.

- 1
2
3 1. Determine a study period for a region (i.e. city, country, clinical commissioning groups,
4 provincial directorate of health and so on). The countries are ranked from the highest
5 number of cases to the lowest. Countries are then selected based on the following
6 criterion: to experience with Coronavirus outbreak for at least five weeks after their
7 100th confirmed case. Table 1 illustrates the countries, total number of cases and dates
8 of 100th confirmed case at the end of week 5th.¹⁹
9
- 10
11
12 2. Extract weekly data for the inputs and outputs in the study. The inputs are categorized
13 into three groups: 1) demography (i.e. number of populations, percentage of people who
14 are 70 or over and median age), 2) cases (i.e. total and weekly confirmed cases of
15 COVID-19) and 3) resources (i.e. number of physicians per 1000 people and hospital
16 beds per 100,000). The outputs are total confirmed deaths, weekly total confirmed
17 deaths and non-mortality rate of COVID-19.
18
- 19
20
21 3. We develop the DEA model by taking into account all the inputs and outputs mentioned
22 in the previous step. A total of 26 constraints related to DMUs (16 constraints) and
23 integer variables (10 constraints) along with the objective function maximizing the
24 output of the model are considered in each model. If any DMU has an efficient score
25 (i.e. 1), the DEA model is modified using the super efficiency method, so that we can
26 rank countries from the most efficient to the least.
27
- 28
29
30 4. The same process is repeated for the next period after the efficiency scores of all DMUs
31 are calculated. We applied this process for five weeks.
32
- 33
34
35 5. Rank and compare the determined regions over the selected period.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 **4. Results and Discussion**

55
56 A total of 80 DEA model was developed, i.e. 16 DEA models for each week. Table 3 and Figure
57
58 2 illustrate the efficiency scores of the DMUs for each week over the study period.
59
60

1
2
3 Most countries have lost the ability to cope with COVID-19 over time (from week 1 to week
4
5 5). While some countries (i.e. USA, UK, Germany and Belgium) in the first week were around
6
7 the efficient boundary, they experienced a dramatic decrease in the following week. The
8
9 efficiency scores gradually decreased in subsequent weeks. Countries (i.e. Spain, Italy,
10
11 Switzerland and Netherlands) having low efficiency scores since the first week failed in the
12
13 five-week period and their efficiency scores decreased further. Some of those countries with
14
15 greater than one efficiency score (i.e. France, China, South Korea, Sweden and Norway) in the
16
17 first week maintained their effectiveness and the rest experienced fluctuations. For example,
18
19 China has become an efficient country consecutively during the five-week period. As of May
20
21 2020, South Korea was still combatting with the disease and maintained efficiency for the entire
22
23 study period, whereas Iran and Singapore were super-efficient countries.

24
25
26
27
28 Table 4 summaries the DEA-based modelling approach. This study provides a periodical-based
29
30 DEA modelling framework for healthcare organizations and governments to more proactively
31
32 manage the Covid19 outbreak process. Thus, in order to overcome this process with less
33
34 material and moral losses, these institutions and organizations will be provided to update
35
36 themselves and renew their perspective. For example, the followed paths and the action plans
37
38 which efficient countries have implemented in combating the pandemic outbreak will shed light
39
40 on inefficient countries.

41
42
43
44
45 It is noteworthy that the number of tests generally performed in countries that can be considered
46
47 successful in the fight against Covid19 is high. However, this may not be sufficient alone and
48
49 the filiation teams investigating the source of Covid19 can be required to be established. The
50
51 filiation teams to be established by the Ministries of Health determine the patients with Covid19
52
53 and reach out to those who are in contact with them. In this way, virus carriers have been
54
55 identified before proceeding to the advanced stages of the disease, but the treatment process is
56
57 started immediately. These people are also prevented from getting around in the community.
58
59
60

1
2
3 In regions where the Covid19 outbreak is partially or fully controlled, if positive cases are
4 observed, the relevant regions (i.e. town, district and province) can be quarantined in a short
5 time. Thus, mobility is prevented.
6
7
8
9

10 There may be reports of new signs about Covid19 in the printed and visual media. As the virus
11 mutates, it may show different characteristics. In this process, countries, which are active in
12 treatment and tracking, constantly update the algorithms in Covid19 Treatment protocols. Thus,
13 they can be successful in reducing mortality rates and minimizing the number of intensive care
14 and intubated patients. Until today, from the beginning of Covid19 outbreak, it has been
15 revealed that herd immunity policy did not work, and this policy could destroy health systems.
16 As a result, the number and percentage of efficient DMUs in Table 4 proved that countries need
17 different policies rather than the current to control the pandemic.
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **5. Conclusion**

33
34 With this DEA-based modelling approach, some results have been reached with this study,
35 where the activities of countries in the fight against Covid19 are taken into consideration. For
36 example, when the inputs of the study are examined, it is seen that Iran is disadvantaged in
37 terms of health system preparedness, and despite its disadvantage, it has the highest efficiency.
38 It can be said that the positive effects of the younger population and outbreak management
39 increased the effectiveness.
40
41
42
43
44
45
46
47
48

49 Although Japan's population average age is high, it has been observed that it is managing well,
50 compared to other countries in terms of strength of its health system and policies they
51 implemented during the pandemic.
52
53
54
55
56
57
58
59
60

1
2
3 While the average number of healthcare workers in countries analysed within the scope of the
4 study is 11.63 per 1,000 people, this number in America is 9,73; the average bed capacity is
5 3.13 per 100,000 people, and 2.57 in USA.
6
7
8
9

10 In all low-productivity countries, such as Spain, Italy, Netherlands and Switzerland, the average
11 age of its population is above the average of the remaining countries in the study. However,
12 they are the more developed countries having advanced and modern health systems. These low-
13 efficient countries need to urgently change their outbreak management policies to turn their age
14 disadvantage into an advantage.
15
16
17
18
19
20

21
22 As can be seen from our inputs and outputs, the mortality rate and spread of the virus do not
23 depend on a single factor. With the combination of a few negative factors, such as elderly
24 population average, health conditions, and poor management policies, the efficiency decreases
25 gradually.
26
27
28
29
30
31

32 This study can guide the organizations on outbreak management policies and the analysis of
33 the current situation of the countries. In addition, the results obtained from the study and
34 especially the epidemic management policies of the countries with high effectiveness have been
35 analysed and the basis has been established for the preparation of a guide for combating the
36 Covid19 outbreak.
37
38
39
40
41
42

43
44 We considered only 16 countries in this study due to the reasons explained in the previous
45 sections. However, the study can be further extended by including more countries, longitudinal
46 data considering more than 5 weeks, additional inputs (e.g. total number of tests, weekly
47 number of tests, number of intensive care units) and output (e.g. total number of recovered
48 patients and weekly number of recovered patients). We only used seven inputs and three
49 outputs, and additional variables could have been considered, such as the number of tests for
50 Covid19 per week. During the pandemic outbreak, researchers might have difficulties in
51
52
53
54
55
56
57
58
59
60

1
2
3 accessing data, however, organizations such as WHO and/or governments can make relevant
4 data available for research purposes.
5
6
7
8
9

10 **Acknowledgement**

11
12
13 Not Applicable
14
15
16
17
18

19 **Conflict of Interest**

20
21
22 The authors declare that there is no conflict of interest
23
24
25
26
27

28 **Ethics Statements**

29
30 The paper does not require any human/animal subjects to acquire ethics approval
31
32
33
34
35

36 **References**

- 37
38
39 1. Coronavirus disease (COVID-19) outbreak. World Health Organization Web Site.
40 [http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-](http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov)
41 [coronavirus-2019-ncov](http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov). Accessed April 25, 2020.
42
43
44
45
46 2. WHO Coronavirus Disease (COVID-19) Dashboard. World Health Organization Web Site.
47 <https://covid19.who.int/> Accessed April 24, 2020.
48
49
50
51 3. Ma Y, Zhao Y, Liu J, et al. Effects of temperature variation and humidity on the death of
52 COVID-19 in Wuhan, China. *Science of the Total Environment*. 2020; 724. doi:
53 10.1016/j.scitotenv.2020.138226
54
55
56
57
58
59
60

- 1
2
3 4. Sahin G, Gokdemir L, Ozturk, D. Global Crisis and Its Effect on Turkish Banking Sector:
4
5 A Study with Data Envelopment Analysis. *Procedia Economics and Finance*. 2016;
6
7 38(2016), 38-48. doi: 10.1016/S2212-5671(16)30174-5
8
9
- 10
11 5. Emrouznejad A, Yang GL. A survey and analysis of the first 40 years of scholarly literature
12
13 in DEA: 1978–2016. *Socio-Economic Planning Sciences*. 2018; 61:4-8. doi:
14
15 10.1016/j.seps.2017.01.008
16
17
- 18 6. Kao L, Lu C, Chiu C. Efficiency measurement using independent component analysis and
19
20 data envelopment analysis. *European Journal of Operational Research*. 2011;210(2011):
21
22 310-317. doi: 10.1016/j.ejor.2010.09.016
23
24
- 25 7. Gok MS, Sezen B. Analyzing the ambiguous relationship between efficiency, quality and
26
27 patient satisfaction in healthcare services: The case of public hospitals in Turkey. *Health*
28
29 *Policy*. 2013;111(2013): 290-300. doi: 10.1016/j.healthpol.2013.05.010
30
31
- 32 8. Girginer N, Kose T, Uckun N. Efficiency Analysis of Surgical Services by Combined Use
33
34 of Data Envelopment Analysis and Gray Relational Analysis. *Journal of Medical Systems*.
35
36 2015;2015: 39-56. doi: 10.1007/s10916-015-0238-y
37
38
- 39 9. Foo CY, Lim KK, Sivasampu S, Dahian KB, Goh PP. Improving the effectiveness of service
40
41 delivery in the public healthcare sector: the case of ophthalmology services in Malaysia.
42
43 *BMC Health Services Research*. 2015;15(349). doi: 10.1186/s12913-015-1011-0
44
45
- 46 10. Levary RR, Ip C. Determining the relative efficiency of gynecological departments using
47
48 DEA. *Financial Modeling Applications and Data Envelopment Applications*. 2015:261-
49
50 273. doi: 10.1108/S0276-8976(2009)0000013016
51
52
- 53 11. Cheng Z, Cai M, Tao H, et al. Efficiency and productivity measurement of rural
54
55 township hospitals in China: a bootstrapping data envelopment analysis. *BMJ Open*. 2016;
56
57 6, e011911. doi: 10.1136/bmjopen-2016-011911
58
59
60

12. Khushalani Y, Ozcan YA. Are hospitals producing quality care efficiently? An analysis using Dynamic Network Data Envelopment Analysis (DEA). *Socio-Economic Planning Sciences*. 2017;60(2017): 5-23. doi: 10.1016/j.seps.2017.01.009
13. Po RW, Gu YY, Yang MS. A new clustering approach using data envelopment analysis. *European Journal of Operational Research*. 2009;199(2009): 276-284. doi: 10.1016/j.ejor.2008.10.022
14. Charnes A, Cooper WW, Rhodes E. Measuring the efficiency of decision-making units. *European Journal of Operational Research*. 1978;2:429–444.
15. Banker RD, Charnes A, Cooper WW. Some models for estimating technical and scale inefficiencies in data envelopment analysis. *Management Science*. 1984;3:1078–1092.
16. Charnes A, Cooper WW, Rhodes E. Evaluating program and managerial efficiency: An application of data envelopment analysis to program follow through. *Management Science*. 1981;27:668–697.
17. Cooper WW, Seiford LM, Tone K. *Data Envelopment Analysis*. 2nd Edition. USA: Springer; Springer, 2007. doi: 10.1007/978-0-387-45283-8
18. Andersen P, Petersen NC. A procedure for ranking efficient units in data envelopment analysis. *Management Science*. 1993;39:1261–1264.
19. Roser M, Ritchie H, Ortiz-Ospina E, Hasell J. Coronavirus Pandemic (COVID-19). Our World in Data Web Site. <https://ourworldindata.org/coronavirus>. Accessed April 11, 2020.
20. Countries by Median Age 2018. World Population Review Web Site. <https://worldpopulationreview.com/countries/median-age/>. Accessed April 11, 2020.

Appendix

$$Max \theta = \frac{u_1 y_{11} + u_2 y_{21} + u_3 y_{31}}{v_1 x_{11} + v_2 x_{21} + v_3 x_{31} + v_4 x_{41} + v_5 x_{51} + v_6 x_{61} + v_7 x_{71}} \quad (1)$$

$$\frac{u_1 y_{1j} + u_2 y_{2j} + u_3 y_{3j}}{v_1 x_{1j} + v_2 x_{2j} + v_3 x_{3j} + v_4 x_{4j} + v_5 x_{5j} + v_6 x_{6j} + v_7 x_{7j}} \leq 1 \quad (2)$$

$$v_1, v_2, v_3, v_4, v_5, v_6, v_7 \geq 0 \quad (3)$$

$$u_1, u_2, u_3 \geq 0 \quad (4)$$

$$Max \theta = \mu_1 y_{11} + \mu_2 y_{21} + \mu_3 y_{31} \quad (5)$$

$$\mu_1 y_{1j} + \mu_2 y_{2j} + \mu_3 y_{3j} \leq v_1 x_{1j} + v_2 x_{2j} + v_3 x_{3j} + v_4 x_{4j} + v_5 x_{5j} + v_6 x_{6j} + v_7 x_{7j} \quad (6)$$

$$v_1, v_2, v_3, v_4, v_5, v_6, v_7 \geq 0 \quad (7)$$

$$\mu_1, \mu_2, \mu_3 \geq 0 \quad (8)$$

where u_s is the weight of output s , v_i is the weight of input i , y_{sj} is the value of output s for DMU j , and x_{ij} is the value of input i for DMU $j, j=1, 2, \dots, 16$

$$\mu_1 y_{16} + \mu_2 y_{26} + \mu_3 y_{36} \leq v_1 x_{16} + v_2 x_{26} + v_3 x_{36} + v_4 x_{46} + v_5 x_{56} + v_6 x_{66} + v_7 x_{76} \quad (9)$$

where u_s is the weight of output s , v_i is the weight of input i , y_{s6} is the value of output s for DMU China, x_{i6} is the value of input i for DMU China.

Table 1: A summary of confirmed COVID-19 cases and dates of the first 100th cases in the selected countries.

Decision Making (DMUs)	Units	Countries	Total Number of Cases	Number of days since the 100th confirmed case (days)	Dates the 100th case was confirmed case
DMU1		United States	501560	39	10.03.2020
DMU2		Spain	157022	39	10.03.2020
DMU3		Italy	147577	47	02.03.2020
DMU4		Germany	117658	41	08.03.2020
DMU5		France	90676	41	08.03.2020
DMU6		China	83004	83	26.01.2020
DMU7		United Kingdom	70272	36	13.03.2020
DMU8		Iran	68192	44	05.03.2020
DMU9		Belgium	26667	35	14.03.2020
DMU10		Switzerland	24228	35	14.03.2020
DMU11		Netherlands	23097	35	14.03.2020
DMU12		South Korea	10450	50	28.02.2020
DMU13		Sweden	9685	35	14.03.2020
DMU14		Norway	6244	35	14.03.2020
DMU15		Japan	5347	49	29.02.2020
DMU16		Singapore	1909	41	08.03.2020

DMU = Decision Making Unit

Table 2: Descriptive statistics of the input and output variables

Variables	Mean	Standard Deviation	Minimum		Maximum		
			Country	Value	Country	Value	
Inputs	Number of Population (\tilde{x}_1)	15047192	341510669.1	Singapore	5850342	China	1439323776
	Median Age (\tilde{x}_2)	40.84	4.22	Iran	30.3	Japan	47.3
	Percentage of people who is 70 or over (\tilde{x}_3)	11.64	3.86	Iran	3.18	Japan	18.49
	Physicians (per 1,000 people) (\tilde{x}_4)	3.13	0.91	Iran	1.49	Norway	4.39
	Hospital beds (per 100,000) (\tilde{x}_5)	4.90	3.34	Iran	1.5	Japan	13.05
	Total confirmed cases of Covid19 (\tilde{x}_6)	65804	87202.12	Singapore	1189	United States	368196
	Weekly total confirmed cases of Covid19 (\tilde{x}_7)	28250	47977	Singapore	386	United States	203576
Outputs	Total confirmed deaths (\tilde{y}_1)	4036	4337.16	Singapore	6	United States	10989
	Weekly total confirmed deaths (\tilde{y}_2)	2225	2538.43	Singapore	3	United States	7819
	Non-mortality rate of Covid19 (\tilde{y}_3)	93.82	4.25	United Kingdom	87.74	Singapore	99.50

Table 3: Efficiency and super efficiency scores of the DMUs for each week of the 5 weeks


Decision Making Units	Countries	1st week	2nd week	3rd week	4th week	5th week
DMU1	United States	0.96	0.86	0.84	0.84	0.82
DMU2	Spain	0.87	0.76	0.74	0.73	0.71
DMU3	Italy	0.76	0.69	0.68	0.67	0.65
DMU4	Germany	0.92	0.72	0.70	0.71	0.69
DMU5	France	1.01	0.79	0.78	0.76	0.72
DMU6	China	1.21	0.84	0.86	0.87	1.07
DMU7	United Kingdom	0.90	0.81	0.79	0.75	0.72
DMU8	Iran	2.11	2.09	2.05	2.01	2.05
DMU9	Belgium	0.93	0.82	0.80	0.77	0.74
DMU10	Switzerland	0.81	0.81	0.81	0.79	0.79
DMU11	Netherlands	0.81	0.78	0.76	0.73	0.72
DMU12	South Korea	1.03	0.85	0.95	0.99	0.99
DMU13	Sweden	1.09	1.05	1.03	1.00	0.97
DMU14	Norway	1.08	1.08	1.08	1.07	1.06
DMU15	Japan	0.96	0.94	0.94	1.33	0.93
DMU16	Singapore	4.52	62.62	54.77	123.67	59.39

Table 4: The summary of the DEA-based modelling


	1st week	2nd week	3rd week	4th week	5th week
Maximum efficiency score	1.00	1.00	1.00	1.00	1.00
Minimum efficiency score	0.76	0.69	0.68	0.67	0.65
Number of efficient DMUs	7	4	4	5	4
Total number of DMUs	16	16	16	16	16
% of efficient DMUs	43.75	25.00	25.00	31.25	25.00

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



The hierarchy of the developed DEA model for the Covid19 outbreak



The Graph for the first 5-week efficiency scores of the countries (Exc. Iran and Singapore) struggling with Covid19

260x153mm (96 x 96 DPI)