

Associations between the portion sizes of food groups consumed and measures of adiposity in the British National Diet and Nutrition Survey

Mary T. Kelly¹, Kirsten L. Rennie^{1,2}, Julie M. W. Wallace¹, Paula J. Robson³, Robert W. Welch¹, Mary P. Hannon-Fletcher¹ and M. Barbara E. Livingstone^{1*}

¹Northern Ireland Centre for Food and Health (NICHE), University of Ulster, Cromore Road, Coleraine BT52 1SA, UK

²Unilever Corporate Research, Colworth House, Sharnbrook, Bedfordshire MK44 1LQ, UK

³School of Public Health, University of Alberta, Edmonton, AB, Canada T6G 2G3

(Received 6 February 2008 – Revised 5 August 2008 – Accepted 5 August 2008 – First published online 10 October 2008)

The objective of the present study was to examine the associations between the portion sizes of food groups consumed with measures of adiposity using data from the National Diet and Nutrition Survey of British adults. Seven-day weighed dietary records, physical activity diaries and anthropometric measurements were used. Foods eaten were assigned to thirty different food groups and analyses were undertaken separately for men and women. The median daily portion size of each food group consumed was calculated. The potential misreporting of dietary energy intake (EI) was identified using the following equation: $EI - \text{estimated energy requirements} \times 100 = \text{percentage of under-reporting (UR) of energy needs}$. Multinomial logistic regression (adjusted for age, social class, physical activity level and UR) was used to determine the portion sizes of food groups most strongly associated with obesity status. Few positive associations between the portion sizes of food groups consumed and obesity status were found. However, UR was prevalent, with a median UR of predicted energy needs of 34 and 33 % in men and women, respectively. After the adjustment was made for UR, more associations between the food groups and obesity status became apparent in both sexes. The present study suggests that the true effect of increased portion size of foods on obesity status may be masked by high levels of UR. Alternatively, these data may indicate that an increased risk of obesity is not associated with specific foods/food groups but rather with an overall increase in the range of foods and food groups being consumed.

Obesity: Portion size: Under-reporting

The prevalence of obesity is rapidly increasing in the UK, with about two-thirds of the adult population now either overweight or obese⁽¹⁾. Overweight and obesity are associated with a host of adverse health outcomes such as diabetes, CVD and certain cancers⁽²⁾. Although the specific underlying causes of weight gain in adults remain unclear, a number of environmental factors have been identified as drivers of an excess energy intake (EI) in the face of reduced energy needs.

One particular alteration in dietary patterns which has been associated with promoting obesity is the trend towards larger food portions. Consumers may be increasingly underestimating their portion sizes and EI as they eat out of the home more often or eat larger portions within the home^(3,4). It has also been suggested that consumers cannot precisely judge the amount of food they consume over the course of a few days or weeks⁽⁵⁾. The passive over-consumption associated with large food portions could potentially lead to increased EI at the time of eating and over subsequent days, and, if not compensated for, could be a significant factor contributing towards excess weight gain and obesity. Thus, the reliance on self-monitoring to control food and EI to maintain a healthy body weight has become increasingly difficult in our obesogenic environment.

Larger food portions have been shown to increase EI in adults at single eating occasions^(6–11) and over the course of 2⁽¹²⁾ and 11 d⁽¹³⁾, under semi-controlled laboratory conditions. To date, only one study⁽¹⁴⁾ has examined the extent to which large food portions affect EI in free-living populations.

The aim of the present study was to examine the association between the portion sizes of foods consumed with indices of obesity status (BMI and waist action level (WAL)) in free-living adults aged 19–64 years in the British National Diet and Nutrition Survey (NDNS).

Methods

Survey design

The NDNS is a programme of cross-sectional surveys of different age groups designed to be representative of Great Britain. This analysis is based on the survey undertaken between July 2001 and June 2002 in adults aged 19–64 years. The survey methodology has been described in detail elsewhere⁽¹⁵⁾. Data from the NDNS adult survey were obtained from the UK Data Archive, University of Essex.

Abbreviations: EER, estimated energy requirements; EI, energy intake; NDNS, National Diet and Nutrition Survey; UR, under-reporting; WAL, waist action level.

* **Corresponding author:** Professor M. Barbara E. Livingstone, fax +44 28 70323023, email mbe.livingstone@ulster.ac.uk

Dietary measures

Subjects completed a 7-d weighed dietary record of all foods and drinks consumed both within and outside of the home. Each subject was issued a set of digital food scales and two recording diaries: the 'home record' and a smaller 'eating and drinking away from home' diary. The subjects were also issued with a pocket-sized notebook for recording any of this information in circumstances where they were reluctant, or it was inappropriate, to carry the 'eating out' diary. They were shown how to weigh and record leftovers and how to record any food that was spilt or otherwise lost. The 'eating out' diary was intended for use only when it was not possible to weigh the food items. In these cases, the subjects were asked to record as much information as they could about each food item consumed, in particular the portion size consumed and an estimate of leftovers, in addition to descriptions, brand names, price and the time and the place where food was purchased. Intakes of nutrients were calculated from the records of food consumption using the Food Standards Agency nutrient databank⁽¹⁵⁾. All foods from the NDNS dataset were assigned to thirty different food groups. Twenty-eight of these food groups were calculated by the Office for National Statistics⁽¹⁵⁾. An additional two food groups – confectionery and soft drinks – were also calculated, because of their potential association with overweight and obesity^(16,17). Only subjects who consumed the foods from a particular food group were included in the analyses of that food group. The total quantity (g/d) of each food group consumed on a daily basis and median daily portion size of each food group consumed were calculated.

Under-reporting

Using data from the physical activity diaries that were kept concurrently with the dietary record, physical activity levels were calculated to assign the appropriate estimated energy requirements (EER) equation to the individual's activity needs⁽¹⁸⁾. Physical activity levels were calculated from the time spent in sleep, light, moderate and vigorous intensity activities for each day of recording for each subject. The time spent in each type of activity was multiplied by a metabolic equivalent to give a total metabolic equivalent hours of activity per day. These were summed up to give a mean metabolic equivalent score for the week and were then divided by 24 h to give a physical activity level. The published equations of the 2002 Institute of Medicine of the National Academies were used to calculate EER on an individual basis⁽¹⁹⁾. These equations, which are derived from doubly labelled water energy expenditure data, are sex and age specific, and are based on the age, weight and height of the subject. They categorise four levels of activity: sedentary; low activity; active; very active. Potential misreporting of dietary EI was identified using the following equation:

$$EI - EER/EER \times 100 = \text{percentage of UR of energy needs.}$$

Anthropometric measurements

Anthropometric measurements recorded have been described previously⁽²⁰⁾. Height was measured in duplicate to the nearest 0.1 cm in bare feet using a portable stadiometer (Leicester

height measure; CMS Weighing Equipment, Ltd., London, UK). Body weight was measured in duplicate to the nearest 0.1 kg using digital scales. Waist circumference was measured in duplicate to the nearest 0.1 cm midway between the top of the hip bone and the lower rib. The subjects were categorised according to the WHO BMI cut-offs for obesity⁽²⁾ as follows: normal weight (BMI 18.5–24.9 kg/m²), overweight (BMI 25.0–29.9 kg/m²) and obese (BMI \geq 30.0 kg/m²). Abdominal obesity categories were defined as WAL, according to Lean *et al.*⁽²¹⁾, as follows: men – below action level = 0–93.99 cm, WAL 1 = 94–101.9 cm, WAL 2 = $>$ 102.0 cm; women – below action level = 0–79.9 cm, WAL 1 = 80.0–87.99 cm, WAL 2 = $>$ 88.0 cm.

Exclusions

At the end of the dietary recording period, the subjects completed questionnaires about illness during the recording period and whether this illness affected their eating. They were also asked questions on whether or not they were dieting to lose weight. Those who reported dieting (*n* 415) or that illness had affected their eating (*n* 210) during the recording period were excluded from the analysis. Subjects who were underweight (BMI $<$ 18.5 kg/m²; *n* 58) and without complete dietary or anthropometry data were also excluded. The analyses were subsequently carried out on 1519 subjects.

Statistical analyses

All analyses were performed using SPSS version 11.5 (Statistical Package for Social Sciences, Chicago, IL, USA). Differences between men and women in baseline characteristics were assessed using *t* tests; weight-status category, WAL and social class category were assessed by χ^2 tests. All the data were assessed for normality and owing to the high level of skewness within the food groups, the groups were split into sex-specific quartiles for analysis. Mann–Whitney *U* tests were used to assess the differences between men and women for the portion size (g) of food groups consumed. UR between weight-status categories and quartiles of portion size was assessed using ANOVA. Multinomial logistic regression analyses were used to determine the associations between the portion sizes of food groups with weight-status category and WAL. This type of analysis was chosen as the proportion of overweight and obese subjects was much greater than that of the normal weight group; therefore, binomial logistic comparisons between normal weight and overweight/obese were not possible. Models were adjusted for age (model 1), social class, physical activity level (model 2) and model 2 plus UR (percentage of UR as a continuous variable; model 3). If model 1 was NS, no further analyses were done. Analyses were done separately for men and women. Results were considered significant at *P* < 0.05.

Results

Subjects

Subject characteristics, anthropometric measurements and social class categories of the 1519 subjects (689 men and 830 women) are presented in Table 1. There was no difference in age between the men and women (*P* = 0.946). Women were significantly

Table 1. Subject characteristics, anthropometric measures and social class groupings of men and women in the National Diet and Nutrition Survey (NDNS) of British Adults 2001

(Mean values and standard deviations or percentages)

	Men* n 689		Women* n 830		P
	Mean	SD	Mean	SD	
Age (years)	42.4	11.9	42.3	11.9	0.946
Weight (kg)	84.4	13.7	69.8	14.9	0.001
Height (m)	1.8	0.7	1.6	0.7	0.001
BMI (kg/m ²)†‡	27.2	4.2	26.7	5.6	0.036
Normal (%)	31.3		44.7		0.001
Overweight (%)	44.3		34.2		0.387
Obese (%)	24.4		21.1		0.705
WAL†‡					
BAL (%)	46.8		44.3		0.110
WAL 1 (%)	24.5		28.2		0.001
WAL 2 (%)	28.7		27.5		0.175
Social class†					
Manual (%)	46.7		32.2		0.023
Non-manual (%)	53.3		67.8		0.001

WAL, waist action level; BAL, below action level.

* Data between men and women are compared using independent *t* tests.† BMI, WAL and social class categories compared using χ^2 tests.‡ BMI categories are as follows: normal weight, 18.5–24.9 kg/m²; overweight, 25.0–29.9 kg/m²; obese, ≥ 30.0 kg/m². WAL categories are as follows: BAL – men < 93.99 cm, women < 79.9 cm; WAL 1 – men = 94–101.9 cm, WAL 1 – women = 80.0–87.99 cm; WAL 2 – men > 102.0 cm, WAL 2 – women > 88.0 cm.

shorter ($P < 0.001$), lighter ($P < 0.001$), had a lower BMI ($P = 0.036$) and were more likely to be of normal weight than men ($P < 0.001$). There were no differences in the proportion of men and women who were classified as either overweight ($P = 0.387$) or obese ($P = 0.705$). Significantly more women than men were classified as having 'WAL 1' ($P < 0.001$). No differences were observed in 'below action level' ($P = 0.110$) or 'WAL 2' ($P = 0.175$) between men and women.

Under-reporting

UR was found to be 34 and 33 % of estimated energy needs in men and women, respectively. UR was found to significantly increase with the weight-status categories (Fig. 1), and was the highest among the obese category in both men and women (37.1 and 38.2 % of energy needs, respectively; $P < 0.001$) compared with the normal and overweight categories.

Portion sizes

Men had significantly higher median intakes for almost all of the food groups compared with women (Table 2). The exceptions were low-fat milks, yoghurts, vegetables, fruits, juices and nuts, fish and beverages where there were no significant differences between the sexes. In the majority of food groups, the highest level of UR was observed in the lowest quartile of portion size.

Eating occasions and energy intakes within and outside of the home

Compared with the men, the women recorded significantly more (8.6 (SD 3.4) v. 7.6 (SD 3.6); $P < 0.001$) in-home eating occasions but recorded significantly less (3.1 (SD 2.7) v. 4.9 (SD 3.2); $P < 0.001$) out-of-home eating occasions.

Table 3 displays the mean daily in-home and out-of-home eating occasions and related EI for men and women across weight-status categories. In-home or out-of-home eating occasions or EI did not vary by weight-status category in men. However, overweight women recorded significantly more in-home eating occasions compared with obese women ($P = 0.027$), while normal weight women recorded significantly more out-of-home eating occasions compared with overweight women ($P = 0.011$). EI consumed outside of the home was also significantly higher for normal weight women compared with overweight women ($P < 0.001$).

Portion sizes and BMI weight status

Few positive associations were found between the portion sizes of food groups and BMI weight-status category (Table 4). In men, only three positive associations were found (whole milk, potatoes, and fresh meat). When the adjustment was made for UR, a further three food groups (breads and rolls, low-fat spreads and vegetables) became significantly positively associated with weight-status groupings. In women, only two positive associations were found (low-fat spreads and fish). When the adjustment was made for UR, a further four food groups (breads and rolls, low-fat milks, chips and processed potatoes and meat products) became significantly positively associated with weight-status groupings.

Portion sizes and abdominal obesity categories

Similar to BMI, few positive associations were found between the portion sizes of food groups with WAL (Table 5). In the men, no positive associations were found until the adjustment was made for UR, when two food groups (breads and rolls and low-fat spreads) became positively associated with WAL. In the women, four food groups (whole milk, chips and processed potatoes, meat products and beverages) were positively associated with WAL. After the adjustment was made for UR, a further three food groups (low-fat spreads, confectionery and soft drinks) were found to be positively associated with WAL.

Discussion

Recently, interest has been focused on the contribution that larger portion sizes may be making to the over-consumption of energy and thus the increasing prevalence of obesity. Studies carried out under semi-controlled laboratory conditions have shown that increasing food portion size can positively influence EI^(6–8,12,13,22) in the short term, but data on associations between the measures of adiposity with the amounts of foods consumed in free-living populations are extremely limited⁽¹⁴⁾.

When these associations were examined in the NDNS (2001) adult survey, portions of some food groups (men: breads and rolls, low-fat spreads and vegetables; women: breads and rolls, low-fat milks, chips and processed potatoes and meat products) were not initially significantly positively associated with obesity status (BMI) but became so after the adjustment for the estimated degree of UR of EI. Similarly, five food groups (men: breads and rolls, low-fat spreads; women: low-fat spreads, confectionery and soft drinks) were




Fig. 1. Estimated under-reporting as a percentage of energy needs by weight-status category in the British National Diet and Nutrition Survey 2001. □, normal; ▨, overweight; ▩, obese. Data are presented as mean (SD), and are analysed using ANOVA. Men: normal–overweight, $P=0.014$; normal–obese, $P<0.001$. Women: normal–overweight, $P<0.001$; overweight–obese, $P=0.010$; normal–obese, $P<0.001$.

only positively and significantly associated with abdominal obesity status (WAL) when the model was adjusted for UR of EI. These data demonstrate the importance of taking into account UR when assessing the association between diet and obesity status if false associations are to be avoided.

The present study has shown that in both men and women, irrespective of their weight-status category, the majority of eating occasions reported over the 7-d recording period occurred within the home. Although the reported frequency of in-home eating occasions was higher than out-of-home

Table 2. Median portion sizes of food groups for men and women in the National Diet and Nutrition Survey (NDNS) of British Adults 2001 (Medians and interquartile ranges (IQR))

Food group	Men				Women				P†
	n*	Percentage*	Median	IQR	n*	Percentage*	Median	IQR	
Rice and pasta	516	74.9	62.8	37, 106.2	598	72	50.6	30.1, 87.7	0.001
Savouries	359	52.1	26.6	11.4, 55.6	411	49.5	18.5	8.7, 35.7	0.001
All breads and rolls	680	98.7	115.1	78.8, 163.4	813	98	78.6	53.1, 105.4	0.001
Breakfast cereals	435	63.1	36.8	19, 66.8	581	70	29.3	15.3, 50	0.001
Biscuits, cakes and pastries	569	82.6	33.4	16.6, 66.3	703	84.7	28.8	14, 50	0.001
Whole milk	254	36.9	99.5	29.1, 242.8	304	36.6	77.0	20.1, 178.8	0.010
Low-fat milks	554	80.4	181.4	81.4, 294.9	682	82.2	165.7	80.6, 259	0.145
Creams, ice creams and desserts	349	50.7	32.8	17.2, 58.9	446	53.7	27.0	16.1, 48.3	0.002
Cheeses	539	78.2	16.6	8.8, 29.3	622	74.9	14.4	8.1, 24.3	0.004
Yoghurts	236	34.3	49.8	23.5, 77	356	42.9	42.8	24.7, 79.1	0.744
Egg and egg dishes	480	69.7	24.4	14.3, 43.6	552	66.5	20.5	11.4, 33	0.001
Butter and spreads	451	65.5	5.8	2.6, 12	528	63.6	4.2	2, 8.9	0.001
Low-fat spreads	479	69.5	11.1	5.8, 18.3	538	64.8	7.1	3.1, 13	0.001
Potatoes, boiled, mashed etc	585	84.9	62.6	35, 98.2	702	84.6	57.1	32.3, 91.1	0.030
Chips and processed potatoes	584	84.8	53.6	29.3, 78.9	661	79.6	40.7	23.6, 64.2	0.001
Vegetable dishes	553	80.3	26.4	13.7, 47.9	673	81.1	23.2	12.8, 41.6	0.012
Vegetables	667	96.8	105.6	65.8, 151.6	818	98.6	98.6	59.5, 151.1	0.119
Fruits, juices and nuts	505	73.3	61.0	26.7, 127.4	660	79.5	67.9	31.8, 135.2	0.065
Fish	374	54.3	30.0	16.3, 52.9	493	59.4	27.2	13.4, 49	0.076
Fish dishes and products	245	35.6	25.7	19.3, 34.2	276	33.3	21.6	14.7, 27.5	0.001
Fresh meat	662	96.1	133.9	90.3, 186.4	763	91.9	96.2	60.9, 143.7	0.001
Meat dishes	210	30.5	17.35	10.5, 30	179	21.6	13.1	6.3, 22.5	0.002
Meat products	573	83.2	48.3	25.8, 84.1	562	67.7	33.6	17.3, 54.5	0.001
Alcoholic beverages	568	82.4	436.3	165.1, 851.5	621	74.8	122.1	42.8, 246.4	0.001
Sugars	517	75.0	17.6	7.1, 35.6	587	70.7	9.4	3.6, 20.5	0.001
Confectionery	409	59.4	14.2	7.6, 35.6	536	64.6	12.3	6.1, 22.8	0.003
Savoury snacks	369	53.6	11.6	5.5, 19.6	465	56	8.3	4.6, 14.3	0.001
Soups and sauces	659	95.6	40.3	18.4, 74.7	786	94.7	34.0	14.8, 69	0.018
Beverages	683	99.1	944.6	641.2, 1289.8	825	99.4	965.7	614.9, 1348.5	0.868
Soft drinks	413	59.9	124.4	51.5, 317.8	454	54.7	99.0	47.1, 218.7	0.008

* Number and percentage of consumers of food group.

† Data between men and women are compared using Mann–Whitney U tests.

Table 3. Daily eating occasions and energy intake (EI) in-home and out-of-home over 7 d by weight-status category*
(Mean values and standard deviations)

	Normal weight†				Overweight‡				Obese‡			
	Eating occasions		Energy (MJ)		Eating occasions		Energy (MJ)		Eating occasions		Energy (MJ)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Men‡												
In-home§	7.25	3.47	6.13	2.45	7.72	3.61	6.34	2.49	7.66	3.56	6.56	2.69
Out-of-home	5.22	3.25	3.76	2.43	4.75	3.29	3.40	2.38	4.65	3.10	3.37	2.27
Women												
In-home§	8.58	3.52	5.07	1.84	9.01	3.43	5.20	1.74	8.15	3.12	5.13	1.88
Out-of-home	3.45	2.84	2.05	1.64	2.84¶	2.54	1.60**	1.33	2.88	2.41	1.77	1.33

* Data are presented as mean daily eating occasions (frequency) and EI (MJ) per eating occasion by weight-status category. Data across weight-status categories are assessed using ANOVA.

** Significantly different from normal weight out-of-home EI.

† Normal weight (BMI 18.5–24.9 kg/m²), overweight (BMI 25.0–29.9 kg/m²), obese (BMI ≥ 30.0 kg/m²).

‡ Significantly different from women for in-home and out-of-home eating occasions.

§ Significantly different from out-of-home eating occasions.

|| Significantly different from overweight in-home eating occasions.

¶ Significantly different from normal weight out-of-home eating occasions.

Table 4. Associations between portion sizes (g) of food groups consumed with BMI status category in men and women in the National Diet and Nutrition Survey (NDNS) of British adults 2001*

Food group	BMI†			
	Men		Women	
	Model 2‡	Model 3‡	Model 2‡	Model 3‡
Rice and pasta	0.267	0.340	0.769	0.543
Savouries	0.797	0.854	0.134	0.097
All breads and rolls	0.393	0.010	0.780	0.042
Breakfast cereals	0.491	0.072	0.623	0.585
Biscuits, cakes and pastries	0.591	0.595	0.345	0.137
Whole milk	0.019	0.049	NS	NS
Low-fat milks	0.849	0.302	0.122	0.032
Creams, ice creams and desserts	0.765	0.481	0.057	0.123
Cheeses	0.332	0.193	0.907	0.715
Yoghurts	0.756	0.806	0.162	0.233
Egg and egg dishes	0.940	0.726	0.868	0.775
Butter and spreads	0.774	0.928	0.316	0.494
Low-fat spreads	0.092	0.006	0.001	0.001
Potatoes, boiled, mashed etc	0.004	0.003	0.780	0.758
Chips and processed potatoes	0.219	0.329	0.189	0.054
Vegetable dishes	0.304	0.206	0.958	0.990
Vegetables	0.124	0.021	0.526	0.296
Fruits, juices and nuts	0.144	0.419	0.073	0.178
Fish	NS	NS	0.020	0.010
Fish dishes and products	NS	NS	0.212	0.162
Fresh meat	0.049	0.006	0.691	0.473
Meat dishes	0.526	0.585	0.861	0.855
Meat products	0.343	0.308	0.092	0.009
Alcoholic beverages	0.940	0.965	0.202	0.071
Sugars	0.718	0.909	0.545	0.948
Confectionery	NS	NS	0.893	0.444
Savoury snacks	0.361	0.358	0.717	0.789
Soups, sauces	0.895	0.897	0.882	0.897
Beverages	0.515	0.729	0.057	0.452
Soft drinks	0.089	0.100	0.791	0.581

* Data are analysed using multinomial logistic regression analysis. Numbers represent *P* values. *P* values are trends across the BMI categories.

† BMI categories are as follows: normal weight, 18.5–24.9 kg/m²; overweight, BMI 25.0–29.9 kg/m²; obese, BMI ≥ 30.0 kg/m².

‡ Model 1 adjusted for age, model 2 adjusted for age, physical activity level and social class, and model 3 = model 2 + adjusted for percentage of UR. All significant associations are positive. *P* > 0.05 in model 1 is NS.

Table 5. Associations between portion sizes (g) of food groups consumed and waist action level (WAL) category in men and women in the National Diet and Nutrition Survey (NDNS) of British adults 2001*

Food group	WAL†			
	Men		Women	
	Model 2‡	Model 3‡	Model 2‡	Model 3‡
Rice and pasta	0.487	0.535	0.948	0.876
Savouries	0.981	0.988	0.113	0.093
All breads and rolls	0.473	0.017	0.161	0.064
Breakfast cereals	0.827	0.772	0.856	0.865
Biscuits, cakes and pastries	0.866	0.641	0.673	0.352
Whole milk	0.486	0.670	0.025	0.015
Low-fat milks	0.853	0.544	0.597	0.204
Creams, ice creams and desserts	0.760	0.620	0.222	0.315
Cheeses	0.616	0.269	0.632	0.531
Yoghurts	0.362	0.454	0.099	0.132
Egg and egg dishes	0.598	0.551	0.759	0.753
Butter and spreads	0.599	0.601	0.625	0.506
Low-fat spreads	0.092	0.004	0.073	0.042
Potatoes, boiled, mashed etc	0.936	0.857	0.427	0.355
Chips and processed potatoes	0.456	0.412	0.019	0.002
Vegetable dishes	0.586	0.453	0.090	0.094
Vegetables	0.410	0.108	0.474	0.552
Fruits, juices and nuts	0.236	0.664	0.074	0.058
Fish	0.624	0.601	0.289	0.220
Fish dishes and products	0.406	0.343	0.704	0.755
Fresh meat	0.709	0.435	0.959	0.884
Meat dishes	0.518	0.437	0.753	0.589
Meat products	0.412	0.322	0.001	0.001
Alcoholic beverages	0.250	0.169	0.255	0.164
Sugars	0.059	0.062	0.881	0.951
Confectionery	0.646	0.407	0.238	0.049
Savoury snacks	0.676	0.673	0.966	0.952
Soups, sauces	0.749	0.746	0.413	0.518
Beverages	0.446	0.661	0.002	0.013
Soft drinks	0.093	0.061	0.177	0.044

BAL, below action level.

* Data are analysed using multinomial logistic regression analysis. Numbers represent *P* values. *P* values are trends across the WAL categories.

† WAL categories are as follows: BAL (men < 93.99 cm, women < 79.9 cm); WAL 1 men = 94–101.9 cm, WAL 1 women 80.0–87.99 cm; WAL 2 = men > 102.0 cm, WAL 2 women > 88.0 cm.

‡ Model 1 adjusted for age, model 2 adjusted for age, physical activity level and social class, and model 3 = model 2 + adjusted for percentage of UR. All significant associations are positive.

eating occasions, it is not clear whether this is a true reflection of their usual eating patterns.

Only one previous observational study has attempted to quantify the risk of excess body fatness with food portion sizes of foods consumed by free-living adults⁽¹⁴⁾. Adiposity was positively associated with the consumption of large portions of specific foods (savory snacks, butter, full-fat spreads, meat products and dishes and chips and processed potatoes) in Irish adults. By contrast, a different range of food groups (breads and rolls, whole milk, low-fat spreads, chips and processed potatoes, fresh meat, meat products) were positively associated with a higher BMI or waist circumference in British adults. Both studies, however, found that increased portion sizes of specific food groups were associated with increased BMI and WAL. The food groups examined in both studies were very similar and some of these food groups (meat products, chips and processed potatoes) were found to be strongly associated with measures of adiposity in both studies.

Although both surveys were conducted in a representative sample of free-living adults using 7-d dietary record methodology, in the present study food intake was quantified by direct weighing, while the earlier study relied on estimating food weights using standardised portion sizes. To what extent the

difference in the methods of assessing portion weights of foods eaten explains these findings are unclear. On the other hand, it may be that there are real dietary differences between the two populations. Both studies adjusted for the potential impact of energy misreporting. McCarthy *et al.*⁽¹⁴⁾ used the ratio of EI to estimated BMR (EI:BMR), whereas a more specific calculation using physical activity data and a continuous variable based on collated doubly labelled water energy expenditure data⁽¹⁹⁾ was used in the present analyses. As previously reported, there is a high level of UR in the NDNS⁽²³⁾. In the present analyses, UR of energy needs was estimated at 34 and 33% in the men and women, respectively, and furthermore there was a higher level of UR among overweight and obese adults compared with those of a normal weight adult, which has been previously observed^(24,25). The adjustment for UR is a vital consideration when assessing possible relationships between dietary intakes and variables of interest, since this may lead to the attenuation of associations when levels of misreporting are high⁽²⁶⁾. The results of the present study emphasised the importance of adjusting for UR when examining the relationships between the portion sizes of food groups consumed with measures of adiposity. The high levels of UR of EI observed in the NDNS and the subsequent adjustment for UR in the statistical models did have an impact on

the associations between the portion sizes of food groups consumed with BMI and WAL in both the men and women. If the UR of energy needs had not been adjusted for, the observed associations may have been missed or it could have been concluded that there are no associations between the portion sizes of food groups consumed with obesity status.

A number of issues of UR are pertinent to the present discussion. First, whether misreporting is due, at least partially, by misrepresentation of usual portion sizes consumed. We found that adjustment for UR significantly affected the interpretation of associations between food portion sizes and measures of body fatness (BMI and WAL). Since dietary intake in free-living individuals in studies is self-reported, there is the opportunity for both measurement error and misreporting of food portion size, which in turn affects the estimate of overall EI.

Second, it is possible that any associations were masked by study participants who used the recording period as an opportunity to change their eating behaviour or to diet. Since under-eating while recording food intake has been observed in lean and obese adults⁽²⁷⁾, we were careful to exclude from the analysis, those subjects who reported dieting during the recording period. Nevertheless, it is likely that the dataset did include subjects who were limiting their intake during the recording period but who consciously or subconsciously failed to declare this. Body weight was only measured once during the recording period; therefore, it was not possible to examine the changes in body weight as a means of detecting under-consumption of food intake.

The third issue is whether UR is associated with the achievement of a self-presentation goal on the part of the study participant. It has been suggested that instead of modifying the diet, specific foods are under-reported or omitted in order to appear in a socially favourable light⁽²⁸⁾. Lafay *et al.*⁽²⁹⁾ suggested that the UR of food intake seems to concern food items that are generally considered 'bad for health'. This may have been the case in the NDNS, where there may have been a possible tendency to under-report or even exclude the foods that are considered to be 'unhealthy' or associated with obesity, such as savoury snacks, creams, ice creams and desserts, biscuits, cakes and soft drinks. However, it was not possible to systematically assess whether there was UR of specific foods in the NDNS as only estimates of UR of EI were feasible. There is also the possibility that foods that were perceived as healthy may have been over-reported; however, there are no available data to support this contention. Consequently, the causal associations between food portion size and obesity remain unclear, owing not only to the cross-sectional nature of food consumption surveys but also to the misreporting of food intake of unknown magnitude and direction.

In the present study, it is highly likely that UR may have masked any true associations between the food portion sizes consumed with measures of obesity. Alternatively, portion sizes of specific foods or food groups in isolation may not be major contributing factors to obesity risk, rather it may be the consumption of a larger range of foods and food groups. Gaining a better understanding of the influence of food portion size on EI and obesity may be a crucial step in identifying a modifiable dietary factor that may be influencing the current epidemic of overweight and obesity.

Acknowledgements

This research was commissioned by the Food Standards Agency (research project N09021). Ethical approval for the NDNS survey was granted by a Multi-centre Research Ethics Committee (MREC). Approval was also sought from National Health Service Local Research Ethics Committees (LREC) in the areas where the fieldwork took place. There are no conflicts of interest. The contributions of the authors were as follows: M. B. E. L. was the principal investigator and is the guarantor. M. T. K. and K. L. R. performed data analysis and wrote the manuscript. M. B. E. L., J. M. W., P. J. R., R. W. W. and M. P. H. F. contributed to data analysis and to the writing of the manuscript.

The present paper forms part of the dissemination of the Food Standards Agency-commissioned research project N09021.

References

1. House of Commons Health Committee (2001) *Tackling Obesity in England*. London: The Stationery Office.
2. World Health Organisation (1998) *Obesity: Preventing and Managing the Global Epidemic. International Obesity Task Force Report*. Geneva: WHO.
3. Nielsen SJ & Popkin BM (2003) Patterns and trends in food portion sizes, 1977–1998. *J Am Med Assoc* **289**, 450–453.
4. Smiciklas-Wright H, Mitchell DC, Mickle SJ, Goldman JD & Cook A (2003) Foods commonly eaten in the United States, 1989–1991 and 1994–1996: are portion sizes changing? *J Am Diet Assoc* **103**, 41–47.
5. Frobisher C & Maxwell SM (2003) The estimation of food portion sizes: a comparison between using descriptions of portion sizes and a photographic food atlas by children and adults. *J Hum Nutr Diet* **16**, 181–188.
6. Rolls BJ, Morris EL & Roe LS (2002) Portion size of food affects energy intake in normal-weight and overweight men and women. *Am J Clin Nutr* **76**, 1207–1213.
7. Rolls BJ, Roe LS, Kral TV, Meengs JS & Wall DE (2004) Increasing the portion size of a packaged snack increases energy intake in men and women. *Appetite* **42**, 63–69.
8. Rolls BJ, Roe LS, Meengs JS & Wall DE (2004) Increasing the portion size of a sandwich increases energy intake. *J Am Diet Assoc* **104**, 367–372.
9. Wansink B & Cheney MM (2005) Super Bowls: serving bowl size and food consumption. *J Am Med Assoc* **293**, 1727–1728.
10. Wansink B & Kim J (2005) Bad popcorn in big buckets: portion size can influence intake as much as taste. *J Nutr Educ Behav* **37**, 242–245.
11. Wansink B, Painter JE & North J (2005) Bottomless bowls: why visual cues of portion size may influence intake. *Obes Res* **13**, 93–100.
12. Rolls BJ, Roe LS & Meengs JS (2006) Larger portion sizes lead to sustained increases in energy intake over 2 days. *J Am Diet Assoc* **106**, 543–549.
13. Rolls BJ, Roe LS & Meengs JS (2007) The effect of large portion sizes on energy intake is sustained for 11 days. *Obesity (Silver Spring)* **15**, 1535–1543.
14. McCarthy SN, Robson PJ, Livingstone MB, Kiely M, Flynn A, Cran GW & Gibney MJ (2006) Associations between daily food intake and excess adiposity in Irish adults: towards the development of food-based dietary guidelines for reducing the prevalence of overweight and obesity. *Int J Obes (Lond)* **30**, 993–1002.
15. Henderson L, Gregory J & Swan G (2002) *The National Diet and Nutrition Survey: adults aged 19 to 64 years, Technical*

- Report. Social Survey Division of the Office for National Statistics **16**, 2002. London: TSO.
16. Bell AC & Swinburn BA (2004) What are the key food groups to target for preventing obesity and improving nutrition in schools? *Eur J Clin Nutr* **52**, 258–263.
 17. Ludwig DS, Peterson KE & Gortmaker SL (2001) Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet* **357**, 505–508.
 18. Rennie KL, Coward A & Jebb SA (2007) Estimating under-reporting of energy intake in dietary surveys using an individualised method. *Br J Nutr* **97**, 1169–1176.
 19. Institute of Medicine of the National Academies (2002) *Dietary Reference Intakes for Energy, Carbohydrate, Fibre, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington, DC: National Academies Press.
 20. Ruston D, Hoare J, Henderson L, Gregory J, Bates C, Prentice A, Birch M, Swan G & Farron M (2000) *The National Diet and Nutrition Survey: Adults Aged 19–64 Years; Nutritional Status (Anthropometry and Blood Analytes) Blood Pressure and Physical Activity*, vol. 4. London: TSO.
 21. Lean MEJ, Han TS & Morrison CE (1995) Waist circumference as a measure for indicating need for weight management. *BMJ* **311**, 158–161.
 22. Rolls BJ, Roe LS & Meengs JS (2004) Salad and satiety: energy density and portion size of a first-course salad affect energy intake at lunch. *J Am Diet Assoc* **104**, 1570–1576.
 23. Rennie KL, Jebb SA, Wright A & Coward WA (2005) Secular trends in under-reporting in young people. *Br J Nutr* **93**, 241–247.
 24. Heitmann BL & Lissner L (1995) Dietary under-reporting by obese individuals – is it specific or non-specific? *BMJ* **311**, 986–989.
 25. Prentice AM, Black AE, Coward WA, Davies HL, Goldberg GR, Murgatroyd PR, Ashford J, Sawyer M & Whitehead RG (1986) High levels of energy expenditure in obese women. *BMJ (Clin Res Ed)* **292**, 983–987.
 26. Livingstone MB & Black AE (2003) Markers of the validity of reported energy intake. *J Nutr* **133**, Suppl. 3, 895S–920S.
 27. Westerterp KR & Goris AHC (2002) Validity of the assessment of dietary intake: problems of misreporting. *Curr Opin Clin Nutr Metab Care* **5**, 489–493.
 28. Hebert JR, Clemow L, Pbert L, Ockene IS & Ockene JK (1995) Social desirability bias in dietary self-report may compromise the validity of dietary intake measures. *Int J Epidemiol* **24**, 389–398.
 29. Lafay L, Mennen L, Basdevant A, Charles MA, Borys JM, Eschwege E & Romon M (2000) Does energy intake underreporting involve all kinds of food or only specific food items? Results from the Fleurbaix Laventie Ville Sante (FLVS) study. *Int J Obes Relat Metab Disord* **24**, 1500–1506.