

Beryllium abundances in stars with planets: Extending the sample[★]

M. C. Gálvez-Ortiz¹, E. Delgado-Mena², J. I. González Hernández^{2,3}, G. Israelian², N. C. Santos⁴, R. Rebolo^{2,5}, and A. Ecuivillon²

¹ Centre for Astrophysics Research, Science and Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB, UK. e-mail: M.Galvez-Ortiz@herts.ac.uk

² Instituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain

³ Dpto. de Astrofísica y Ciencias de la Atmósfera, Facultad de Ciencias Físicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain

⁴ Centro de Astrofísica, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal

⁵ Consejo Superior de Investigaciones Científicas, Spain

Received November 2009; accepted

ABSTRACT

Context. Chemical abundances of light elements as beryllium in planet-host stars allow us to study the planet formation scenarios and/or investigate possible surface pollution processes.

Aims. We present here an extension of previous beryllium abundance studies. The complete sample consists of 70 stars hosting planets and 30 stars without known planetary companions. The aim of this paper is to further assess the trends found in previous studies with less number of objects. This will provide more information on the processes of depletion and mixing of light elements in the interior of late type stars, and will provide possible explanations for the abundance differences between stars that host planets and “single” stars.

Methods. Using high resolution UVES spectra, we measure beryllium abundances of 26 stars that host planets and 1 “single” star mainly using the λ 3131.065 Å Be II line, by fitting synthetic spectra to the observational data. We also compile beryllium abundance measurements of 44 stars hosting planets and 29 “single” stars from the literature, resulting in a final sample of 100 objects.

Results. We confirm that the beryllium content is roughly the same in stars hosting planets and in “single” stars at temperatures $T_{\text{eff}} \gtrsim 5700$ K. The sample is still small for $T_{\text{eff}} \lesssim 5500$ K, but it seems that the scatter in Be abundances of dwarf stars is slightly higher at these cooler temperatures.

Conclusions. We search for distinctive characteristics of planet hosts through correlations of Be abundance versus Li abundance, age, metallicity and oxygen abundance. These could provide some insight in the formation and evolution of planetary systems, but we did not find any clear correlation.

Key words. stars: abundances — stars: fundamental parameters — stars: planetary systems — stars: planetary systems: formation — stars: atmospheres

1. Introduction

Since the discovery of the first extrasolar planets, many efforts have been made to characterise planet-host stars and to find the features that can distinguish them from stars that do not have any known planetary companion (e.g. Santos et al. 2004a; Fischer & Valenti 2005). The study of chemical composition and abundances in general (e.g. Ecuivillon et al. 2004a,b, 2006; Gilli et al. 2006; Neves et al. 2009), and, in particular, of the light elements Li, Be and B (e.g. Santos et al. 2002, 2004c; Israelian et al. 2004, 2009), provided some hints on the influence of a planetary companion in the composition and evolution of stars. These light elements are destroyed by (p, α)-reactions at relatively low temperatures (about 2.5, 3.5 and 5.0×10^6 K for Li, Be and B, respectively). Thus, light elements and their abundance ratios are good tracers of stellar internal structure and allow us to extract valuable information about

the mixing mechanism and rotation behaviour in these stars (e.g. Pinsonneault et al. 1990).

Several groups (e.g. Gonzalez et al. 2001; Santos et al. 2001, 2004a) have found a correlation between stellar metallicity and the presence of giant planetary companion among solar-like stars. It has been also largely discussed whether the trend is due to “primordial” origin, i.e. the frequency of planetary companion is a function of the proto-planetary disc metal composition (see e.g. Santos et al. 2004a); or is due to a posterior “pollution”, i.e. the metallicity excess is due to matter accretion after reaching the main sequence (see e.g. Pasquini et al. 2007). Most studies suggest the “primordial” scenario although some examples may indicate that both cases occur (Israelian et al. 2001, 2003; Ashwell et al. 2005; Laws et al. 2003).

However, it is not clear whether a direct relationship between metallicity and probability of planet formation exists or if it is only applicable to the “type” of planets that have been discovered so far. Santos et al. (2004a) suggested that two distinct populations of exoplanets can be present depending on whether or not the planet is formed by a metallicity-dependent process.

[★] Based on observations obtained with UVES at VLT Kueyen 8.2 m telescope in programme 074.C-0134(A)

Recently, Haywood (2009) has also proposed that the metallicity excess in the sample of stars hosting planets may come from a dynamical effect of galactic nature (i.e. migration of stars in the galactic disc), and may not be related with the giant planets formation process.

Israeli et al. (2004) found that exoplanet hosts are significantly more Li-depleted than comparison stars but only in the range 5600–5850 K. This result has been confirmed by Chen & Zhao (2006), Takeda et al. (2007), and Gonzalez (2008). On the other hand, Gonzalez (2008) proposed that planet-host stars hotter than 5900 K have more Li than the comparison ones. There is observational evidence that Li depletion is connected with the rotational history of the star (e.g. García López 1994; Randich et al. 1997). Thus, similar stars, i.e. with similar stellar parameters and age, should have different depletion rates depending on the proto-planetary disc mass and composition and its effect on the rotation of the parent star (Bouvier 2008).

More recently, Israeli et al. (2009) has confirmed that Li is more depleted in Sun-like stars hosting planets than in similar stars without detected planets. These stars with and without planets, which fall in the effective temperature range 5700–5850 K, do not show any differences in the correlation patterns of their Li abundances with age and metallicity, pointing to a connection between the presence of planets and the low Li abundance measured in Sun-like planet-host stars.

Whatever the metallicity “excess” in planet-host stars is due to, light elements can still provide insights on the influence of planets on the parent star evolution and in the internal evolution of stars in general, they serve as a key for understanding pollution “events”, and also as tracers of the possible planet-disc interaction.

In previous works directly related with this paper (Santos et al. 2002, 2004b,c), a comparison between stars hosting planets and stars without known planets revealed that, with some exceptions, the two samples follow approximately the same behaviour in the variation of Be abundance with the temperature, similar to Li trend. This behaviour shows a Be abundance maximum near 6100 K, decreasing towards higher and lower temperatures and a Be “gap” for solar-temperature stars. Santos et al. (2004c) also remarked the possible tendency of the planet hosts to be more Be-rich, at least for $T_{\text{eff}} \gtrsim 5700$ K.

In this paper we derive beryllium abundances for 26 stars with planets and one star without planets, and collect the lithium abundances for most of them. This sample enlarges the previous samples of 44 stars with planets and 29 stars without known planetary companions reported in García López & Pérez de Taoro (1998); Santos et al. (2004c). Thus, we end up with a sample of roughly 100 stars where 70 host planets. We studied the behaviour of these abundances with the spectral type and luminosity class in both samples of stars in order to further assess the trends found before, and to study possible interactions between extrasolar planets and their parent stars and its connection to the light-element abundances through time.

2. Observations and data reduction

We obtained near-UV high-resolution spectra of the targets using the UVES spectrograph at the 8.2-m Kueyen VLT (UT2) telescope (run ID 074.C-0134(A)) on 21, 22 December 2004, in the blue arm with a wavelength coverage of $\lambda\lambda 3025\text{--}3880$ Å. These spectra have a spectral resolution $\lambda/\delta\lambda \sim 70,000$.




Fig. 1. HR diagram: surface gravity vs effective temperatures of the stars in our sample including the stars in Tables 1, 2 and 3. We overplot solar metallicity tracks from Girardi et al. (2000) for 0.7 to 1.8 M_{\odot} . The luminosity classes of the stars in the sample are assigned according with each star position in this diagram (See Sect. 3). Metallicity is represented by the size of the symbol.

All the data were reduced using IRAF¹ tools in the echelle package. Standard background correction, flat-field, and extraction procedures were used. The wavelength calibration was done using a ThAr lamp spectrum taken during the same night. Finally, we normalized the spectra by a low-order polynomial fit to the observed continuum.

3. Sample

The total sample of ~ 100 stars, including 70 planet-host stars, cover a wide temperature range from 4500 to 6400 K. Thus, the stars hosting planets have spectral types between F6 to K2 and three luminosity classes (V, IV and III). The luminosity classes were assigned following a criterion based on the position of the star in the Hertzsprung-Russell (HR) diagram, and also on their surface gravity, i.e. dwarfs (V) have typically $\log g > 4$, subgiants (IV) have $3.3 < \log g < 4$ and giants (III) have $\log g < 3.3$. We built an effective temperature-surface gravity plot with solar metallicity tracks from Girardi et al. (2000) for 0.7 to 1.8 M_{\odot} and according to each star position in this diagram (see Fig. 1), we assigned a luminosity class, presented in column 10 of Table 1 and column 7 of Tables 2 and 3. In Fig. 1 we represent the metallicity of the objects by increasing the symbol size according to three possible metallicity ranges from smaller to bigger size: $[\text{Fe}/\text{H}] < -0.5$, $-0.5 < [\text{Fe}/\text{H}] < 0.0$ and $0.0 < [\text{Fe}/\text{H}] < 0.5$, respectively. Thus, our sample of planet-host stars with new Be abundance measurements contains 3 giants, 2 subgiants and 21 dwarfs that give a total of 4 giants, 6 subgiants and 60 dwarfs in this study. The 30 stars without known planets of our sample cover the same spectral types but only two of them are a subgiant star. All the new objects are provided in Table 1 (with 26 stars with planets and one star without known planetary companions). Stars with and without known planets from literature are listed in Tables 2 and 3, respectively.

¹ IRAF is distributed by the National Optical Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation.




Fig. 2. Best synthetic spectral fits to the observed spectra in the Be II line region. From upper to lower from left to right, the planet-host dwarfs HD 33636 and HD 114729, the subgiant HD 99492 and the giant HD 47536 are displayed.

4. Stellar parameters

Stellar parameters were taken from the detailed analysis of Sousa et al. (2008) when available, otherwise from Santos et al. (2004a, 2005). These parameters are determined from ionization and excitation equilibrium of Fe I and Fe II. They used the 2002 version of the code MOOG² (Snedden 1973) and a grid of local thermodynamical equilibrium (LTE) model atmospheres (Kurucz 1993). The adopted parameters, effective temperature, T_{eff} , surface gravity, $\log g$, metallicities, $[\text{Fe}/\text{H}]$, and masses are listed in Table 1. The mean values of the uncertainties on the parameters from Santos et al. (2004a, 2005) are of the order of 44 K for T_{eff} , 0.11 dex for $\log g$, 0.08 km s^{-1} for ξ_t , 0.06 dex for metallicity and the adopted typical relative error for the masses is $0.05 M_{\odot}$. Whereas from Sousa et al. (2008) these mean uncertainties are 25 K for T_{eff} , 0.04 dex for $\log g$, 0.03 km s^{-1} for ξ_t and 0.02 dex for metallicity and $0.10 M_{\odot}$ for the masses. We refer to Sousa et al. (2008) and Santos et al. (2004a, 2005) for further details. In Tables 2 and 3 we give the main data of the sample of stars with and without planets from Santos et al. (2004b,c). We note here the uniformity of the adopted stellar parameters (see Section 5 in Sousa et al. 2008).

One should note that stellar masses are estimated from theoretical isochrones which are strongly sensitive to the adopted helium and metal content of the star. In addition, the stellar mass, although being a fundamental parameter, is relatively

more uncertain that other stellar parameters, because it depends on the effective temperature, surface gravity and metallicity of the star. However, unlike the stellar age, the stellar mass for main-sequence stars is typically better constrained from theoretical isochrones than for subgiant and/or giant stars (see e.g. Allende Prieto et al. 2004).

5. Beryllium and lithium abundances

The Be abundances were derived by fitting synthetic spectra to the data. These synthetic spectra were convolved with a Gaussian smoothing profile and a radial-tangential profile to take into account the spectral resolution and the macroturbulence, respectively. The latter was varied between 1.0 and 5.0 km s^{-1} , between K and F dwarfs (Gray 1992), respectively. A rotational profile was also added to account for the projected rotational velocity, $v \sin i$, of the sample stars, given in Table 1. These were calculated from the width of the CORALIE cross-correlation function (see appendix of Santos et al. 2002), or taken from the literature (mostly Fischer & Valenti 2005). A limb-darkening coefficient of 0.6 was adopted in all cases, and the overall metallicity was scaled to the iron abundance.

To derive Be abundances, we fitted all the observed spectral range between 3129.5 and 3132 Å, where two Be II lines are located ($\lambda 3130.420$ and $\lambda 3131.065$ Å). We used mainly the Be II $\lambda 3131.065$ Å line as the feature for deriving the best-fit Be abundance, and Be II $\lambda 3130.420$ Å was used only for checking, since it is highly blended with other elements lines.

² The source code of MOOG 2002 can be downloaded at <http://verdi.as.utexas.edu/moog.html>




Fig. 3. Be abundances as a function of effective temperature (top panel) and as a function of mass (bottom panel) for the planet-host stars in our sample plus the stars from García López & Pérez de Taoro (1998); Santos et al. (2002, 2004b,c). Filled and open symbols represent stars with and without planets. Circles, squares and triangles, are depicted for dwarfs (luminosity class V), subgiants (IV) and giants (III). In the top panel we superimpose the Be depletion isochrones (case A) of Pinsonneault et al. (1990) for solar metallicity and an age of 1.7 Gyr. From top to bottom, the lines represent a standard model (solid line) and 4 models (dashed lines) with different initial angular momentum. We also depict the Be depletion model (dotted line) including gravity waves provided in Montalbán & Schatzman (2000).

We adopted the same line list as in Santos et al. (2004c), which provides a solar photospheric Be abundance of $\log N(\text{Be}) = 1.1 \text{ dex}^3$, using the same models and tools as in this work. The main source of error is probably the placement of the continuum. An uncertainty on the continuum position of 3% yields to an error in the Be abundance of $\sim 0.05 \text{ dex}$ for solar type stars (see e.g. Randich et al. 2002). We estimate this uncertainty to be $\sim 0.06 \text{ dex}$ at $T_{\text{eff}} = 5500 \text{ K}$, $\sim 0.11 \text{ dex}$ at $T_{\text{eff}} = 5200 \text{ K}$, and $\sim 0.20 \text{ dex}$ at $T_{\text{eff}} = 4900 \text{ K}$. Another weak point is the possible 0.3 dex difference between the solar photospheric Be abundance and the meteoritic Be abundance. It has been suggested that this could be due to our inability to properly account for all continuum opacity sources in the UV (Balachandran & Bell 1998). However, other authors have argued that Kurucz atmospheric models are able to reproduce

³ Here we use the notation $\log N(\text{Be}) = \log[N(\text{Be})/N(\text{H})] + 12$




Fig. 4. Be abundances as a function of effective temperature (top panel) and as a function of mass (bottom panel) for only dwarfs with (filled circles) and without known planetary companion (open circles). Overplotted are the best linear fits found for both population together.

the near-UV absolute continuum, at least, for stars in the T_{eff} range 4000–6000 K (Allende Prieto & Lambert 2000). In fact, Balachandran & Bell (1998) argue that the Fe I bound-free opacity should be increased by a factor of 1.6, which is equivalent to an increase of 0.2 dex in the Fe abundance. Smiljanic et al. (2009) have tested this possibility for a model of $[\text{Fe}/\text{H}] -0.5 \text{ dex}$ and they find that the difference in Be abundance is small, $\Delta \log N(\text{Be}) \sim 0.02 \text{ dex}$.

For stars with effective temperatures below 5100 K, the spectral region surrounding the Be II $\lambda 3131 \text{ \AA}$ line begins to be dominated by the contribution of other species (Mn I, C, OH lines, etc.) and for the giant stars also the thulium line (Tm II $\lambda 3131.255 \text{ \AA}$) may play an important role (see Melo et al. 2005). This makes it more difficult to fit the observed spectra and thus, to measure Be abundance. Although we took into account some of these other elements when fitting cooler and giant objects, we assume that Be abundances are not so accurate in these cases. In addition, the sensitivity of the Be II to the Be abundance decreases towards lower temperatures. Thus, the total error, including mainly the uncertainties on the effective temperature and the continuum location, in the Be abundance is of the order of 0.1 dex at $T_{\text{eff}} \sim 6000 \text{ K}$, 0.12 dex at $T_{\text{eff}} \sim 5500 \text{ K}$, 0.17 dex at $T_{\text{eff}} \sim 5200 \text{ K}$, and almost 0.3 dex at $T_{\text{eff}} \sim 4900 \text{ K}$.

All the Be abundances measured in this work are listed in Table 1. We have also added the Be abundance measurements

of the 44 stars hosting planets and 29 “single” stars from previous studies (García López & Pérez de Taoro 1998; Santos et al. 2002, 2004b,c), listed in Tables 2 and 3. We have compiled Li abundance measurements, obtained from the $\text{Li I } \lambda 6708 \text{ \AA}$ line, from the literature (Israelian et al. 2004, 2009). We also measure new Li abundances in four objects (three giants and a subgiant, see Table 1) in the same way as in (Israelian et al. 2004, 2009). In Fig. 2 we display some synthetic spectral fits to the observed spectra of four stars of our sample.

5.1. Ages

We have gathered together the ages of some of our stars hosting planets from Saffe et al. (2005). Saffe et al. (2005) studied the correlations between stellar properties with age. They measured the chromospheric activity in a sample of 49 stars with planetary companions and combining with the literature, they obtained age estimates for 112 objects. They applied the calibrations reported in Donahue (1993) and Rocha-Pinto & Maciel (1998) for the chromospheric activity-age relation but also other methods based on isochrones, lithium abundances, metallicities, and kinematics, and they compared them with the chromospheric results. They concluded that chromospheric activity and isochrone methods give comparable results, and claimed that isochrone technique is, in practice, the only tool currently available to derive ages for the complete sample of planet-host stars, as chromospheric activity calibrations of Donahue (1993) are limited for ages < 2 Gyr and most of the planet-host stars sample are > 2 Gyr age. Lithium cannot be used to derived stellar ages greater than 1 Gyr. Here, we have adopted the ages derived from isochrones when available for objects with age < 2 Gyr, otherwise we used those derived using the chromospheric activity technique with Donahue (1993) calibrations. The adopted ages are provided in Table 1. We note that the typical uncertainty on the age determination in field stars is very large, between 1 and 3 Gyr, especially for dwarf stars.

6. Discussion

6.1. Be and Li abundance versus effective temperature and mass

In Fig. 3 we plot the derived Be abundances as a function of effective temperature (top panel) and as a function of mass (bottom panel) for the stars in our sample together with those for the stellar samples reported in García López & Pérez de Taoro (1998); Santos et al. (2002, 2004b,c). We also distinguish among stars with and without planets and among different luminosity classes, including dwarfs (V), subgiants (IV) and giants (III). We note that we did not find in the literature any determination of the stellar mass for all stars in this study, as for instance, no stellar mass was available for subgiant and giant planet-host stars with $T_{\text{eff}} < 4900$ K.

Subgiant and especially giant stars have probably changed their effective temperatures considerably from their “initial” value on the main sequence and their convective envelopes have also expanded and reached deeper and hotter regions in the stellar interiors. Extra mixing may have already occurred and thus, the dilution and/or depletion of their light elements do not follow the normal trend. Therefore, we will take evolved objects into account separately (see Sect. 6.1.2).

Considering only dwarf stars, the addition of the new Be abundance of 26 planet-host stars in Fig. 3 shows no significant changes to what Santos et al. (2004c) found. The overall impres-




Fig. 5. Same as Fig.3 but for Li abundance measurements taken from Israelian et al. (2004, 2009).

sion is that no clear difference seems to exist between the two stellar populations, planet hosts and “single” stars.

Be abundances decrease from a maximum near $T_{\text{eff}} = 6100$ K towards higher and smaller temperatures, similarly to the behaviour of Li abundances versus effective temperature (see Fig. 5). The steep decrease with increasing temperature resembles the well known Be gap for F stars (e.g. Boesgaard & King 2002), while the decrease of the Be content towards lower temperatures is smoother, and may show evidence for continuous Be burning during the main-sequence evolution of these stars (see Santos et al. 2004c, and references therein).

In the top panel of Fig. 3 we have also posed a set of Yale beryllium-depletion isochrones from Pinsonneault et al. (1990). We depict the depletion isochrones (case A) for solar metallicity and an age of 1.7 Gyr, the standard model and 4 models with different initial angular momentum (see Table 3-6 of Pinsonneault et al. 1990) assuming an initial $\log N(\text{Be}) = 1.26$ (intermediate value between solar, 1.10, and meteoritic, 1.42) for all the stars (see Santos et al. 2004b, and references therein). As already noticed in Santos et al. (2004b,c), these models agree with the observations above roughly 5600 K, but while the observed Be abundance decreases towards lower temperatures when $T_{\text{eff}} < 5600$ K, these models predict either constant or increasing Be abundances. Several possibilities were also discussed for this discrepancy between models and Be observations: the increasing difficulty to measure Be abundances at low temperatures; the lack of near-UV line-opacity in the spectral synthesis; the presence of the planetary companions; possible ac-

cretion of $\approx 0.5M_{\odot}$ by stars in solar neighborhood (Murray et al. 2001); and possible correlation with oxygen abundances.

All of them dismissed as the main possible cause although we will further discuss the last one in Sect. 6.4. The better explanation is probably that these models do not predict correctly the behaviour of the Be abundances at the lowest temperatures.

Mixing by internal gravity waves may provide an explanation to the Li and Be depletion in cool dwarf stars (García Lopez & Spruit 1991; Montalbán 1994; Montalbán & Schatzman 1996; Charbonnel & Talon 2005). In Fig. 3 we also depict one model, that includes mixing by gravity waves, from Montalbán & Schatzman (2000). Santos et al. (2004b) already pointed out that models including mixing by internal waves (Montalbán & Schatzman 2000), reproduce the decrease of Be content in the lower temperature regime although still overestimate the Be abundances with respect to the observed values.

In the bottom panel of Fig. 3 we display the Be abundances as a function of the stellar masses. The Be abundances show, as expected, a similar trend as compared to that of the top panel of Fig. 3. However, some stars move with respect to the other stars in the figure, specially the evolved stars. As already mention in Sect. 6.1, the four giants and one subgiant with $T_{\text{eff}} < 4900$ K do not have available mass determinations. A priori, one would expect to find the subgiant stars at high masses, but this picture changes when having stars with different metallicities. Some subgiant stars with planets stay with relatively low masses due to their low metal content, whereas other subgiants, like HD 38529 with a mass of $1.6 M_{\odot}$ and $[\text{Fe}/\text{H}] = 0.4$, and with an upper-limit Be measurement, have very high masses partially due to their high metallicity. In the following, we will concentrate on the Be abundances in main-sequence stars.

6.1.1. Main-sequence stars

In top panel of Fig. 4 we display Be abundances as a function of effective temperature only for unevolved stars with and without known planetary companion. Here we have excluded the stars with Be abundances below 0.4 dex which lie in some kind of Be-gap (see Santos et al. 2004b). We perform a linear fit, also overplotted in Fig. 4, and get a correlation coefficient of 0.74 with standard deviation of 0.15. If one discriminates between the planet hosts and the comparison sample dwarf stars, no significant difference in slope is found when fitting all stars in the whole temperature range.

In Fig. 4 (top panel) one can see that whereas in the range $5700 \lesssim T_{\text{eff}} \lesssim 6200$ K, the stars with and without planets mostly show similar abundances, at the temperature range $5100 \lesssim T_{\text{eff}} \lesssim 5500$ K there seems to be some indication that planet-host stars may be more Be depleted than “single” stars.

Israeli et al. (2009), based in the Li study of unbiased sample of solar-analogue stars with and without detected planets, find that planet host have lower Li abundances, being most of them upper-limit measurements, than “single” stars, suggesting that the presence of planets may increase the amount of mixing and deepen the convective zone to such an extent that the Li can be burned.

If this is true, we should see the same effect at lower temperatures in Be burning regime. This may be reflected in the apparent overabundance of Be in “single” stars respect to planet host stars at $5100\text{--}5500$ K T_{eff} range, in the sense that most of the Be measurements in planet hosts are upper-limits.

However, the sample with T_{eff} below 5500 K is still too small. In addition, as shown in Sect. 5, the expected error on

the Be abundances in this T_{eff} range is 0.12–0.17 dex. Therefore, this calls for new observations to add more Be measurements in dwarf stars at these cool temperatures.

In Fig. 4 (bottom panel) one can see that many points have move to a different relative position according to the effective temperature due to the different metallicity of the stars with and without planets. In particular, the stars without planets which on average have lower metallicity appear to have lower masses. So, the previous picture with the points well concentrated in a relatively narrow T_{eff} range, are now spread in a relatively larger mass range. In particular, the main-sequence stars with planets in the T_{eff} range $5100\text{--}5500$ K have a mean metallicity of $[\text{Fe}/\text{H}] \sim 0.18$ (with a $\sigma_{\text{Fe}} = 0.22$) whereas “single” stars have a mean $[\text{Fe}/\text{H}] \sim -0.22$ ($\sigma_{\text{Fe}} = 0.20$). This explains why the slight signature in stars with and without planets seen in the T_{eff} plot gets smudged in the $\log(M/M_{\odot})$ plot. We note that the star BD-103166, at $T_{\text{eff}} \sim 5320$ K and with $\log N(\text{Be}) < 0.5$, is not display in Figs. 3 and 4. We did not find any mass determination for this star, but we may estimate its mass to be roughly $0.91 M_{\odot}$, and thus having $\log(M/M_{\odot}) \sim -0.04$.

On the other hand, the stellar masses trace the stellar positions at the beginning of the stellar life and not at their present states, which are established by the stellar temperatures. As we notice in Sect. 6.3, the stellar metallicity may have an impact on the Be burning rate but we do not expect that this explains why most of the Be measurements at the T_{eff} range $5100\text{--}5500$ K in planet hosts are only upper-limits. Unfortunately, we cannot track the effect of metallicity in the pre-main sequence and main-sequence evolution of Be abundances from theoretical models at different metallicities since most of the standard models do not predict any Be depletion at these cool temperatures irrespective of the metal content of the star (see e.g. Siess et al. 2000).

In Fig. 5 we display the Li abundances of the stars in the sample versus effective temperature (top panel) and versus mass (bottom panel). All the stars in this plot follow the general trend: stars with low Be abundances have also their Li severely depleted. This general trend seen in the top panel of Fig. 5, is less clear in the bottom panel due to the larger spread of the Li abundance measurements in the $\log(M/M_{\odot})$ axis. However, there is a small group of both stars with and without planets at temperatures $T_{\text{eff}} \lesssim 5670$ K which show relatively high values of Li abundance, $\log N(\text{Li}) > 1.2$, whereas their Be abundances are close to the solar value, $\log N(\text{Be}) \sim 1.1$. Two of these stars are dwarf planet-host stars: HD 1237, with a relatively high chromospheric activity index, $\log R_{\text{HK}} = -4.496$ (Gray et al. 2006), classified as an active star, and may be a young object, with an age below 1Gyr; and HD 65216, with a slightly high Li abundance, $\log N(\text{Li}) = 1.28$, and a low chromospheric index, $\log R_{\text{HK}} = -4.92$. We note that the Sun has $\log R_{\text{HK}} \sim -4.75$. Among these stars there are also four “single” dwarf stars: three of them, HD 36435, HD 43162, and HD 74576, with also high chromospheric activity indexes, $\log R_{\text{HK}} = -4.499, -4.480$ and -4.402 , respectively. However, the other object, HD 43834, show a relatively low activity ($\log R_{\text{HK}} = -4.940$). This star, with a $T_{\text{eff}} = 5594$ K, has a “normal” Be abundance, $\log N(\text{Be}) = 0.94$ dex and a high Li abundance, $\log N(\text{Li}) = 2.30$ dex, for its temperature, according to the general trend. This makes this star interesting in the context of pollution. We refer here the studies by Santos et al. (2004b,b) where these four “single” stars have been already discussed.

It is also worth remarking one planet-host dwarf star of our sample, HD 142, with a temperature $T_{\text{eff}} \approx 6400$ K, with higher Be content than expected from the typical trend, being nearly as high as the maximum at $T_{\text{eff}} \sim 6100$ K. This star also has a




Fig. 6. Be versus Li abundances of the stars in the sample with and without planets. Different colors represent different luminosity classes. The stars are splitted into different temperature ranges associated with different symbols.




Fig. 7. Same as Fig. 6 but for Be abundance versus metallicity.

higher Li abundance than expected from its effective temperature (see Fig. 5). We also note that the Be abundance maximum may need to be slightly shifted towards lower temperatures by one of the planet-host stars presented in this work, HD 213240, which is located at $T_{\text{eff}} = 5982$ K and have $\log N(\text{Be}) \sim 1.4$. In other words, one would have to say that stars with maximum Be abundances have temperatures in the range $5950 \leq T_{\text{eff}} \leq 6150$ K.

The simplest explanation may be that HD 142, classified as F7V by SIMBAD⁴, is just a young star which may be supported by its derived and relatively low surface gravity, $\log g = 4.62$. However, according to the age estimates in Eggenberger et al. (2007), this star has an age of ≈ 2.8 -5.93 Gyr which still makes lithium content too high. In addition, this star has a relatively low chromospheric activity index $\log R_{\text{HK}} = -4.853$ (Gray et al. 2006), which may indicate that, in fact, the age determined by Eggenberger et al. (2007) may be correct. Gray et al. (2006) classified HD 142 as a chromospherically inactive star. HD 142 lies in the called Be "gap" according to its derived effective temperature of $T_{\text{eff}} \sim 6400$ K, but has nearly keep all its Be. In the literature one can find stars with similar or even higher Be content but typically at $T_{\text{eff}} \leq 6350$ K, for instance, in the Hyades

⁴ The SIMBAD database can be accessed at <http://simbad.u-strasbg.fr/simbad/>

cluster with an age of 600 Myr (Boesgaard & King 2002) and in the intermediate-age open cluster IC 4651 (Smiljanic et al. 2010) with an age of 1.7 Gyr, both with a metallicity of ~ 0.1 dex, very similar to the metallicity of HD 142. We note, however, that the mass of HD 142, $1.29 M_{\odot}$, places this star very close to other main-sequence stars in the bottom panel of Figs. 3 and 4.

New observations of Be in stars with and without planets are also needed in this temperature range to study possible pollution effects. We note that the depletion rates may be different for different proto-planetary disc masses and composition, and therefore, the rotational history of the star (e.g. Balachandran 1995; Chen et al. 2001). Thus we will need to find targets with similar conditions, i.e. temperature, age, etc.

6.1.2. Evolved stars

Following the criterion established in Sect. 3, we have four objects clearly classified as giants and eight that could also have evolved off the main sequence.

Five of these subgiants have Li abundance measurements and four of them (the planet hosts HD 10697, HD 88133 and HD 117176 and the "single" star HD 23249) show relatively high values, with respect to the Li trend described by most of the dwarf stars. However, these four subgiant stars have Be abundances consistent with the Be trend shown in dwarf stars (see Fig. 3).

HD 10697 and HD 117176, as well as the "single" star HD 23249 were already discussed in Santos et al. (2002, 2004c). These authors discussed the possibility that pollution by invoking planet engulfment (see e.g. Israelian et al. 2001, 2003), at least for the two planet hosts, could explain the high Li content and the relative "normal" Be content (Siess & Livio 1999). However, recent models show that many accretion events of planetary material could cause Li depletion instead of Li enhancement (see e.g. Théado et al. 2010; Baraffe & Chabrier 2010).

The high Li content and also moderate Be in the metal-rich planet host HD 88133, $[\text{Fe}/\text{H}] = 0.33$, may be also explained by pollution effects.

The four giant stars, HD 47536, HD 59686, HD 219449, and HD 27442, are all of them planet hosts and only present upper limits for Li and Be.

6.2. Beryllium versus lithium

A beryllium versus lithium diagram can provide information on their different depletion rates in main-sequence stars. We have already introduced the relationship between Be and Li abundance in the previous Sect. 6.1, but here, we will focus on Fig. 6, which depicts the Be abundances versus Li abundances of the stars in our sample. In general, there seems to be almost constant relation between the Be and Li abundances in dwarf stars, except for very few dwarfs in the comparison sample. However, both stars with and without planets seem to follow the same behaviour.

In Fig. 6 we also split the Be and Li measurements in several stellar temperature ranges. Thus, the objects with $T_{\text{eff}} \geq 6000$ K are situated in the upper right corner, and define the abundance maximum, for both Be and Li, and follow a positive correlation. The stars with temperatures between 5600 and 6000 K are situated in the middle of the diagram, in a nearly flat Be abundance through Li abundances ranging from 0.05 to 2.5 dex. The objects with temperatures between 5600 and 5100 K are mostly in




Fig. 8. Same as Fig. 6 but for Be abundance versus age.

the left upper-middle panel, also follow an almost constant Be abundance at $\log N(\text{Be}) \sim 0.9$ with only upper-limit Li measurements. This temperature range involve all those stars where Be depletion has already taken place and therefore Li is severely depleted. Thus, this group shows a Be level lower than that at higher temperatures. Nevertheless, 5 objects in this temperature regime (two of them are subgiants), both with and without planetary companions, are located in the right upper side showing both high levels of Be and Li abundances. Finally, we separate the objects with temperatures below 5100 K as Be abundances cannot be determined accurately. These stars can be in turn divided in three sets: (i) low Be and Li abundances that correspond to the giant stars; (ii) low Li abundances but still relatively high Be abundances; and (iii) intermediate abundances of both Li and Be. The latter contains two objects, one of them being a subgiant (see Sect. 6.1.2).

Fig. 6 confirms what found in Santos et al. (2004c), that Be and Li burning seem to follow the same trend for the hottest and coolest objects. Li and Be are depleted in a similar way up to 5600 K, where Be burning stops while Li burning continues up to ~ 6000 K, where again both Li and Be are destroyed in a similar way. Li burning starts to be severe from $T_{\text{eff}} \lesssim 5900$ K while Be keeps its low-burning rate until $T_{\text{eff}} \sim 5600$ K. Giant stars show both Li and Be severely depleted with exception of one object, HD 27442, that still keeps a relatively high upper limit to its Be abundance.

In general, Li is more depleted than Be in stars with smaller effective temperatures and all stars with and without planetary companions show a similar behaviour. With the addition of the 26 planet-host stars and one “single” star, the apparent “lack” of planet-host stars in the range of Li abundances $\log N(\text{Li}) \sim 1.5 - 2.5$, noted by Santos et al. (2004c), has almost disappeared. There seems to be roughly the same number of stars with and without planets in this range of Li abundances, but this might also be due to the small number of stars without planets, 30, in this sample, in comparison with the 70 planet-host stars.

6.3. Be abundance versus $[\text{Fe}/\text{H}]$ and age

In Fig. 7 we display the Be abundances versus the metal content of the stars with and without planets. We do not see any clear correlation but just an increasing dispersion of the Be content towards higher metallicities. The differences in the Be content are mainly due to the differences in T_{eff} , being lower at lower




Fig. 9. *Top:* Be versus $[\text{O}/\text{H}]$ abundances in dwarf stars hosting planets in the sample and the best linear fit. *Bottom:* Be abundance corrected using the linear relation $\text{Be}-T_{\text{eff}}$.

temperatures, but roughly constant for all metallicities. This was already noticed by Santos et al. (2004c).

At the highest metallicities, the opacity of the convective zone changes considerably, producing a higher rate of Li burning and also, although less significant, Be burning. This might explain why the Be abundance in planet-host stars has a maximum at $[\text{Fe}/\text{H}] \sim 0.20$ and then decreases towards higher metallicities. On the other hand, Galactic Be abundances are known to increase with the metallicity (e.g. Rebolo et al. 1988; Molaro et al. 1997; Boesgaard et al. 1999), which in fact is seen in Fig. 7, although the dispersion in Be abundances and the effective temperature spread is probably masking this Galactic effect (see also Santos et al. 2004c, and references therein).

In Fig. 8 we display the Be abundances versus ages of the planet-host stars including those from the literature. We have objects in the same temperature regime but with different ages. There is no correlation, indicating that the age, as well as the metallicity, is a secondary effect on the level of the Be content behind a primary effect driven by the effective temperature of the star. However, any interpretation from this plot must be taken with caution since the age determinations, specially for main-sequence stars, have large error bars and may not be reliable.

6.4. Be versus $[\text{O}/\text{H}]$

Beryllium is mainly produced in the spallation of CNO nuclei (see e.g. Tan et al. 2009, and references therein). In particular,

oxygen gives the largest contribution to the spallation process responsible for forming Be. Thus, a comparison between O and Be abundances can provide alternative information on the behaviour of beryllium. We have collected the oxygen abundances in stars hosting planets of our sample from Ecuivillon et al. (2006). They found average oxygen abundances, $[O/H]$, 0.1-0.2 dex higher in stars hosting planets with respect to stars without planets. However, although planet accretion was not excluded, the possibility that it is the main source of the observed oxygen enhancement in planet-host stars is unlikely. Thus, it seems to be related to the higher metal content of planet-harboring stars and both Galactic trends of stars with and without planet are nearly indistinguishable.

In (top panel) of Fig. 9 we display the beryllium versus oxygen abundances relative to the Sun only in dwarf planet-host stars. We took the $[O/H]$ values derived from the near-UV OH bands in the spectral range 3167–3255 Å (Ecuivillon et al. 2006), because these lines are located close to the Be_{II} 3131 Å line and therefore would be affected for the same continuum opacity. We note that the stellar parameters used in this work to derive Be abundances are not exactly the same as those used by Ecuivillon et al. (2006) to derive oxygen abundances, but in general the differences are very small. We adopted the mean value derived from the four OH features as given in Ecuivillon et al. (2006). We also note that for some stars in Fig. 3 there is no available O measurement.

The Be content is stratified in different values as the temperature changes. In the temperature range 5100–5600 K Be abundance versus $[O/H]$ seems to have a positive slope, just due to one dwarf star located at $[O/H] \sim -0.25$. However, all these trends are affected by the strong influence of the effective temperature on the Be abundances in each star, irrespective of metallicity and oxygen abundance.

In the top panel of Fig. 9, similarly to Fig. 4, we have also displayed the best linear fit, showing a correlation of 0.44 with a standard deviation of 0.18. To study the influence of the effective temperature in this Be-O relation, we subtract from each Be measurement the Be content given by the $Be-T_{eff}$ linear relation in Fig. 4. We thus obtain the Be abundance “corrected” from the effect of the effective temperature, $\log N(Be) - \log N(Be(T_{eff}))$, which is shown in the bottom panel of Fig. 9. The Be content does not seem to depend much on the initial oxygen abundance of the star, as the trend disappears when removing the T_{eff} effect.

7. Summary and Conclusions

We present here new Be abundances of 26 stars harbouring planets and one star without detected planets, from unpublished UVES spectra, added to the previous sample of García López & Pérez de Taoro (1998); Santos et al. (2002, 2004b,c). We have also compiled Li abundances for these objects. The complete sample contains 100 objects, including dwarfs, subgiants and giants, with 70 stars hosting planets. We study the behaviour of Be abundances of the stars in the sample with effective temperature, Li abundance, metallicity, age, oxygen abundance and through different spectral types and luminosity classes.

In general, the work concludes that the beryllium content in stars with and without planetary companion is similar and behaves in the same way. As the burning process is sensitive to the mixing mechanism, the presence of a planetary companion is not necessary affecting this process in a notorious way, at least, at $T_{eff} \gtrsim 5500$ K.

Below $T_{eff} \approx 5500$ K, the stellar sample is still small but appears to be an increase in the dispersion of the Be abundances of dwarf stars. In addition, most of the planet-host Be measurements are not indeed measurements but upper limits. However, there is no way to check if the apparent higher scatter of Be content at $T_{eff} \lesssim 5500$ K is due to the different stellar metallicities since most of the standard theoretical models predict no Be depletion at these temperatures for any metallicity.

Be depletion depends on the effective temperature more than on the age and metal content. The fact that the stars which host planets are richer in different metal contents than “single” stars, but have the same Be content, supports the idea of primordial origin of these over-abundances in planet hosts.

Acknowledgements. M.C.G.O. acknowledges financial support from the European Commission in the form of a Marie Curie Intra European Fellowship (PIEF-GA-2008-220679) and the partial support by the Spanish MICINN under the Consolider-Ingenio 2010 Program grant CSD2006-00070: First Science with the GTC (<http://www.iac.es/consolider-ingenio-gtc>). N.C.S. would like to thank the support by the European Research Council/European Community under the FP7 through a Starting Grant, as well from Fundação para a Ciência e a Tecnologia (FCT), Portugal, through a Ciência2007 contract funded by FCT/MCTES (Portugal) and POPH/FSE (EC), and in the form of grant reference PTDC/CTE-AST/098528/2008 from FCT/MCTES. J.I.G.H. thanks financial support from the Spanish Ministry project MICINN AYA2008-00695. E.D.M and G.I. would like to thank financial support from the Spanish Ministry project MICINN AYA 2008-04874. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This work has also made use of the IRAF facility.

References

- Allende Prieto, C., & Lambert, D. L. 2000, *AJ*, 119, 2445
 Allende Prieto, C., Barklem, P. S., Lambert, D. L., & Cunha, K. 2004, *A&A*, 420, 183
 Ashwell, J. F., Jeffries, R. D., Smalley, B., Deliyannis, C. P., Steinhauer, A., & King, J. R. 2005, *MNRAS*, 363, L81
 Balachandran, S. C. 1995, *AJ*, 446, 203
 Balachandran, S. C., & Bell, R. A. 1998, *Nature*, 392, 791
 Baraffe, I., & Chabrier, G. 2010, *A&A*, 521, A44
 Bertelli, G., Girardi, L., Marigo, P., & Nasi, E. 2008, *A&A*, 484, 815
 Boesgaard, A. M., Deliyannis, C. P., King, J. R., Ryan, S. G., Vogt, S. S., & Beers, T. C. 1999, *AJ*, 117, 1549
 Boesgaard, A. M., & King, J. R. 2002, *ApJ*, 565, 587
 Bouvier, J. 2008, *A&A*, 489, L53
 Charbonnel, C., & Talon, S. 2005, *Science*, 309, 2189
 Chen, Y. Q., Nissen, P. E., Benon, T., & Zhao, G. 2001, *A&A*, 371, 943
 Chen, Y. Q. and Zhao, G., 2006, *AJ*, 131, 1816
 Donahue, R. A. 1993, PhD. Thesis, New Mexico State University
 Ecuivillon, A., Israelian, G., Santos, N. C., Mayor, M., García López, R. J., & Randich, S. 2004a, *A&A*, 418, 703
 Ecuivillon, A., Israelian, G., Santos, N. C., Mayor, M., Villar, V., & Bihain, G. 2004b, *A&A*, 426, 619
 Ecuivillon, A., Israelian, G., Santos, N. C., Shchukina, N. G., Mayor, M., & Rebolo, R. 2006, *A&A*, 445, 633
 Eggenberger, A., Udry, S., Chauvin, G., Beuzit, J.-L., Lagrange, A.-M., Ségransan, D., & Mayor, M. 2007, *A&A*, 474, 273
 Fischer, D. A., & Valenti, J. 2005, *ApJ*, 622, 1102
 García López, R. J., & Spruit, H. C. 1991, *ApJ*, 377, 268
 García López, R. J., Rebolo, R., & Martín, E. L., 1994, *A&A*, 282, 518
 García López, R. J. & Pérez de Taoro, M. R. 1998, *A&A*, 334, 599
 Gilli, G., Israelian, G., Ecuivillon, A., Santos, N. C., & Mayor, M. 2006, *A&A*, 449, 723
 Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, *A&AS*, 141, 371G
 Gonzalez, G., Laws, C., Tyagi, S., & Reddy, B. E. 2001, *AJ*, 121, 432
 Gonzalez, G., 2008, *MNRAS*, 386, 928
 Gray, D. F. 1992, *Camb. Astrophys. Ser.*, Vol. 20
 Gray, R. O., Corbally, C. J., Garrison, R. F., McFadden, M. T., Bubar, E. J., McGehee, C. E., O’Donoghue, A. A., & Knox, E. R. 2006, *AJ*, 132, 161
 Haywood, M. 2009, *ApJ*, 698, L1
 Israelian, G., Santos, N. C., Mayor, M., & Rebolo, R. 2001, *Nature*, 411, 163
 Israelian, G., Santos, N. C., Mayor, M., & Rebolo, R. 2003, *A&A*, 405, 753
 Israelian, G., Santos, N. C., Mayor, M., & Rebolo, R. 2004, *A&A*, 414, 601

Table 1. Stellar parameters and Be abundances measured in this work. We measured Li abundances for 4 stars. Li abundances without a label were taken from Israelian et al. (2009).

Name	T_{eff} [K]	$\log g$ [cm s ⁻²]	ξ_t km s ⁻¹	[Fe/H]	Mass M _⊙	$V_{\text{ sini}}$ km s ⁻¹	V	SpT ^d	LC ^e	logN(Be) [dex]	logN(Li) [dex]	[O/H] ^f [dex]	Age ^g Gyr
HD142 ^a	6403	4.62	1.74	0.09	1.29	11.3	5.70	F7V	V	1.31	3.00	0.07±0.13	3.6
HD1237 ^a	5514	4.50	1.09	0.07	0.94	5.5	6.59	G8.5Vk:	V	1.14	2.19	0.07±0.10	0.15
HD4208 ^a	5599	4.44	0.78	-0.28	0.83	2.8	7.79	G7VFe-1H-05	V	0.94	<0.20	-0.14±0.08	4.47
HD23079 ^a	5980	4.48	1.12	-0.12	1.01	3.4	7.12	F9.5V	V	1.10	2.21	0.02±0.09	8.4
HD28185 ^a	5667	4.42	0.94	0.21	0.98	2.5	7.80	G6.5IV-V	V	0.98	<0.26	0.23±0.10	12.2
HD30177 ^b	5588	4.29	1.08	0.39	1.01	2.96 ¹	8.41	G8V	V	0.96	<0.35 ²	0.31±0.12	8.30
HD33636 ^{b*}	6046	4.71	1.79	-0.08	1.16	3.08 ¹	7.06	G0VH-03	V	1.34	2.64	0.02±0.11	8.1
HD37124 ^b	5546	4.50	0.80	-0.38	0.75	1.22	7.68	G4IV-V	V	0.94	<0.31	-0.06±0.08	0.57
HD39091 ^a	6003	4.42	1.12	0.09	1.10	3.3	5.67	G0V	V	1.11	2.39	0.23±0.08	6.0
HD47536 ^b	4554	2.28	1.82	-0.54	-	1.93 ³	5.26	K0III	III	<-1.32	<-0.46 ³	-0.29±0.12	-
HD50554 ^b	6026	4.41	1.11	0.01	1.09	3.88 ¹	6.84	F8V	V	1.18	2.57	0.17±0.09	7.0
HD59686 ^c	4871	3.15	1.85	0.28	-	0.96 ³	5.45	K2III	III	<-0.75	<0.07 ³	0.14±0.16	-
HD65216 ^a	5612	4.44	0.78	-0.17	0.87	2.3	7.97	G5V	V	1.05	1.28	0.00±0.08	-
HD70642 ^a	5668	4.40	0.82	0.18	0.98	2.8	7.18	G6VCN+05	V	0.90	<0.46	0.17±0.09	10.2
HD72659 ^b	5995	4.30	1.42	0.03	1.16	2.21 ¹	7.46	G0V	V	1.25	2.38	0.21±0.09	8.2
HD73256 ^a	5526	4.42	1.11	0.23	0.97	3.1	8.08	G8IV-VFe+05	V	0.75	<0.54	0.11±0.11	15.9
HD74156 ^b	6112	4.34	1.38	0.16	1.27	4.32	7.61	G0	V	1.33	2.75	0.32±0.08	3.2
HD88133 ^c	5438	3.94	1.16	0.33	0.98	2.17 ¹	8.06	G5IV	IV	0.65	1.91 ³	0.22±0.09	9.56
HD99492 ^c	4810	4.21	0.72	0.26	-	1.36 ¹	7.57	K2V	IV	<-0.22	<-0.01	0.01±0.08	4.49
HD106252 ^b	5899	4.34	1.08	-0.01	1.02	1.93 ¹	7.36	G0V	V	0.95	1.66	0.09±0.07	9.2
HD114729 ^a	5844	4.19	1.23	-0.28	0.94	2.3	6.69	G0V	V	0.99	2.00	-0.01±0.07	11.9
HD117207 ^a	5667	4.32	1.01	0.22	0.98	1.8	7.26	G7IV-V	V	0.94	<0.12	0.19±0.09	16.1
HD117618 ^a	5990	4.41	1.13	0.03	1.08	3.2	7.17	G0V	V	1.10	2.29	0.14±0.09	6.72
HD213240 ^a	5982	4.27	1.25	0.14	1.21	4.0	6.80	G0/G1V	V	1.41	2.54	0.25±0.08	3.60
HD216435 ^a	6008	4.20	1.34	0.24	1.33	5.9	6.03	G0V	V	1.26	2.77	0.23±0.12	5.40
HD216437 ^b	5887	4.30	1.31	0.25	1.20	2.6	6.06	G1VFe+03	V	1.32	1.96	0.22±0.10	8.7
HD219449 ^c	4757	2.71	1.71	0.05	-	5.10	4.21	K0III	III	<-1.1	<-0.16 ³	-0.10±0.13	-

^a Values taken from Sousa et al. (2008).^b Values taken from Santos et al. (2004a).^c Values taken from Santos et al. (2005).^d Values taken from SIMBAD.^e Luminosity class assigned by comparing the stellar parameters with isochrones from Girardi et al. (2000)^f Values taken from Ecuivillon et al. (2006)^g Values taken from Saffe et al. (2005)¹ Data taken from Fischer & Valenti (2005).² Data taken from Israelian et al. (2004).³ Data measured in this work.

* stars without planet companion.

- Israelian, G., Delgado Mena, E., Santos, N. C., Sousa, S. G., Mayor, M., Udry, S., Domínguez Cerdeña, C., Rebolo, R., & Randich, S. 2009, *Nature*, 462, 189
- Kurucz, R. 1993, *ATLAS9 Stellar Atmosphere Programs and 2 km/s grid*. Kurucz CD-ROM No. 13. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993, 13
- Laws, C., Gonzalez, G., Walker, K. M., Tyagi, S., Dodsworth, J., Snider, K., & Szentzeff, N. B. 2003, *AJ*, 125, 2664
- Melo, C. H. F., de Laverny, P., Santos, N. C., Israelian, G., Randich, S., Do Nascimento, J. D., Jr., & de Medeiros, J. R. 2005, *A&A*, 439, 227
- Molaro, P., Bonifacio, P., Castelli, F., & Pasquini, L. 1997, *A&A*, 319, 593
- Montalbán, J. 1994, *A&A*, 281, 421
- Montalbán, J., & Schatzman, E. 1996, *A&A*, 305, 513
- Montalbán, J., & Schatzman, E. 2000, *A&A*, 354, 943
- Murray, N., Chaboyer, B., Arras, P., Hansen, B., & Noyes, R. W. 2001, *ApJ*, 555, 801
- Neves, V., Santos, N. C., Sousa, S. G., Correia, A. C. M., & Israelian, G. 2009, *A&A*, 497, 563
- Pasquini, L., Döllinger, M. P., Weiss, A., Girardi, L., Chavero, C., Hatzes, A. P., da Silva, L., & Setiawan, J. 2007, *A&A*, 473, 979
- Pinsonneault, M. H., Kawaler, S. D., & Demarque, P. 1990, *ApJS*, 74, 501
- Randich, S., Aharpour, N., Pallavicini, R., Prosser, C. F., & Stauffer, J. R. 1997, *A&A*, 323, 86
- Randich, S., Primas, F., Pasquini, L., & Pallavicini, R. 2002, *A&A*, 387, 222
- Rebolo, R., Abia, C., Beckman, J. E., & Molaro, P. 1988, *A&A*, 193, 193
- Rocha-Pinto, H.-J., & Maciel, W.-J. 1998, *MNRAS*, 298, 332
- Saffe, C., Gómez, M., & Chavero, C. 2005, *A&A*, 443, 609
- Santos, N. C., Israelian, G., & Mayor, M. 2001, *A&A*, 373, 1019
- Santos, N. C., García López, R. J., Israelian, G., Mayor, M., Rebolo, R., García-Gil, A., Pérez de Taoro, M. R., & Randich, S. 2002, *A&A*, 386, 1028
- Santos, N. C., Israelian, G., Mayor, M., Rebolo, R., & Udry, S. 2003, *A&A*, 398, 363
- Santos, N. C., Israelian, G., & Mayor, M. 2004a, *A&A*, 415, 1153
- Santos, N. C., Israelian, G., Randich, S., García López, R. J., & Rebolo, R. 2004b, *A&A*, 425, 1013
- Santos, N. C., Israelian, G., García López, R. J., Mayor, M., Rebolo, R., Randich, S., Ecuivillon, A., & Domínguez Cerdeña, C. 2004c, *A&A*, 427, 1085
- Santos, N. C., Israelian, G., Mayor, M., Bento, J. P., Almeida, P. C., Sousa, S. G., & Ecuivillon, A., 2005, *A&A*, 437, 1127
- Siess, L., & Livio, M. 1999, *MNRAS*, 308, 1133
- Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 593
- Smiljanic, R., Pasquini, L., Charbonnel, C., & Lagarde, N. 2010, *A&A*, 510, A50
- Smiljanic, R., Pasquini, L., Bonifacio, P., Galli, D., Barbuy, B., Gratton, R., & Randich, S. 2009, *A&A*, 499, 103
- Snedden, C. 1973, PhD Dissertation, Univ. of Texas, Austin
- Sousa, S. G., et al. 2008, *A&A*, 487, 373
- Takeda, Y., Kawanomoto, S., Honda, S., Ando, H., & Sakurai, T. 2007, *A&A*, 468, 663
- Tan, K. F., Shi, J. R., & Zhao, G. 2009, *MNRAS*, 392, 205
- Théado, S., Bohuon, E., & Vauclair, S. 2010, *IAU Symposium*, 268, 427

Table 2. Stellar parameters and Be and Li abundances from literature (García López & Pérez de Taoro 1998; Santos et al. 2002, 2004c; Sousa et al. 2008): Stars hosting planets.

Name	T_{eff} [K]	$\log g$ [cm s^{-2}]	[Fe/H]	Mass ^a M_{\odot}	SpT ^b	LC ^c	$\log N(\text{Be})$ [dex]	$\log N(\text{Li})$ [dex]	[O/H] ^d [dex]	Age ^e Gyr
BD-103166	5320	4.38	0.33	-	K0V	V	<0.5	-	-	5.39
HD6434	5835	4.6	-0.52	0.82	G2/G3V	V	1.08	<0.85	-0.18±0.1	13.3
HD9826	6212	4.26	0.13	1.30	F8V	V	1.05	2.55	0.22±0.12	-
HD10647	6143	4.48	-0.03	1.20	F9V	V	1.19	2.8	-0.06±0.09	4.8
HD10697	5641	4.05	0.14	1.22	G5IV	IV	1.31	1.96	-	1.17
HD12661	5702	4.33	0.36	1.05	K0V	V	1.13	<0.98	-	7.05
HD13445	5163	4.52	-0.24	0.70	K1V	V	<0.4	<-0.12	-0.25±0.12	-
HD16141	5801	4.22	0.15	1.06	G5IV	V	1.17	1.11	0.19±0.10	11.2
HD17051	6252	4.61	0.26	1.26	F9VFe+03	V	1.03	2.66	0.33±0.11	1.47
HD19994	6190	4.19	0.24	1.37	F8V	V	0.93	1.99	0.31±0.12	4.7
HD22049	5073	4.43	-0.13	0.85	K2V	V	0.77	<0.25	-0.23±0.10	-
HD27442	4825	3.55	0.39	-	K2III	III	<0.3	<-0.47	0.07±0.14	-
HD38529	5674	3.94	0.4	1.60	G4V	IV	<-0.1	<0.61	-	5.09
HD46375	5268	4.41	0.2	0.82	K11V	V	<0.8	<-0.02	0.04±0.10	16.4
HD52265	6103	4.28	0.23	1.20	G0III-IV	V	1.25	2.88	0.21±0.10	3.8
HD69830	5410	4.38	-0.03	0.82	K0V	V	0.79	<0.47	-	-
HD75289	6143	4.42	0.28	1.24	F9VFe+03	V	1.36	2.85	0.22±0.11	4.0
HD75732A ^f	5150	4.15	0.29	0.87	G8V	IV	0.55	<0.04	-	-
HD82943	6016	4.46	0.3	1.15	F9VFe+05	V	1.27	2.51	0.29±0.11	3.5
HD83443	5454	4.33	0.35	0.95	K0V	V	<0.7	<0.52	0.12±0.13	2.94
HD92788	5821	4.45	0.32	1.03	G5	V	1.19	1.34	-	9.6
HD95128	5954	4.44	0.06	1.07	G1V	V	1.23	1.83	-	-
HD108147	6248	4.49	0.2	1.26	F8VH+04	V	0.99	2.33	-	1.98
HD114762	5884	4.22	-0.7	0.81	F9V	V	0.82	2.2	-	11.8
HD117176	5560	4.07	-0.06	0.93	G5V	IV	0.86	1.88	-	-
HD120136	6339	4.19	0.23	1.33	F6IV	V	<0.25	-	-	-
HD121504	6075	4.64	0.16	1.13	G2V	V	1.33	2.65	-	1.62
HD130322	5392	4.48	0.03	0.88	K0III	V	0.95	<0.13	-	1.24
HD134987	5776	4.36	0.3	1.03	G5V	V	1.22	<0.74	-	11.1
HD143761	5853	4.41	-0.21	0.95	G0Va	V	1.11	1.46	-0.09±0.08	-
HD145675	5311	4.42	0.43	0.90	K0V	V	<0.65	<0.03	-	7.6
HD168443	5617	4.22	0.06	0.96	G6V	V	1.11	<0.78	-	10.6
HD169830	6299	4.1	0.21	1.42	F7V	V	<-0.4	<1.16	0.22±0.12	2.3
HD179949	6260	4.43	0.22	1.28	F8.5V	V	1.08	2.65	0.26±0.11	2.05
HD186427 ^f	5700	4.35	0.06	0.99	G3V	V	1.3	0.46	-	-
HD187123	5845	4.42	0.13	1.04	G5	V	1.08	1.21	-	7.3
HD192263	4947	4.51	-0.02	0.69	K2V	V	<0.9	<-0.39	-0.17±0.09	0.57
HD195019	5842	4.32	0.08	1.06	G3IV-V	V	1.15	1.47	-	10.6
HD202206	5752	4.5	0.35	1.04	G6V	V	1.04	1.04	0.20±0.09	2.04
HD209458	6117	4.48	0.02	1.15	G0V	V	1.24	2.7	0.03±0.07	6.6
HD210277	5532	4.29	0.19	0.90	G0V	V	0.91	<0.3	0.13±0.12	6.93
HD217014	5804	4.42	0.2	1.05	G5V	V	1.02	1.3	-0.06±0.11	-
HD217107	5646	4.31	0.37	1.02	G8IV	V	0.96	<0.4	0.29±0.13	7.32
HD222582	5843	4.45	0.05	0.97	G5	V	1.14	<0.59	0.09±0.08	11.1

^a Values taken from Sousa et al. (2008) and Santos et al. (2004a)^b Values taken from SIMBAD.^c Luminosity class assigned by comparing the stellar parameters with isochrones from Girardi et al. (2000)^d Values taken from Ecuillon et al. (2006)^e Values taken from Saffe et al. (2005)^f Values taken from García López & Pérez de Taoro (1998)

Table 3. Stellar parameters and Be and Li abundances from literature (see García López & Pérez de Taoro 1998; Santos et al. 2002, 2004c; Sousa et al. 2008): "Single Stars"

Name	T_{eff} [K]	$\log g$ [cm s ⁻²]	[Fe/H]	Mass ^a M_{\odot}	SpT ^b	LC ^c	$\log N(\text{Be})$ [dex]	$\log N(\text{Li})$ [dex]
HD870	5447	4.57	-0.07	0.86	K0V	V	0.8	<0.2
HD1461	5768	4.37	0.17	1.02	G0V	V	1.14	<0.51
HD1581	5956	4.39	-0.14	1.00	F9.5V	V	1.15	2.37
HD3823	5948	4.06	-0.25	1.01	G0VFe-09H-04	IV	1.02	2.41
HD4391	5878	4.74	-0.03	1.11	G5VFe-08	V	0.64	<1.09
HD7570	6140	4.39	0.18	1.20	F9VFe+04	V	1.17	2.91
HD10700	5344	4.57	-0.52	0.63	G8V	V	0.83	<0.41
HD14412	5368	4.55	-0.47	0.73	G8V	V	0.8	<0.44
HD20010	6275	4.4	-0.19	1.33	F6V	V	1.01	2.13
HD20766	5733	4.55	-0.21	0.93	G2V	V	<-0.09	<0.97
HD20794	5444	4.47	-0.38	0.70	G8V	V	0.91	<0.52
HD20807	5843	4.47	-0.23	0.95	G0V	V	0.36	<1.07
HD23249	5074	3.77	0.13	0.83	K0IV	IV	<0.15	1.24
HD23484	5176	4.41	0.06	0.82	K2V _k :	V	<0.70	<0.4
HD26965A	5126	4.51	-0.31	0.67	K1V	V	0.76	<0.17
HD30495	5868	4.55	0.02	1.10	G1.5VH-05	V	1.16	2.44
HD36435	5479	4.61	0.00	0.98	G9V	V	0.99	1.67
HD38858	5752	4.53	-0.23	0.90	G4V	V	1.02	1.64
HD43162	5633	4.48	-0.01	1.00	G6.5V	V	1.08	2.34
HD43834	5594	4.41	0.1	0.93	G7V	V	0.94	2.3
HD72673	5242	4.5	-0.37	0.70	G9V	V	0.7	<0.48
HD74576	5000	4.55	-0.03	0.78	K2.5V _k :	V	0.7	1.72
HD76151	5803	4.5	0.14	1.04	G2V	V	1.02	1.88
HD84117	6167	4.35	-0.03	1.15	F8V	V	1.11	2.64
HD186408 ^d	5750	4.2	0.11	-	G1.5V _b	V	1.1	1.24
HD189567	5765	4.52	-0.23	0.87	G2V	V	1.06	<0.82
HD192310	5069	4.38	-0.01	0.80	K2+v	V	<0.6	<0.2
HD211415	5890	4.51	-0.17	0.93	G0V	V	1.12	1.92
HD222335	5260	4.45	-0.16	0.77	G9.5V	V	0.66	<0.31

^a Values taken from Sousa et al. (2008) and Santos et al. (2004a)^b Values taken from SIMBAD.^c Luminosity class assigned by comparing the stellar parameters with isochrones from Girardi et al. (2000)^d Values taken from García López & Pérez de Taoro (1998)