

LETTER TO THE EDITOR

Observation of H₂O in a strongly lensed *Herschel*-ATLAS source at $z = 2.3$ [★]

A. Omont^{1,2}, R. Neri³, P. Cox³, R. Lupu⁴, M. Guélin³, P. van der Werf⁵, A. Weiß²², R. Ivison¹⁶, M. Negrello⁶, L. Leeuw¹⁹, M. Lehnert¹⁸, I. Smail¹¹, A. Verma²¹, A. J. Baker³⁴, A. Beelen⁷, J. E. Aguirre⁴, M. Baes³¹, F. Bertoldi⁸, D. L. Clements⁹, A. Cooray¹⁰, K. Coppin³³, H. Dannerbauer¹², G. De Zotti¹³, S. Dye¹⁴, N. Fiolet^{1,2,7}, D. Frayer²³, R. Gavazzi^{1,2}, D. Hughes¹⁵, M. Jarvis³⁵, M. Krips³, M. J. Michałowski²⁵, E. J. Murphy²⁸, D. Riechers²⁸, S. Serjeant³⁶, A. M. Swinbank¹¹, P. Temi²⁰, M. Vaccari²⁴, J. D. Vieira²⁸, R. Auld¹⁴, B. Buttiglione²⁶, A. Cava³², A. Dariush^{27,14}, L. Dunne¹⁷, S. A. Eales¹⁴, J. Fritz²⁶, H. Gomez¹⁴, E. Ibar¹⁶, S. Maddox¹⁷, E. Pascale¹⁴, M. Pohlen¹⁴, E. Rigby¹⁷, D. J. B. Smith^{17,35}, J. Bock^{28,29}, C. M. Bradford^{28,29}, J. Glenn³⁰, K. S. Scott⁴, and J. Zmuidzinas^{28,29}

(Affiliations can be found after the references)

Received 18 March 2011 / Accepted 11 April 2011

ABSTRACT

The *Herschel* survey, H-ATLAS, with its large areal coverage, has recently discovered a number of bright, strongly lensed high- z submillimeter galaxies. The strong magnification makes it possible to study molecular species other than CO, which are otherwise difficult to observe in high- z galaxies. Among the lensed galaxies already identified by H-ATLAS, the source J090302.9-014127B (SDP.17b) at $z = 2.305$ is remarkable because of its excitation conditions and a tentative detection of the H₂O 2_{02-1₁₁} emission line (Lupu et al. 2010, ApJ, submitted). We report observations of this line in SDP.17b using the IRAM interferometer equipped with its new 277–371 GHz receivers. The H₂O line is detected at a redshift of $z = 2.3049 \pm 0.0006$, with a flux of 7.8 ± 0.5 Jy km s⁻¹ and a *FWHM* of 250 ± 60 km s⁻¹. The new flux is 2.4 times weaker than the previous tentative detection, although both remain marginally consistent within 1.6 σ . The intrinsic line luminosity and ratio of H₂O(2_{02-1₁₁})/CO(8-7) are comparable with those of the nearby starburst/enshrouded-AGN Mrk 231, and the ratio I(H₂O)/L_{FIR} is even higher, suggesting that SDP.17b could also host a luminous AGN. The detection of a strong H₂O 2_{02-1₁₁} line in SDP.17b implies an efficient excitation mechanism of the water levels that must occur in very dense and warm interstellar gas probably similar to Mrk 231.

Key words. galaxies: high-redshift – galaxies: starburst – galaxies: active – infrared: galaxies – submillimeter: galaxies – radio lines: galaxies

1. Introduction

Gravitationally lensed sources have played an important role in infrared and submillimeter studies of high- z galaxies since the discovery of IRAS FSC10214+4724 (hereafter IRAS F10214; Rowan-Robinson et al. 1991). The studies of this and two other bright strongly-lensed QSOs, APM 08279+5255 (Downes et al. 1999) and the Cloverleaf (H1413+117; Barvainis et al. 1994), demonstrate the utility of using high gravitational magnification to investigate the detailed properties of distant galaxies. These sources allowed pioneering detections of the infrared and submillimeter continuum and lines of CO, HCO⁺, HCN, HNC, CN, and H₂O (see e.g., Solomon & Vanden Bout 2005; Guélin et al. 2007; Riechers et al. 2010, 2011a). These three lensed sources also provided the first few spatially resolved measurements on scales of hundreds of parsecs. Before *Herschel* these sources were without peer, except for SMMJ2135-0102, MM 18423+5938 and SXDF1100.001 (Swinbank et al. 2010; Lestrade et al. 2010; Ikarashi et al. 2010).

Now the wide area surveys from space – especially H-ATLAS and HerMES with *Herschel*, which will observe 570 deg² and 70 deg², respectively (Eales et al. 2010; Oliver et al. 2010) – and from the ground, for example, with the South Pole Telescope (Vieira et al. 2010), are increasing the area of

submillimeter surveys by factors of hundreds over previous surveys. Correspondingly, we expect that the number of strongly lensed submillimeter sources will also increase by a very large factor. From the analysis and intensive follow-up of the five strongest lenses found in the H-ATLAS science demonstration (SDP) field of 14.5 deg² (Negrello et al. 2010, hereafter Ne10), the H-ATLAS team has shown that these lenses are relatively easy to identify. With $S_{500\mu\text{m}} > 100$ mJy and $S_{1.2\text{mm}} > \sim 10$ mJy, their number exceeds that of the unlensed high- z submillimeter sources (Negrello et al. 2007). The number of similar high- z strongly lensed galaxies in the full H-ATLAS survey will be larger than 100.

CO lines are almost the only molecular lines currently detected with the very broad band spectrometers such as Z-Spec (Lupu et al. 2010, hereafter Lu10; Scott et al. 2011) and Zpectrometer (Frayer et al. 2011) and with interferometers (Cox et al. 2011; Riechers et al. 2011b; Leeuw et al., in prep.). Apart from ~ 20 CO-line detections in high- z *Herschel* galaxies, the only other molecule detection reported in these sources is the tentative detection with Z-Spec of the emission line of para-H₂O 2_{02-1₁₁} (rest LST frequency 987.93 GHz) in the $z = 2.305$ source H-ATLAS J090302.9-014127B – hereafter SDP.17b (Lu10). However, the noise is high in this part of the Z-Spec spectrum and, in addition, the line is partially blended with another strong feature that was identified as CO(5-4) emission from the intervening massive lensing galaxy at $z = 0.942$ H-ATLAS J090302.9-014127A – hereafter SDP.17a – as

[★] *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

discussed in Lu10 and Ne10. Here we report new measurements of the H₂O 2₀₂–1₁₁ line in SDP.17b using the IRAM Plateau de Bure interferometer (PdBI), which confirm the line and enable a more detailed study of its properties.

We adopt a cosmology with $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.27$, $\Omega_\Lambda = 0.73$ (Spergel et al. 2003).

2. Water lines in high- z galaxies

Conducting studies of H₂O in high- z galaxies is important. If is not locked in grains, H₂O may be one of the most abundant molecules in the gas. It is known to be a tracer of the dense, warm gas and possibly of strong infrared radiation because its large dipole and high energy levels make its excitation difficult and sensitive to the interstellar conditions (González-Alfonso et al. 2010, hereafter G-A10).

Spectra from the Infrared Space Observatory (ISO) have shown that the far-infrared H₂O lines are prominent in local ultra-luminous infrared galaxies (ULIRGs) and composite AGN/starburst galaxies, such as Mrk 231 and Arp 220, which show a series of H₂O lines in absorption (González-Alfonso et al. 2008, 2004). The recent *Herschel* SPIRE FTS submillimeter spectrum of Mrk 231 provides a wealth of molecular emission lines including high- J CO lines up to $J = 13$, seven rotational lines of H₂O almost comparable in strength to the CO emission lines, as well as other species (OH⁺, H₂O⁺, and HF) which had not been observed before in external galaxies (van der Werf et al. 2010 (hereafter vdW10); G-A10); Fischer et al. 2010). A high intensity of the H₂O lines implies a high H₂O column density in a compact nuclear component, and thus a H₂O abundance approaching 10^{-6} (G-A10). This is most probably the consequence of shocks, cosmic rays, dense hot cores, possibly a moderated XDR (X-ray dominated, see e.g. Meijerink & Spaans 2005) chemistry, and/or an “undepleted chemistry” where grain mantles have evaporated. Therefore a composite model might be necessary to explain the water emission – shocks excited by the mechanical energy of the starburst or perhaps an AGN may enhance the gas phase abundance of water, while a strong infrared radiation field from an intense starburst or AGN (similar to Mrk 231) may be responsible for the high excitation. As stressed by G-A10, the high H₂O/CO ratio makes it unlikely that the H₂O emission originates in classical photon-dominated regions (PDR), since in the Orion Bar, the proto-typical Galactic PDR, CO lines are a factor ≥ 50 stronger than the H₂O lines. Similarly, in the starburst galaxy M82, this ratio is ≈ 40 (Weiß et al. 2010). Cosmic-ray-dominated chemistry (see e.g. Papadopoulos 2010) appears to be excluded because it is unable to heat the gas to sufficiently high temperatures. The favoured model for explaining these extraordinary features implies a high H₂O abundance and the presence of a small star-forming disk, composed of clumps of dense gas exposed to strong ultraviolet radiation, dominating the emission of CO lines up to $J = 8$ (vdW10, G-A10). X-rays from the accreting supermassive black hole in Mrk 231 are likely to contribute significantly to the excitation and chemistry of the inner disk, as shown by the presence of OH⁺ and H₂O⁺ lines.

The number density of galaxies similar to Mrk 231 is expected to be at least two orders of magnitude higher at high than at low redshift (Fabian et al. 2000; Alexander et al. 2005). They should represent a significant fraction of all submillimeter galaxies (SMGs), and thus of lensed SMGs detected by *Herschel*. At high- z , the far-infrared lines of molecular species such as H₂O are redshifted into the atmospheric sub/millimeter windows. For strongly lensed sources, these molecular lines are within the detection reach of present sub/millimeter interferometers such as




Fig. 1. Spectrum of the para H₂O 2₀₂–1₁₁ emission line towards SDP.17b, where the velocity scale is centred on its observed frequency at 298.93 GHz (corresponding to $z = 2.3049$). The rms noise is ~ 4.7 mJy/beam in 31.6 MHz channels. A Gaussian fit to the H₂O spectrum is shown as a solid line while the dotted line shows the underlying dust continuum emission. The H₂O line is clearly asymmetric and not well fit by a Gaussian profile.

the Plateau de Bure Interferometer (PdBI) and future facilities such as ALMA and NOEMA. Tentative detections of H₂O were reported in the Cloverleaf for the 2₂₀–1₁₁ transition (Bradford et al. 2009, Table 2) and IRAS F10214 for the 2₁₁–2₀₂ transition (Casoli et al. 1994). In addition, luminous water masers ($\nu_{\text{rest}} = 22.2$ GHz) were detected in a lensed quasar at $z = 2.64$ (Impellizzeri et al. 2008) and tentatively in F10214 (McKean et al. 2011). The H₂O 1₁₀–1₀₁ line was also detected in absorption at $z = 0.685$ towards B0218+357 by Combes & Wiklind (1997). It is likely that in the near future, the whole set of molecular lines (including the water lines) seen in Mrk 231 and other local sources (vdW10, Fischer et al. 2010) will be detectable with ALMA in high- z lensed galaxies, provided they are comparable to Mrk 231.

3. Observations and results

In order to confirm the detection towards SDP.17b of the redshifted H₂O 2₀₂–1₁₁ emission line, we used the PdBI with six antennae and the new “Band 4” receiver, which covers the frequency range 277–371 GHz. Because the wide-band correlator, WideX, provides a contiguous frequency coverage of 3.6 GHz in dual polarization, it allowed us to include the frequency of 297 GHz at the edge of the bandpass where Lu10 reported a second strong, but partially blended line, which they identified as the CO(5–4) emission of the lensing galaxy SDP.17a at $z = 0.942 \pm 0.004$.

First observations were made in the compact D-configuration on 2011 January 3 in conditions with good atmospheric phase stability (seeing of $0.7''$) and reasonable transparency ($\text{PWV} \leq 0.5$ mm). They were complemented by observations in extended A- and B-configurations in February and March 2011. With a total of ~ 6.2 h on-source integration, a strong signal was detected both in the continuum and in the purported H₂O line (Fig. 1). The dust continuum flux density at 1.0 mm is 32.3 ± 2 mJy, which agrees well with the value derived from Z-Spec by Lu10, and used in the SED fits of Ne10 and Lu10. However, the respective contributions of SDP.17b and SDP.17a to this value remain uncertain. The maximum flux density and integrated intensity of the H₂O line are 29 mJy and 7.8 ± 0.5 Jy km s^{–1}, respectively, with a $FWHM$ of 250 ± 60 km s^{–1}. The line central frequency of 298.93 GHz corresponds to $z = 2.3049 \pm 0.0006$, which is consistent with the value reported by Lu10, 2.308 ± 0.011 .

Table 1. Observed parameters of the H₂O 2₀₂–1₁₁ emission line in SPD.17b

Source	ν_{rest} [GHz]	ν_{obs} [GHz]	z	S_ν [mJy]	ΔV_{FWHM} [km s ⁻¹]	I [Jy km s ⁻¹]	L^a [10 ⁷ L _⊙]	$L^a/10^{10}$ [K km s ⁻¹ pc ²]
SDP.17b	987.93	298.93	2.3049 ± 0.0006	29	250 ± 60	7.8 ± 0.5	85 ± 0.6	2.5 ± 0.2

Notes. Quoted errors are the statistical errors derived from a Gaussian fit to the line profile. ^(a) Equations (1) and (3) of Solomon et al. (1997). The line luminosities are not corrected for the magnification. Typical amplifications of ~10 (or more) are derived for the lensed sources detected by *Herschel* (Ne10).

Table 2. Properties of the para H₂O 2₂₀–1₁₁ emission line in active galaxies

Source	z	$I(\text{H}_2\text{O})^a$ [Jy km s ⁻¹]	$L(\text{H}_2\text{O})^a$ [10 ⁷ L _⊙]	$I_{\text{H}_2\text{O}}^a/I_{\text{CO}}^b$
SDP.17b	2.305	7.8 ± 0.5	85/μ _L	0.5 ± 0.2
Mrk 231 ¹	0.042	718	2.4	0.75 ± 0.23
Cloverleaf ²	2.565	20.3 ± 6.1	21(11/μ _L)	0.4 ± 0.2

Notes. ^(a) H₂O 2₀₂–1₁₁ line. ^(b) I(H₂O 2₀₂–1₁₁)/I(CO[8–7]) from Lu10. ⁽¹⁾ G-A10; ⁽²⁾ Bradford et al. (2009).

Compared to the value previously reported by Lu10 of 19 ± 7 Jy km s⁻¹, the intensity is lower by a factor 2.4 and marginally consistent within 1.6σ. The relatively low angular resolution (≈1′′) did not allow us to study the spatial properties of the signal. The source does not seem to be resolved and it is unlikely that significant flux is missed, either from the H₂O line or the continuum flux, as shown by the consistency of the latter with the Z-Spec value.

Clearly, there is no other strong line in this spectral range with an intensity approaching that of H₂O. In particular, this precludes the presence of a CO(5–4) line of SDP.17a stronger than about 3 Jy km s⁻¹, i.e. 1/10 of the tentative detection reported by Lu10 (29 ± 9 Jy km s⁻¹), in the redshift range 0.922 < z < 0.944. However, emission from SDP.17a in the CO(5–4) line is not ruled out and could be present at a redshift $z > 0.944$, outside of the bandpass of the current observations.

4. Discussion: high-excitation gas

4.1. Properties of SDP.17b

In order to assess the implication of the detection of H₂O emission in SDP.17b, it is important to summarize the current information on the properties of SDP.17b and to consider this source in relation to other submillimeter lensed sources. SDP.17b is one of the five prominent high- z lensed SMGs found by Ne10 in the H-ATLAS SDP field, with an apparent infrared luminosity $L_{\text{IR}} = 4 \times 10^{13} L_{\odot}$ (8–1000 μm, Lu10). The information on the lensing system remains limited because no high-resolution sub/millimeter image is available yet. However, Ne10 suggested that the deflector SDP.17a could consist of two foreground lensing masses at similar high redshifts ($z \sim 0.8$ –0.9). The amplification factor of SDP.17b is still unknown and could reach values of ≈10 or more (Ne10). The infrared luminosity of SDP.17b is thus comparable to typical ULIRGs, including Mrk 231 ($L_{\text{IR}} = 4 \times 10^{12} L_{\odot}$). The mid-infrared photometry of SDP.17b is unknown, but, because it is detected at 100 μm (Ne10), there might be an AGN contribution.

The most remarkable property of SDP.17 is the richness of its 200–300 GHz Z-Spec spectrum (Lu10). Besides the H₂O line, it displays three CO lines ($J = 6$ –5, 7–6(+[CI]) and 8–7) at $z = 2.3$. Despite the relatively low individual S/N ratios (~1.5–3) of these line intensities, their distribution has no clear sign of a turnover up to $J = 8$ –7, pointing to a similarity with both Mrk 231 and the Cloverleaf (Lu10). In fact, the CO spectral line energy distribution of Mrk 231 also has a peak between the CO(5–4) and CO(6–5) transitions, but it has in addition a strong high-excitation component as seen in vdW10. However, there is as yet no information about CO lines above $J = 8$ –7 in SDP.17b. Note that SDP.17b is one of the rare examples among the strongly lensed galaxies detected by *Herschel* (together with HERMESJ 105751.1+573027 (HLSW-01), Scott et al. 2011) that display strong high- J CO lines. While the information about high- J CO lines is still lacking in many of them, several other well studied high- z lensed sources either from H-ATLAS (J091304.9-005344 (SDP.130), Lu10; J142413.9+022304 (SDP.15.141), Cox et al. 2011) or from elsewhere (Weiß et al. 2005; Danielson et al. 2011; Lestrade et al. 2010) have a clear turnover at lower J values.

Nothing is published yet about the results of observations of lower- J CO millimeter lines of SDP.17b. However, several studies are in progress at CARMA, the GBT (with Zspectrometer) and PdBI, including observations of the $J = 1$ –0, 3–2 and 4–3 lines (Leeuw et al., in prep.; Frayer et al., in prep.; Cox et al., in prep.). These results will provide information on both the molecular gas of SDP.17b and the lensing system. It will be interesting to compare the profiles and the spatial distributions of the H₂O and CO emission lines and to study the connection between the warm gas (emitting in H₂O lines) and the colder gas (traced in CO), in particular the cold gas traced by the CO(1–0) line.

The FIRST radio survey (Becker et al. 1995) yields $S_{1.4\text{GHz}} = 464 \pm 145 \mu\text{Jy}$ for SDP.17. From the L_{IR} value and using the definition of Sajina et al. (2008), we find for the IR/radio parameter $q = 2.53$. This value is well within the range of q values for $z \sim 2$ starburst sources (e.g. Sajina et al. 2008; Fiolet et al. 2009; Ivison et al. 2010). This suggests that SDP.17b is not a radio-loud source and that star formation is responsible for most of its far-infrared luminosity (similar conclusions could apply to SDP.17a). However, it would be important to check whether the actual radio spectral index could suggest that some fraction of the radio emission is powered by an AGN, as in the case of SPD.81 (H-ATLAS J090311.6+003907; Valtchanov et al. 2011).

4.2. Implication of H₂O emission in SDP.17b

As discussed in Sect. 2, the detection of H₂O in SDP.17b implies special excitation conditions in an intense infrared radiation field and a warm dense gas, similar to Mrk 231 (e.g. with $n(\text{H}_2) \sim 10^6 \text{ cm}^{-3}$ and $T_{\text{dust}} \sim 100 \text{ K}$). Note, however,

that the density is far below the critical densities of H_2O ($n(\text{H}_2)_{\text{crit}} > 10^8 \text{ cm}^{-3}$). It is thus impossible to determine the density because the excitation is dominated by the radiation field. Indeed there is currently no information for SDP.17b about other strong emission in H_2O lines with higher excitation, as in the case of Mrk 231, and it remains difficult to infer the detailed conditions of the gas from the detection of a single line. However, despite the uncertainty in the amplification factor, clearly the intrinsic H_2O line luminosity is at least comparable to that of Mrk 231 (Table 2). Similarly, the ratio $I(\text{H}_2\text{O}(2_{02-111}))/I(\text{CO}(8-7))$ found for SDP.17b (~ 0.5) is consistent with that found for Mrk 231 (~ 0.75), but is completely different from standard PDRs (Sect. 2). In addition, the ratio $L(\text{H}_2\text{O})/L_{\text{IR}}$ is three times higher in SDP.17b than in Mrk 231 (2.1×10^{-5} vs. 0.6×10^{-5}). All the evidence suggests that SDP.17b and Mrk 231 have similar properties. SDP.17b may thus display excitation conditions as special as and a chemistry as rich as those of Mrk 231 (see Sect. 2), suggesting the influence of a luminous AGN.

However, strictly speaking, without observations of other transitions, the excitation of H_2O in SDP.17b could be lower than in Mrk 231 and mostly limited to levels with energy ~ 100 K, as the 2_{02} level. As discussed by G-A10, the 2_{02-111} line does not provide unique constraints on the overall excitation conditions, because it can also have a large contribution from gas excited by cooler dust ($T_{\text{dust}} \sim 40$ K). This excitation might be achieved in less extreme conditions found in warm dense gas over a more extended region, through dense hot cores, and/or via shocks, and is not necessarily associated with excitation by a powerful AGN. But it remains unclear if the above conditions could boost the $\text{H}_2\text{O}/\text{CO}$ line intensity ratio to the values that are observed.

We conclude that SDP.17b is likely to be an analogue of Mrk 231. Most of the water lines detected by vdW10 in Mrk 231 could already be detected in SDP.17b or in other lensed sources detected by *Herschel* (including SDP.81 and HLSW-01) using the PdBI today or during ALMA's early science phase.

The results reported in this paper are an example of the studies that can be initiated when many bright, lensed high- z submillimeter galaxies become available (through *Herschel* or other facilities). Follow-up observations of these sources, especially with the increased sensitivities afforded by ALMA or NOEMA, will allow the undertaking of comprehensive studies of molecular lines in sources similar to SDP.17b and provide new insights into the physical conditions of the dense warm molecular gas of high- z SMGs and their AGN.

Acknowledgements. Based on observations carried out with the IRAM Plateau de Bure interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). The authors are grateful to the IRAM staff for their support. US participants in H-ATLAS acknowledge support from NASA through a contract from JPL. Italian participants in H-ATLAS acknowledge a financial contribution from the agreement ASI-INAF I/009/10/0. The SPIRE development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC (UK); and NASA (USA). We also thank the anonymous referee for very helpful suggestions.

References

Alexander, D. M., Smail, I., Bauer, F. E., et al. 2005, *Nature*, 434, 738
 Barvainis, R., Tacconi, L., Antonucci, R., et al. 1994, *Nature*, 371, 586
 Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
 Bradford, C. M., Aguirre, J. E., Aikin, R., et al. 2009, *ApJ*, 705, 112
 Casoli, F., Gerin, M., Encrenaz, P. J., & Combes, F. 1994, *A&A*, 287, 716
 Combes, F., & Wiklind, T. 1997, *ApJ*, 486, L79

Cox, P., Neri, R., Krips, M., Omont, A., et al. 2011, *ApJ*, submitted
 Danielson, A., Swinbank, A., Smail, I., et al., 2011, *MNRAS*, 410, 1687
 Downes, D., Neri, R., Wiklind, T., et al. 1999, *ApJ*, 513, L1
 Eales, S., Dunne, L., Clements, D., et al. 2010, *PASP*, 122, 499
 Fabian, A. C., Smail, I., Iwasawa, K., et al. 2000, *MNRAS*, 315, L8
 Fiolet, N., Omont, A., Polletta, M., et al. 2009, *A&A*, 508, 117
 Fischer, J., Sturm, E., González-Alfonso, E., et al. 2010, *A&A*, 518, L41
 Frayer, D. T., Harris, A. I., Baker, A. J., et al. 2011, *ApJ*, 726, L22
 González-Alfonso, E., Smith, H. A., Fischer, J., & Cernicharo, J. 2004, *ApJ*, 613, 247
 González-Alfonso, E., Smith, H. A., Ashby, M. L. N., et al. 2008, *ApJ*, 675, 303
 González-Alfonso, E., Fischer, J., Isaak, K., et al. 2010, *A&A*, 518, L43 [G-A10]
 Guélin, M., Salomé, P., Neri, R., et al. 2007, *A&A*, 462, L45
 Ikarashi, S., Kohno, K., Aguirre, J. E., et al. 2010 [arXiv:1009.1455]
 Impellizzeri, C. M. V., McKean, J. P., Castangia, P., et al. 2008, *Nature*, 456, 927
 Ivison, R. J., Magnelli, B., Ibar, E., et al. 2010, *A&A*, 518, L31
 Lestrade, J.-F., Combes, F., Salomé, P., et al. 2010, *A&A*, 522, L4
 Lupu, R. E., Scott, K. S., Aguirre, J. E., et al. 2010, *ApJ*, submitted [arXiv:1009.5983] [Lu10]
 McKean, J. P., Impellizzeri, C. M. V., Bauer, F. E., et al. 2011, *MNRAS*, 410, 2506
 Meijerink, R., & Spaans, M. 2005, *A&A*, 436, 397
 Negrello, M., Perrotta, F., González-Nuevo, J., et al. 2007, *MNRAS*, 377, 1557
 Negrello, M., Hopwood, R., De Zotti, G., et al. 2010, *Science*, 330, 800 [Ne10]
 Oliver, S. J., Wang, L., Smith, A. J., et al. 2010, *A&A*, 518, L21
 Papadopoulos, P. P. 2010, *ApJ*, 720, 226
 Riechers, D. A., Weiß, A., Walter, F., & Wagg, J. 2010, *ApJ*, 725, 1032
 Riechers, D. A., Walter, F., Carilli, C. L., et al. 2011a, *ApJ*, 726, 50
 Riechers, D. A., Cooray, A., Omont, A., et al. 2011b, *ApJ*, submitted
 Rowan-Robinson, M., Broadhurst, T., Oliver, S. J., et al. 1991, *Nature*, 351, 719
 Sajina, A., Yan, L., Lutz, D., et al. 2008, *ApJ*, 683, 659
 Scott, K., Lupu, R. E., Aguirre, J. E., et al. 2011, *ApJ*, in press
 Solomon, P. M., & Vanden Bout, P. 2005, *ARA&A*, 43, 677
 Solomon, P. M., Downes, D., Radford, S. J. E., & Barrett, J. W. 1997, *ApJ*, 478, 144
 Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175
 Swinbank, M., Smail, I., Longmore, S., et al. 2010, *Nature*, 464, 733
 Valtchanov, I., Virdee, J., Ivison, R., et al. 2011, *MNRAS*, in press
 van der Werf, P., Isaak, K. G., Meijerink, R., et al. 2010, *A&A*, 518, L42 [vdW10]
 Vieira, J. D., Crawford, T. M., Switzer, E. R., et al. 2010, *ApJ*, 719, 763
 Weiß, A., Walter, F., Scoville, N. Z., et al. 2005, *A&A*, 438, 533
 Weiß, A., Requena-Torres, M. A., Gristen, R., et al. 2010, *A&A*, 521, L1

¹ UPMC Univ Paris 06, UMR7095, Institut d'Astrophysique de Paris, 75014 Paris, France

e-mail: omont@iap.fr

² CNRS, UMR7095, Institut d'Astrophysique de Paris, 75014 Paris, France

³ Institut de Radioastronomie Millimétrique (IRAM), 300 rue de la Piscine, 38406 Saint-Martin-d'Hères, France

⁴ Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, US

⁵ Leiden Observatory, Leiden University, Post Office Box 9513, 2300 RA Leiden, The Netherlands

⁶ Department of Physics and Astronomy, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

⁷ Univ Paris-Sud and CNRS, Institut d'Astrophysique Spatiale, UMR8617, 91405 Orsay, France

⁸ Argelander Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

⁹ Astrophysics Group, Physics Department, Blackett Lab, Imperial College London, Prince Consort Road, London SW7, UK

¹⁰ Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

¹¹ Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE, UK

¹² Laboratoire Astrophysique, Instrumentation et Modélisation Paris Saclay, Commissariat à l'Énergie Atomique (CEA)/Direction des Sciences de la Matière (DSM) – CNRS – Université Paris Diderot, Institut de recherche sur les lois fondamentales de l'Univers

- (Irfu)/Service d'Astrophysique, CEA Saclay, Orme des Merisiers, 91191 Gif-sur-Yvette Cedex, France
- ¹³ INAF – Osservatorio Astronomico di Padova, Vicolo Osservatorio 5, 35122 Padova, Italy; and SISSA, via Bonomea 265, 34136 Trieste, Italy
- ¹⁴ School of Physics and Astronomy, Cardiff University, The Parade, Cardiff, CF24 3AA, UK
- ¹⁵ Instituto Nacional de Astrofísica, Óptica y Electrónica, Apartado Postal 51 y 216, 72000 Puebla, Mexico
- ¹⁶ UK Astronomy Technology Center, Royal Observatory Edinburgh, Edinburgh, EH9 3HJ, UK
- ¹⁷ School of Physics and Astronomy, University of Nottingham, University Park, Nottingham NG7 2RD, UK
- ¹⁸ GEPI, Observatoire de Paris, CNRS, Université Paris Diderot, 5 Place Jules Janssen, 92190 Meudon, France
- ¹⁹ SETI Institute, 515 North Whisman Avenue, Mountain View CA, 94043, USA
- ²⁰ Astrophysics Branch, NASA Ames Research Center, MS 245-6, Moffett Field, CA 94035, USA
- ²¹ Oxford Astrophysics, Denys Wilkinson Building, University of Oxford, Keble Road, Oxford, OX1 3RH, UK
- ²² Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
- ²³ National Radio Astronomy Observatory Post Office Box 2, Green Bank, WV 24944, USA
- ²⁴ INAF-IASF Milano, via E. Bassini 15, 20133, Italy
- ²⁵ Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK
- ²⁶ University of Padova, Department of Astronomy, Vicolo Osservatorio 3, 35122 Padova, Italy
- ²⁷ School of Astronomy, Institute for Research in Fundamental Sciences (IPM), PO Box 19395-5746, Tehran, Iran
- ²⁸ California Institute of Technology, Pasadena, CA 91125, USA
- ²⁹ Jet Propulsion Laboratory, Pasadena, CA 91109, USA
- ³⁰ University of Colorado, CASA 389-UCB, Boulder, CO 80303, USA
- ³¹ Sterrenkundig Observatorium, Universiteit Gent, Krijgslaan 281-S9, 9000 Gent, Belgium
- ³² Departamento de Astrofísica, Facultad de CC. Físicas, Universidad Complutense de Madrid, 28040 Madrid, Spain
- ³³ Department of Physics, McGill University, Ernest Rutherford Building, 3600 Rue University, Montréal, Québec, H3A 2T8, Canada
- ³⁴ Department of Physics and Astronomy, Rutgers, The State University of New Jersey, Piscataway, NJ 08854-8019, USA
- ³⁵ Centre for Astrophysics, Science & Technology Research Institute, University of Hertfordshire, Hatfield, Herts, AL10 9AB, UK.
- ³⁶ Department of Physics and Astronomy, The Open University, Milton Keynes, MK7 6AA, UK