

# Mapping of loci from *Solanum lycopersicoides* conferring resistance or susceptibility to *Botrytis cinerea* in tomato

Joel Davis · Daozhan Yu · Wendy Evans ·  
Tufan Gokirmak · Roger T. Chetelat · Henrik U. Stotz

Received: 18 November 2008 / Accepted: 8 April 2009  
© The Author(s) 2009. This article is published with open access at Springerlink.com

**Abstract** Cultivated tomato (*Solanum lycopersicum*, syn. *Lycopersicon esculentum*) is susceptible to the necrotrophic ascomycete and causal agent of gray mold, *Botrytis cinerea*. Resistance to this fungal pathogen is elevated in wild relatives of tomato, including *Solanum lycopersicoides*. An introgression line population (IL) containing chromosomal segments of *S. lycopersicoides* within the background of tomato cv. VF36 was used to screen the genome for foliar resistance and susceptibility to *B. cinerea*. Based on this screen, putative quantitative trait loci (QTL) were identified, five for resistance and two for susceptibility. Four resistance QTL decreased infection frequency while the fifth reduced lesion diameter. One susceptibility QTL increased infection frequency whereas the other increased lesion diameter. Overlapping chromosomal segments provided strong evidence for partial resistance on chromosomes 1

and 9 and for elevated susceptibility on chromosome 11. Segregation analysis confirmed the major resistance QTL on the long arm of chromosome 1 and susceptibility on chromosome 11. Linkage of partial resistance to chromosome 9 could not be confirmed. The usefulness of these data for resistance breeding and for map-based cloning of foliar resistance to *B. cinerea* is discussed.

## Introduction

*Botrytis cinerea* (teleomorph: *Botryotinia fuckeldiana* (de Bary) Whetzel) is a necrotrophic ascomycete that attacks more than 200 host species (Elad et al. 2004). This fungus infects various fruit, flower, and vegetable crops in the field and/or after harvest. The cultivated tomato, *Solanum lycopersicum* (formerly *Lycopersicon esculentum*), is an agronomically important host, but variation for resistance to *B. cinerea* among cultivars is small. Sources for resistance to *B. cinerea* have therefore been sought among wild relatives of tomato. *Solanum lycopersicoides* exhibits high resistance to *B. cinerea* (Rick 1987; Rick and Chetelat 1995, Chetelat et al. 1997). Importantly, accession LA2951, which was used to develop a population of introgression lines (ILs) (Canady et al. 2005), proved to express strong foliar resistance to *B. cinerea* (Guimaraes et al. 2004).

Other resistance traits are important for tomato production. Specifically for greenhouse production, stem resistance is a critical trait because pruning provides sites for fungal penetration. Hence, *Solanum habrochaites* (formerly *L. hirsutum*) and *Solanum neorickii* (formerly *L. parviflorum*) have been used to identify quantitative trait loci (QTL) for stem resistance to *B. cinerea* (Finkers et al. 2007, 2008). QTL mapping has also been done in *Arabidopsis thaliana* (Denby et al. 2004; Rowe and Kliebenstein 2008) and, although the genes

---

Communicated by I. Paran.

---

J. Davis · W. Evans  
Department of Horticulture, Oregon State University,  
Corvallis, OR 97331, USA

D. Yu  
School of Medicine,  
University of Maryland, Baltimore, MD 21201, USA

T. Gokirmak  
Plant Molecular and Cellular Biology Program,  
University of Florida, Gainesville, FL 32611, USA

R. T. Chetelat (✉)  
Department of Plant Sciences,  
University of California, Davis, CA 95616, USA  
e-mail: trchetelat@ucdavis.edu

H. U. Stotz  
RIKEN Plant Science Center, Yokohama,  
Kanagawa 230-0045, Japan

that are responsible for quantitative resistance to *B. cinerea* remain at large, camalexin has been shown to contribute to defense against this fungus (Kliebenstein et al. 2005; Rowe and Kliebenstein 2008). Approximately half of the resistance QTL recently identified were associated with camalexin accumulation (Rowe and Kliebenstein 2008). *Arabidopsis* studies provide valuable background information that may aid in the identification of candidate genes that control resistance QTL in crop species such as tomato. The relationship to induced phytoalexin biosynthesis is particular intriguing in light of rishitin's role in UV protection of fruits against *B. cinerea* (Charles et al. 2008).

The objectives of the present study were to locate resistance factors from *S. lycopersicoides* using a nearly complete set of ILs. Based on a genome-wide screen for foliar resistance to *B. cinerea*, three chromosomal regions were selected for further characterization in segregating progeny. The results presented herein establish the ILs as a potentially useful resource for *B. cinerea* resistance breeding in tomato.

## Materials and methods

### Biological material

Introgression lines (ILs) of an original backcross *Solanum lycopersicum* cv. 'VF36' × *Solanum lycopersicoides* 'LA2951' were used for linkage mapping of resistance to *B. cinerea*. The development and description of this population of ILs have been reported (Canady et al. 2005). Seeds of the IL's and parental lines as well as in vitro-propagated clones of the F<sub>1</sub> hybrid were provided by the Tomato Genetics Resource Center (TGRC), University of California, Davis. *B. cinerea* strain B05.10 was a gift of Jan van Kan (Wageningen University, The Netherlands).

### Growth conditions

Seeds were treated with bleach (Guimaraes et al. 2004) and sown in soil (Sunshine Mix SB40, SunGrow). Plants were grown in a greenhouse with supplemental lighting. Day-time and night-time temperatures were set at 21 and 16°C, respectively. Slow release fertilizer (Osmocote 15-9-12) was added to the soil after 1 month of growth. Seedlings were grown for a minimum of 6–8 weeks prior to leaf removal for inoculation trials. A completely randomized design was used to grow seedlings in 5 × 5 flats consisting of 0.5 l pots. Multiple flats were used per experiment. Consecutive numbers were given to the seedlings that were planted. It was, therefore possible to conduct "blind" assays because identification of genotypes was avoided.

*B. cinerea* was grown on potato dextrose agar from conidia stored at –80°C. The fungus was grown in

darkness for 3–4 days and then exposed to fluorescent light (34 μmol m<sup>-2</sup> s<sup>-1</sup>) to promote sporulation. Conidia were harvested according to published protocols (Benito et al. 1998; Guimaraes et al. 2004).

### Foliar inoculation

For the genome wide screen, a minimum of five and ten individuals per homozygous and heterozygous IL were screened for resistance to *B. cinerea*, respectively. The youngest, fully expanded leaves (typically the fourth leaf from the apex) were detached from the plant and inserted into moist florist foam to maintain turgidity. Prior to inoculation, conidia (10<sup>6</sup> ml<sup>-1</sup>) were incubated in Gamborg's medium containing 10 mM sucrose and 10 mM potassium phosphate, pH 6 for 2–3 h at room temperature. A total of ten droplets (10<sup>3</sup> conidia inoculum<sup>-1</sup>) were applied per leaf. Inoculated leaves were maintained in clear plastic boxes at room temperature under saturating humidity and a 12 h light/12 h dark regime at 34 μmol m<sup>-2</sup> s<sup>-1</sup>. Each box contained both *S. lycopersicum* 'VF36' and *S. lycopersicoides* 'LA2951' as controls. Lesions were evaluated 2–4 days after inoculation based on two parameters. First, the frequency of lesion expansion was based on the ability of the fungus to spread beyond the initial necrotic region that is confined to a diameter of 2 mm. This assessment provides an indicator of healthy versus diseased leaf tissues. Second, lesion diameters were measured using a caliper.

For the marker cosegregation analysis a minimum of 50 segregating progeny of the IL's were inoculated using the same inoculation procedure, but with two modifications. A total of ten droplets (5 × 10<sup>2</sup> conidia inoculum<sup>-1</sup>) were applied to each leaf, and the frequency and diameter of lesions expanding more than 3 mm were recorded.

### Statistical analysis

The SAS (Cary, NC, USA) program package was used to analyze all of the data. Data analysis of the frequency of expanding lesions was based on a binomial distribution using a generalized linear model (GENMOD). Average numbers of infections per leaf over 2–3 days period (3–4 days for the cosegregation test) were used as response variables per ten inoculations. IL was used as a covariate and the model was based on a logistic regression. The DSCALE option was used to limit overdispersion of the model.

The fixed effect model of an analysis of variance (ANOVA) was used to determine differences in lesion diameter among ILs. Levene's Test was used to assess the homogeneity of variances and only if the data passed this test, contrasts and least squares means (LSMEANS) were used to identify significant differences among means. Each date of evaluation was tested separately.

## DNA Extraction

For Southern blot and CAPS marker analysis, DNA extraction was performed according to the protocol of Miklas et al. (1993) with minor modifications. DNA samples were quantified using a UV fluorimeter (Hoefer).

## Marker analysis

Southern blots were prepared by digesting DNA from the ILs, VF36, LA2951, and F<sub>1</sub> with *EcoRI* followed by electrophoresis on 1% agarose gels. PCR primers derived from cloned and sequenced tomato RFLP's (Table 3) were used to amplify fragments from the VF36 parent. Amplification conditions were optimized using a gradient thermocycler (BioRad) to determine appropriate annealing temperatures (Table 3). The PCR profile consisted of a 3 min denaturing at 94°C followed by 35 cycles of 30 s at 94°C, 45 s annealing, and 2 min at 72°C, a final extension step at 72°C for 7 min, and then held at 4°C. Fragments were gel purified and labeled with the AlkPhos labeling kit (Amersham Biosciences). Hybridization and signal detection were performed using the manufacturer's instructions.

CAPS markers were generated by PCR amplification of the parents and F<sub>1</sub> using the same RFLP primers, followed by restriction digests of aliquots of the amplification products with enzymes, such as BamHI, BsrDI, DraIII, HaeIII, HindIII, NotI, and XhoI. After visualization on agarose gels, the restriction enzymes that allowed identification of the parental and F<sub>1</sub> genotypes were used for subsequent genotyping of the IL progeny (Table 3).

## Results

### Screening of the *S. lycopersicoides* genome for foliar resistance to *B. cinerea*

*S. lycopersicoides* ILs were screened for resistance to *B. cinerea* over a period of more than 2 years (Table 1). Seven trials were conducted to evaluate all of the ILs, which together represent more than 96% of the map units in the *S. lycopersicoides* genome (Canady et al. 2005). A total of 58 ILs were tested, 49 of those belonging to a 'primary set' of ILs (Canady et al. 2005). This allowed for complete coverage of the *S. lycopersicoides* genome with the exception of a region encompassing the marker TG65 on chromosome 4 (Fig. 1) and two additional small segments on chromosomes 2 and 3 (Canady et al. 2005). Given that *S. lycopersicoides* restricts the formation of expanding lesions (Guimaraes et al. 2004), all of the seven trials included data on frequencies of lesion expansion (Table 1). In addition, diameters of expanding lesions were recorded during the last four trials (Table 2).

**Table 1** Identification of *S. lycopersicoides* introgression lines with altered resistance to *B. cinerea*

Screen	Season Year	LA	IL <sup>a</sup>	<i>n</i>	RIF <sup>b</sup>	<sup>b, c</sup>	<i>P</i> value
1	Winter 03	4268	9A	6	0.53	6.81	0.0091
2	Summer 03	4233	1B	16	0.65	13.2	0.0003
2	Summer 03	4237	2	16	0.6	15.94	<0.0001
2	Summer 03	4269	9B	16	0.73	9.31	0.0023
3	Summer 03	4242	3	10	0.35	9.35	0.0022
3	Summer 03	4279	11C	10	1.62	8.75	0.0031
5	Summer 04	4264	8A	12	1.27	4.78	0.0299
6	Summer 04	3866	1A	7	2.19	7.45	0.0063
6	Summer 04	4234	1C	11	0.53	5.91	0.0151
6	Summer 04	4277	11A	17	1.52	8.17	0.0043
7	Summer 05	4233	1B	25	0.77	6.72	0.0095
7	Summer 05	4234	1C	20	0.77	5.84	0.0156
7	Summer 05	4268	9A	10	0.5	21.2	<0.0001


Statistically significant differences in frequency of infection are based on GENMOD

<sup>a</sup> Introgression lines are designated by chromosome number and a letter based on chromosomal location from top to bottom

<sup>b</sup> Relative Infection frequency in comparison to *S. lycopersicum* cv. VF36

<sup>c</sup>  $\chi^2$ -tests are based on contrasts between individual introgression lines and *S. lycopersicum* cv. VF36

Six *S. lycopersicoides* ILs were significantly more resistant to *B. cinerea* than *S. lycopersicum* 'VF36' when frequency of infection was used as a variable. Three of these lines were verified in two independent trials. These lines are referred to as IL-1B and IL-1C because they contain introgressed segments from chromosome 1, and as IL-9A with an introgressed segment of chromosome 9. Another line, IL-9B, was more resistant to *B. cinerea* than 'VF36' when 16 individuals were tested during the second trial (Table 1), but this line was not significantly different from the tomato parent when eight or five individuals were tested in 2003 and 2005, respectively. IL-9A and IL-9B share a polymorphism for the marker TG9 and, therefore, contain a region of overlap (Canady et al. 2005). Nevertheless, the phenotype of IL-9A was more robust and the magnitude of the phenotypic effect was larger in this line than in IL-9B. This suggests either that a susceptibility locus is located on the chromosomal segment that was introgressed into IL-9B or that more than one resistance locus is present on the region that was introgressed into IL-9A. The former explanation appears more likely because line IL-9C (LS10-6A) contains part of the chromosomal region of IL-9B and has a phenotype that is similar to tomato 'VF36' (Fig. 1). Moreover, IL-3, which shares a region containing TG18 with IL-9A, is resistant to *B. cinerea*. These data suggest that a quantitative trait locus (QTL) is located near the 'top' of chromosome 9. IL-2 was



**Fig. 1** Screen of the *S. lycopersicoides* genome for resistance to *B. cinerea*. Introgression lines of the **a** primary and **b** secondary set were screened. Columns indicate chromosomes with markers (not to scale), rows indicate introgression lines. *Lines in bold* were tested at least twice. *Italicized lines in bold* were only scored as hyper-resistant once out of two trials (IL-2) or twice out of three trials (IL-9B). *Solid and hatched boxes* indicate homozygous and heterozygous segments,

respectively. Reduction in frequency of infection and rate of lesion expansion are color-coded *red* and *yellow*, respectively. Elevation in frequency of infection and rate of lesion expansion are color-coded *blue* and *purple*, respectively. The consensus shown beneath all introgression lines shows the location of resistance and susceptibility QTL. In each of these indicated consensus regions differences in phenotypes outnumbered similarities to *S. lycopersicum* cv. VF36

evaluated as resistant to *B. cinerea* when 16 individuals were analyzed during the summer but not when seven individuals were analyzed during the winter of 2003. It is possible that there is a seasonal influence or resistance at this locus. We have observed that resistance of *S. lycopersico-*

*ides* can weaken during winter months under conditions in Oregon, but we have not analyzed this phenomenon further. Collectively, these data suggest that QTL controlling frequency of infection by *B. cinerea* are located on chromosomes 1, 2, and 9, and possibly on chromosome 3.

**Table 2** Identification of *S. lycopersicoides* introgression lines with altered resistance to *B. cinerea*

Screen	Season year	LA	IL <sup>a</sup>	<i>n</i>	LER <sup>b</sup>	<i>F</i> value <sup>c</sup>	<i>P</i> value	Transformation <sup>d</sup>
4	Winter 04	4246	4	6	0.57	4.94	0.0275	Log
4	Winter 04	4278	11B	10	1.45	5.33	0.0221	Log
4	Winter 04	4281	12	11	1.57	4.52	0.0349	Log
6	Summer 04	4249	5	8	1.33	4.39	0.0405	Inv
7	Summer 05	3889	8B	12	1.32	4.06	0.0467	Log

Statistically significant differences in rates of lesion expansion are based on ANOVA

<sup>a</sup> Introgression lines are designated by chromosome number and a letter based on chromosomal location from top to bottom

<sup>b</sup> Lesion expansion rates as compared to *S. lycopersicum* cv. VF36

<sup>c</sup> Contrasts were used to determine statistical differences between means of introgression lines and *S. lycopersicum* cv. VF36

<sup>d</sup> Logarithmic and inverse transformations were used to fulfill the homogeneity criterion of variances

Four *S. lycopersicoides* ILs were significantly more susceptible to *B. cinerea* than *S. lycopersicum* ‘VF36’ when frequency of infection was evaluated. IL-11A and IL-11C both contain introgressed segments located on chromosome 11 and share segments with IL-11B, an IL which is more susceptible to *B. cinerea* than the tomato parent based on lesion expansion (Table 2). A higher frequency of expanding lesions was also observed for lines IL-1A and IL-8A.

Besides IL-11B, three other lines, IL-5, IL-8B, and IL-12, were more susceptible than tomato ‘VF36’ when lesion expansion was evaluated (Table 2). Based on the consensus of all lines evaluated, a susceptibility QTL may be present on chromosome 5 (Fig. 1), but this inference is solely based on the phenotype of IL-5 because no other line contained the *S. lycopersicoides* polymorphism for CD64 (Canady et al. 2005).

IL-4 was scored as more resistant to *B. cinerea* than ‘VF36’ based on lesion expansion data. The consensus data of all the lines suggests a resistance QTL is linked to CT50 on chromosome 4. In hindsight, it would have been advisable to include LA4250 (Canady et al. 2005) for comparison with IL-4 to verify this prediction. However, line LA4250 contains additional introgressed segments from chromosomes 5, 8, and 9. LA4250 was not included in the original screen because the introgressed region was covered by IL-4.

Segregation analysis of a foliar *B. cinerea* resistance QTL on chromosome 1

Based on genome-wide analysis of resistance to *B. cinerea*, a QTL altering infection frequency was identified on chromosome 1 (Table 1). Fewer expanding lesions were observed when accession IL-1B was challenged with *B. cinerea* compared to tomato cv. ‘VF36’. In order to determine whether this difference in frequency of infection was linked to marker TG17 on chromosome 1, a segregation analysis was performed in the progeny of heterozygous IL-1B. TG17 is located at the bottom of an approximately 43 cM-long segment that has been introgressed into IL-1B (Canady et al. 2005). Segments introgressed from *S. lycopersicoides* recombine at only a very low rate with corresponding tomato chromosomes (Canady et al. 2006). Recombination rates are generally only 0–10% of expected frequencies (i.e. 90–100% suppressed). In the absence of significant recombination, single markers are sufficient to genotype each IL in segregating populations.

Segregation of TG17 was not significantly different from the Mendelian 1:2:1 expectation. Individuals carrying the *S. lycopersicoides* allele were significantly more resistant to *B. cinerea* both in terms of infection frequency and in terms of lesion diameter (Table 4). Resistance of the heterozy-

**Table 3** CAPS markers used for genotyping introgression lines (ILs)

Marker <sup>a</sup>	Chromosome/IL	Primer sequences (5’-3’) <sup>1</sup>	Size <sup>b</sup> (bp)	Anneal. temp. (°C)	Enzyme
TG17	1B	cggctgtgtactatctgga aaaatcaattgaaccggctgt	2,200	55	<i>Hind</i> III
TG18	9A	ctcaagctccagctgtttcc gctccttctgcaatgggtaa	1,700	58	<i>Bam</i> HI
TG46	11C	atccaacctctgagcacac gttcttggaaaccgatattgc	1,400	62	<i>Hae</i> III

<sup>a</sup> Marker and primer sequences were obtained from Solanaceae Genomics Network database (<http://www.sgn.cornell.edu>)

<sup>b</sup> PCR product size before digestion with restriction enzyme

gous progeny to *B. cinerea* appeared intermediate relative to homozygous segregants. Nevertheless, heterozygotes were significantly more resistant than homozygotes carrying the *S. lycopersicum* allele (Table 4). Conversely, resistance of heterozygotes to *B. cinerea* was not significantly different from homozygotes carrying the *S. lycopersicoides* allele. With respect to lesion diameter, no significant differences in resistance were observed between heterozygous and homozygous progenies, indicating that genetic control of this trait may be different from infection frequency.

Accession IL-1B apparently harbors a major QTL linked to marker TG17. Infection frequency of homozygotes for the *S. lycopersicoides* allele was reduced by 55% relative to the difference between tomato cv. VF36 and *S. lycopersicoides* LA2951 (Fig. 2a). This approximation compared favorably with estimates that were obtained during the genome-wide screen for *B. cinerea* resistance (Table 1). In the latter case, infection frequency was reduced by 45 and 23% during 2003 and 2005 seasons. More importantly, segregation analysis demonstrates that the QTL on chromosome 1 explains approximately 21% of the phenotypic variation (Table 4). By comparison, the effect on lesion size was smaller (11% of the phenotypic variation) and not observed during the genome-wide screen, perhaps a result of the smaller sample size or lower expression of the trait during the 2005 season.

Resistance to *B. cinerea* is not linked to TG18 in IL-9A

Genome wide analysis of resistance to *B. cinerea* indicated a QTL decreasing infection frequency on chromosome 9. To determine linkage between marker TG18 and this trait, homozygous IL-9A and cv. VF36 were crossed, then self-pollinated to generate a segregating population. TG18 is

located 14 cM from the top of chromosome 9. The segment that has been introduced into IL-9A is approximately 18 cM in size (Canady et al. 2005).

Segregation of TG18 in the F<sub>2</sub> progeny of IL-9A differed significantly from the 1:2:1 expectation. Marker segregation was skewed in favor of the allele from cultivated tomato. Infection frequency (Fig. 3a) and lesion diameter (Fig. 3b) did not significantly differ among the three genotypes that were defined by TG18 segregation. Nevertheless, all the three IL-9A genotypes were significantly more resistant to *B. cinerea* than cv. VF36. Both infection frequency and lesion diameter were reduced (Fig. 3). The magnitude of reduction in an infection frequency was similar to the 50 and 53% reduction observed during the genome-wide screen (Table 1). The progeny was not genotyped based on phenotype, but on molecular marker data. In the absence of discernible phenotypic segregation, the most parsimonious explanation is that this partial resistance locus is not linked to the segment on chromosome 9 but located elsewhere in the *S. lycopersicoides* genome even though markers used by Canady et al. (2005) did not detect the presence of other introgressed segments in IL-9A.

Analysis of a putative susceptibility locus linked to TG46 on chromosome 11

Genome wide analysis of the interactions between *S. lycopersicoides*-derived ILs and *B. cinerea* indicated a susceptibility locus on chromosome 11. Marker TG46 was used to analyze segregation for susceptibility to *B. cinerea* in progeny of heterozygous line IL-11C. TG46 is located approximately in the middle of a ca. 50 cM segment introgressed into IL-11C. Surprisingly, seedlings homozygous for the *S. lycopersicoides* allele were not recovered between

**Table 4** Segregation analysis of introgression line F<sub>2</sub> progeny

IL	Genotype <sup>a</sup>	Infection freq. <sup>b</sup>	<i>P</i> value <sup>c</sup>	% var. <sup>d</sup>	Lesion diameter (mm) <sup>e</sup>	<i>P</i> value	% var.
1B	+/+	6.9 (17)	–	21.4	6.2 (17)	–	11.0
	+/S	4.9 (30)	0.0226		5.3 (26)	0.1003	
	S/S	2.8 (8)	0.0009		4.4 (6)	0.0318	
9A	+/+	4.9 (27)	–	4.0	6.6 (24)	–	6.0
	+/S	5.5 (17)	0.1952		6.8 (17)	0.6163	
	S/S	3.8 (10)	0.3458		5.9 (8)	0.2039	
11C	+/+	4.7 (21)	–	3.8	5.2 (17)	–	6.6
	+/S	5.8 (28)	0.1977		6.4 (27)	0.0580	


<sup>a</sup> +/+ is homozygous for cv. VF36 allele, S/S is homozygous for *S. lycopersicoides* allele, and +/S is heterozygous

<sup>b</sup> Infection frequency is the mean number of expanding lesions per leaf (10 inoculation droplets per leaf); values in parentheses are the numbers of F<sub>2</sub> progeny of each genotype included in the statistical analysis

<sup>c</sup> *P* values compare the +/S or S/S genotypes against the +/+ genotype

<sup>d</sup> Percentage of phenotypic variation explained by each QTL in each F<sub>2</sub> population


<sup>e</sup> Lesion diameter is the mean diameter in mm of leaf lesions; values in parentheses are the numbers of F<sub>2</sub> progeny of each genotype included in the statistical analysis



**Fig. 2** Segregation for resistance to *B. cinerea* on chromosome 1. **a** Numbers of expanding lesions and **b** lesion diameters were scored among progeny of heterozygous IL-1B and compared to the parents LA2951 and VF36. Alleles of *S. lycopersicum* cv. ‘VF36’ and *S. lycopersicoides* line LA2951 are abbreviated + and S, respectively. Heterozygous +/S and homozygous +/+ and S/S individuals were identified using the CAPS marker TG17/*Hind*III. Means, standard errors, and sample sizes are indicated. Significant differences to +/+ genotypes are indicated at  $P < 0.05$  (\*) and  $P < 0.001$  (\*\*\*)

germinating and surviving individuals. Segregation of TG46 was not significantly different from an expectation of 1:2:0. Although we do not know whether embryo lethality occurs in this genotype because we did not attempt embryo rescue, homozygosity at this locus still causes a selective disadvantage.

The remaining genotypes were analyzed for susceptibility to *B. cinerea* (Fig. 4). Frequency of infection was not significantly different between genotypes that were heterozygous or homozygous for the *S. lycopersicum* allele. Lesion diameters were elevated in the heterozygous genotype compared to individuals homozygous for the *S. lycopersicum* allele and cv. VF36, but this difference was not statistically significant. It might be argued that a one-tailed test could be used based on the expectation of the genome-wide screen. In this case, the heterozygous genotype would be scored as susceptible. When compared to cv. VF36, susceptibility was clearly elevated (Fig. 4), probably a reflection of the larger number of cultivar seedlings tested. Lesion size was 25% larger in the case of IL-11C when compared in the heterozygous state to cv. VF36. Collectively, these data let us conclude that susceptibility locus exists on chromosome 11.




**Fig. 3** Segregation analysis for resistance to *B. cinerea* on chromosome 9. **a** Numbers of expanding lesions and **b** log N-transformed lesion diameters were scored among progeny of heterozygous IL-9A and compared to the parents LA2951 and VF36. +/+ is homozygous for the VF36 allele, S/S is homozygous for the LA2951 allele, and +/S is heterozygous based on CAPS marker TG18/*Bam*HI. Means, standard errors, and sample sizes are indicated

## Discussion

Analysis of foliar susceptibility to *B. cinerea* among 58 ILs derived from *S. lycopersicoides* has resulted in identification of four putative resistance QTL for infection frequency and one putative resistance QTL for lesion diameter. In addition, putative susceptibility QTL for infection frequency or lesion diameter was identified. Several of these loci are referred to *Fbc* for foliar *B. cinerea* resistance or susceptibility in Fig. 5.

### Colocation of foliar *B. cinerea* resistance and susceptibility QTL with known resistance loci

Several of the putative foliar *B. cinerea* resistance QTL identified herein are linked to previously recognized QTL for stem resistance to *B. cinerea*. The closest association was observed between *Fbc3* (Fig. 5) and *Snbc3*, the latter from *S. habrochaites* (Finkers et al. 2008). Although we did not find other regions of overlap, *Fbc2* and *Fbc4* are closely linked to *Rbcq2* and *Rbcq4b*, respectively (Finkers et al.




**Fig. 4** Segregation for susceptibility to *B. cinerea* on chromosome 11. Genotypes homozygous for the LA2951 allele were not recovered. **a** Numbers of expanding lesions and **b** lesion diameters were scored among the remaining progeny of heterozygous IL-11C and compared to the parents LA2951 and VF36. +/+ is homozygous for the VF36 allele, and +/S is heterozygous based on the CAPS marker TG46/*Hael*-II. Means, standard errors, and sample sizes are indicated. Significant difference to VF36 is indicated at  $P < 0.01$  (\*\*)

2007). Homologous QTL for stem resistance to *B. cinerea* were previously identified (Finkers et al. 2008), suggesting that a limited number of QTL specify resistance to *B. cinerea* in different *Solanum* species. Collectively, data on stem resistance (Finkers et al. 2008; Finkers et al. 2007) and on foliar resistance from *S. lycopersicoides* presented here indicate genomic specialization for *B. cinerea* resistance on chromosomes 1, 2, 3, 4, and 9 of *Solanum*. Nevertheless, comparative analysis (Fig. 5) also demonstrates that stem and foliar resistance are controlled by separate QTL. The map locations of resistance genes and QTL controlling interactions with other pathogens are shown in Fig. 5, but due to the large sizes of introgressed *S. lycopersicoides* fragments, any inferences about potential coincidences would be speculative.

#### Confirmed QTL for foliar resistance or susceptibility to *B. cinerea*

Three putative QTL were characterized further because overlapping segments of ILs strongly suggested the presence of loci that have a quantitative influence on interactions



**Fig. 5** Synopsis of resistance and susceptibility loci mapped to chromosomes 1–5 and to chromosome 11. The genetic map has previously been published (Canady et al. 2005). Solid vertical bars delimit locations of putative foliar *B. cinerea* (*Fbc*) resistance loci on chromosomes 1 through 4 or susceptibility QTL on chromosomes 5 and 11. Approximate locations of previously reported qualitative resistance genes (*R* genes) or resistance QTL (dashed vertical lines or underlined) are also shown. Resistance loci on chromosomes 3 and 4 have previously been designated (Finkers et al. 2008); QTL3 and 4 were renamed to *Snbc3* and 4 in reference to *Solanum neorickii*. Following references assign remaining resistance loci: *Cf9* (Parniske et al. 1997), *Rbcq1* and 2 (Finkers et al. 2007), *Cml-1* (Sandbrink et al. 1995), *rx-2* (Yu et al. 1995), *Tm-1* (Young and Tanksley 1989), *Bs4* (Ballvora et al. 2001), *Pto* and *Prf* (Salmeron et al. 1996), *lb5b* and *lb11b* (Brouwer and St Clair 2004), *Sm* (Behare et al. 1991), and *I2* (Ori et al. 1994)

with *B. cinerea* in leaves. *Fbc1*, a major QTL for foliar resistance to *B. cinerea*, was identified in this fashion. This QTL was initially identified based on its reduction of infection frequency (Table 1). *Fbc1* explains approximately 21% of the phenotypic variation in susceptibility to *B. cinerea*, which is larger than the magnitudes of previously reported stem resistance QTL (Finkers et al. 2007, 2008). Analysis of a segregating IL-1B population also revealed a significant difference in lesion diameter between homozygous progenies, but this effect was much smaller than the infection frequency (Table 4). The relatively small effect on lesion size probably explains why this trait was not detected in the previous genome-wide screen (Tables 1, 2). Based on



the large effect on infection frequency, *Fbc1* would make a good candidate for fine-scale mapping and eventual cloning of the resistance gene. Although suppression of recombination in ILs represents a genetic barrier (Canady et al. 2005), *S. pennellii* has successfully been used as a bridge species to overcome this barrier and increase the rate of recombination between *S. lycopersicoides* and cultivated tomato (Canady et al. 2006).

A second resistance QTL was examined in detail based on the observation that IL-9A and IL-9B, with overlapping fragments of chromosome 9, were more resistant to *B. cinerea*. Surprisingly, segregation of resistance was not observed using marker TG18 in progeny of heterozygous IL-9A, although all of the progeny were more resistant than tomato cv. VF36 (Fig. 3). This suggests that the resistance locus either exists somewhere else in the genome or that linkage between TG18 and the QTL has been broken. With respect to the former possibility, it is interesting to note that markers used for mapping ILs were separated by 18.4 cM on average with a range of 2.5–64 cM (Canady et al. 2005). Although Canady et al. (2005) reported introgression of a single segment on chromosome 9 in IL-9A, it can therefore not be excluded that an unlinked small piece was introduced from the *S. lycopersicoides* genome. This could explain our inability to detect segregation for partial resistance to *B. cinerea* using TG18.

Genome wide screening for *B. cinerea* susceptibility indicated that the chromosomal region of *S. lycopersicoides* in IL-11C increases infection frequency (Table 1). However, segregation analysis showed that *Lbc11* significantly increased lesion diameter. Lesion size was not evaluated during the primary screen. An effect on lesion diameter may, therefore, have been missed during this earlier assessment. The population size of the segregating progeny was larger than the number of individuals used for the primary screen. Quantitative differences in lesion diameter are therefore assumed to be reliable. Variability between disease incidence and lesion growth was also observed for QTL9 from *S. neorickii* (Finkers et al. 2008). Although the authors argued that in their case QTL9 was expected to reduce disease incidence, the cause for different experimental observations in separate screens was not resolved. The size of the phenotypic effect appeared to be relatively small in the case of QTL9 (Finkers et al. 2008). Similarly, *Fbc11* appeared to be a relatively minor QTL, at least in comparison to *Fbc1*. Thus, in the case of minor QTL, large samples appear to be needed to reliably assign resistance or susceptibility traits. However, in the case of *Fbc11* no surviving homozygotes for the *S. lycopersicoides* were recovered, possibly indicating a major defect-related to plant development and defense.

## Conclusion

Our studies confirm that *S. lycopersicoides* is a potent source of foliar resistance to *B. cinerea*. Several partially resistant ILs were identified in this study. These ILs could be used in combination with marker-assisted selection to develop improved tomato cultivars.

**Acknowledgments** We thank Rejane Guimarães for advice in resistance screening. The California Tomato Commission provided financial support.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Ballvora A, Pierre M, van den Ackerveken G, Schornack S, Rossier O, Ganal M, Lahaye T, Bonas U (2001) Genetic mapping and functional analysis of the tomato *Bs4* locus governing recognition of the *Xanthomonas campestris* pv. *vesicatoria* AvrBs4 protein. Mol Plant Microbe Interact 14:629–638
- Behare J, Laterrot H, Sarfatti M, Zamir D (1991) Restriction-fragment-length-polymorphism mapping of the *Stemphylium* resistance gene in tomato. Mol Plant Microbe Interact 4:489–492
- Benito E, ten Have A, van't Klooster J, van Kan J (1998) Fungal and plant gene expression during synchronized infection of tomato leaves by *Botrytis cinerea*. Eur J Plant Pathol 104:207–220
- Brouwer DJ, St Clair DA (2004) Fine mapping of three quantitative trait loci for late blight resistance in tomato using near isogenic lines (NILs) and sub-NILs. Theor Appl Genet 108:628–638
- Canady MA, Meglic V, Chetelat RT (2005) A library of *Solanum lycopersicoides* introgression lines in cultivated tomato. Genome 48:685–697
- Canady MA, Ji Y, Chetelat RT (2006) Homeologous recombination in *Solanum lycopersicoides* introgression lines of cultivated tomato. Genetics 174:1775–1788
- Charles MT, Mercier J, Makhlof J, Arul J (2008) Physiological basis of UV-C-induced resistance to *Botrytis cinerea* in tomato fruit—I. Role of pre- and post-challenge accumulation of the phytoalexin-rishitin. Postharvest Biol Technol 47:10–20
- Chetelat RT, Cisneros P, Stamoia L, Rick CM (1997) A male-fertile *Lycopersicon esculentum* × *Solanum lycopersicoides* hybrid enables direct backcrossing to tomato at the diploid level. Euphytica 95:99–108
- Denby KJ, Kumar P, Kliebenstein DJ (2004) Identification of *Botrytis cinerea* susceptibility loci in *Arabidopsis thaliana*. Plant J 38:473–486
- Elad Y, Williamson B, Tudzynski P, Delen N (2004) *Botrytis*: biology, pathology, and control. Kluwer, Dordrecht
- Finkers R, van den Berg P, van Berloo R, ten Have A, van Heusden AW, van Kan JAL, Lindhout P (2007) Three QTLs for *Botrytis cinerea* resistance in tomato. Theor Appl Genet 114:585–593
- Finkers R, Bai YL, van den Berg P, van Berloo R, Meijer-Dekens F, ten Have A, van Kan J, Lindhout P, van Heusden AW (2008) Quantitative resistance to *Botrytis cinerea* from *Solanum neorickii*. Euphytica 159:83–92
- Guimaraes RL, Chetelat RT, Stotz HU (2004) Resistance to *Botrytis cinerea* in *Solanum lycopersicoides* is dominant in hybrids with

- tomato, and involves induced hyphal death. *Eur J Plant Pathol* 110:13–23
- Kliebenstein DJ, Rowe HC, Denby KJ (2005) Secondary metabolites influence *Arabidopsis/Botrytis* interactions: variation in host production and pathogen sensitivity. *Plant J* 44:25–36
- Miklas PN, Delorme R, Riley R (1993) Identification of QTL conditioning resistance to white mold in snap bean. *J Am Soc Hortic Sci* 128:564–570
- Ori N, Paran I, Aviv D, Eshed Y, Tanksley S, Zamir D, Fluhr R (1994) A genomic search for the gene conferring resistance to *Fusarium* wilt in tomato. *Euphytica* 79:201–204
- Parniske M, Hammond-Kosack KE, Goldstein C, Thomas CM, Jones DA (1997) Novel disease resistance specificities result from sequence exchange between tandemly repeated genes at the *Cf-4/9* locus of tomato. *Cell* 91:821–832
- Rick C (1987) Genetic resources in *Lycopersicon*. In: Nevins DJ, Jones RA (eds) *Plant biology*. Alan R. Liss, New York, pp 17–26
- Rick C, Chetelat R (1995) Utilization of related wild species for tomato improvement. *Acta Hort* 412:21–38
- Rowe HC, Kliebenstein DJ (2008) Complex genetics control natural variation in *Arabidopsis thaliana* resistance to *Botrytis cinerea*. *Genetics* 180:2237–2250
- Salmeron JM, Oldroyd GED, Rommens CMT, Scofield SR, Kim H-S, Lavelle DT, Dahlbeck D, Staskawicz BJ (1996) Tomato *Prf* is a member of the leucine-rich repeat class of plant disease resistance genes and lies embedded within the *Pto* kinase gene cluster. *Cell* 86:123–133
- Sandbrink JM, van Ooijen JW, Purimahua CC, Vrieling M, Verkerk R, Zabel P, Lindhout P (1995) Localization of genes for bacterial canker resistance in *Lycopersicon peruvianum* using RFLPs. *Theor Appl Genet* 90:444–450
- Young ND, Tanksley SD (1989) RFLP analysis of the size of chromosomal segments retained around the *Tm-2* locus of tomato during backcross breeding. *Theor Appl Genet* 77:353–359
- Yu ZH, Wang JF, Stall RE, Vallejos CE (1995) Genomic localization of tomato genes that control a hypersensitive reaction to *Xanthomonas campestris* pv. *vesicatoria* (Doidge) dye. *Genetics* 141:675–682