

Discovery of shocked H₂ around OH 231.8+4.2

K.P. Forde^{1*}, T.M. Gledhill¹

¹Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK

Accepted 2011 Month ??; Received 2011 Month ??; in original form 2011 Month ??

ABSTRACT

We present *K*-band integral field observations of the circumstellar envelope of the evolved star OH 231.8+4.2. Spatial and spectral information were simultaneously acquired using the SINFONI integral field unit, with adaptive optics, on the Very Large Telescope. The observations reveal the discovery of H₂ emission (1) around the centre of the nebula and (2) located in clumps along the Western side of the Northern lobe, presumably associated with the strong shocks that stimulate the previously reported H α emission at the same location. An observed H₂ 1 \rightarrow 0/2 \rightarrow 1S(1) line ratio of 8.3 ± 1.9 was calculated for the central field, a value consistent with shock excitation.

Key words: molecular data – circumstellar matter – stars: AGB and post-AGB – stars: individual(OH 231.8+4.2) – shock waves

1 INTRODUCTION

OH 231.8+4.2 (hereafter OH231) is an O-rich late spectral type (M) central star (Mira variable, QX Pup) with bipolar high-velocity dust and gas outflows (Alcolea et al. 2001), filamentary structures observed in scattered and molecular line emission, and large angular size ($10''\times 60''$). Often labeled as a post-AGB object or pre-planetary nebula, the presence of both a Mira central star and a main-sequence companion of spectral type A (Sánchez Contreras et al. 2004) seems to contradict this classification. OH231 is more likely a D-type bipolar symbiotic system (Frew & Parker 2010). However in some cases, morphological similarities do exist between post-AGB and symbiotic objects, most strikingly the presence of highly collimated bipolar nebulae. It is via fast-collimated outflows that these stars shape their surrounding nebula. Understanding the development and origin of these fast outflows is critical for advancing hydrodynamical models of wind interaction. Recent work by Lee et al. (2009) attempting to reproduce the high velocity molecular emission in AFGL 618 using collimated fast wind models, emphasises the need for further observations and model development in this area.

OH231 has been the subject of many studies spanning multiple wavelength ranges, for example: Cohen et al. (1985) were first to propose the existence of a binary companion; Bujarrabal et al. (2002) imaged the shape of the shocks using H α (reproduced in Fig. 1 [a]) detected with the *Hubble Space Telescope* (*HST*); Meakin et al. (2003) report *HST*/NICMOS NIR images of the dust distribution and hence a high resolution map of the extinction through the

nebula. Matsuura et al. (2006) using the MIDI and NACO instruments on the Very Large Telescope (VLT) detected a compact circumstellar disc. The envelope of OH231 is also known to be rich in molecular species (e.g. H₂O, OH, and SiO) however previous studies in the NIR have all returned null detections of H₂ (e.g. Weintraub et al. 1998; Matsuura et al. 2006).

In this Letter, we present the results of preliminary observations of OH231 at NIR (*K*-band) wavelengths showing for the first time the presence of shock-excited H₂. Throughout this work we assume OH231 is a member of the open cluster M46 at a distance of 1.3 kpc (Jura & Morris 1985). The origin of the coordinate system used in Figures 1,3, and 4 is given by the location of the SiO maser emission at RA=07^h42^m16^s.93, Dec=-14°42'50".2 (J2000) (Sánchez Contreras et al. 2002), and the inclination angle of the bipolar axis is 36° to the plane of the sky (Kastner et al. 1992).

2 OBSERVATIONS AND DATA REDUCTION

The data were taken using the SINFONI (Eisenhauer et al. 2003) instrument located on UT4 at VLT at Paranal, Chile, on the 1st/2nd Feb 2010. We use the lowest resolution mode (LRM), corresponding to the widest field-of-view ($8''\times 8''$) with adaptive optics (AO) and a natural guide star (NGS). A plate scale of 250×125 mas pixel⁻¹, and a spectral and velocity resolution of 4580 and 66 km s⁻¹, respectively, are available at this resolution (for a dispersion of 2.45 Å/pix and line FWHM of 1.96 pixels). All observations utilised the *K*-band (2.2 μ m) filter which covers many ro-vibrational H₂ emission lines. The ambient seeing varied from $\sim 0.6''$

* E-mail: k.p.forde@herts.ac.uk

Table 1. Details of the VLT/SINFONI observations for OH231 and the telluric standard stars taken on 1st/ 2nd February 2010.

	OH231			HD		
	South	Central	North	75004	63487	
Tot. Exp. (secs)	600	600	480	800	1	10
K (mag)	–	–	–	–	7.268 [†]	7.674 [†]
Night	1	1	2	1	1	2

[†] 2mass magnitude from: SIMBAD

to 1.1'' during the observations. The OH231 observations consisted of three fields focused on (1) the edge of the Northern lobe, (2) the central region, and (3) the middle of the Southern lobe (labelled N, C, S in Fig. 1 [b]). No H₂ was detected in the Southern field and will not be discussed further. Table 1 summarises exposures for each of the three fields. Telluric standard stars used for calibration are HD 75004 (G0V), and HD 63487 (G2V) for night one and two, respectively. An average AO-corrected PSF of ~ 340 mas FWHM is estimated from the standard stars. The data were reduced using the ESO common pipeline library to a wavelength-calibrated datacube and further analysed using both PyRAF¹. Wavelength calibration errors were corrected by comparison of OH emission lines with a high resolution template. Quoted velocities were adjusted to local standard of rest (LSR) velocities using 22.88 and 23.30 km s⁻¹ corrections for night one and night two, respectively. Line maps (Fig. 1 [c,d]) were generated by fitting the H₂ emission lines with a Gaussian profile; initial fit parameters (FWHM₀, central wavelength₀, etc.) were determined by manually fitting an individual H₂ line, only lines with a FWHM \approx FWHM₀ make up the final line maps. Signal-to-Noise (S/N) of the line maps was enhanced by smoothing the data with a 2 \times 2 pixel boxcar filter. Line rest wavelength information is from Black & van Dishoeck (1987).

3 RESULTS

We report the detection of several H₂ emission lines arising from both the centre and Northern lobe of OH231. In Figure 1 (c,d) we present continuum-subtracted line maps of the 1 \rightarrow 0S(1) transition for both fields, showing clearly H₂ emission arising from the centre of OH231 (Fig. 1 [d]) and from knots of material in the Northern lobe (Fig. 1 [c]). In Fig. 1 (a) the contours of the detected H₂ are shown in relation to H α emission, while Fig. 1 (b) gives the location of the H₂ relative to the strong continuum emission as imaged with a 2.12 μ m filter. An integrated spectrum of the Western region of the centrally located H₂ is shown in Figure 2, from which we note, a) several S- and Q-branch ro-vibrational H₂ lines, and b) a CO bandhead absorption feature (~ 2.3 μ m).

Channel maps, extracted from the SINFONI datacube, are presented in Figures 3 and 4 for the central and Northern regions respectively, showing how the distribution of the H₂ changes across the line profile.

Table 2 presents the flux measurements for the central and Northern fields, for lines with flux errors less than ~ 50

Table 2. Line fluxes, F , (in units of 10^{-19} W m⁻²) of the observed H₂ lines for the central and Northern fields. Measured peak line wavelengths (in μ m) are given for the central field.

Line	Central		Northern
	λ_{meas}	F	F
1 \rightarrow 0S(3)	1.9579	67.2 \pm 14.2	11.5 \pm 1.10
1 \rightarrow 0S(2)	2.0342	9.71 \pm 3.36 $^\circ$	–
1 \rightarrow 0S(1)	2.1222	91.5 \pm 6.16	9.69 \pm 0.76
2 \rightarrow 1S(1)	2.2481	7.73 \pm 3.38 [‡]	–
1 \rightarrow 0Q(1)	2.4071	46.4 \pm 14.5	11.9 \pm 2.95
1 \rightarrow 0Q(3)	2.4241	42.2 \pm 24.0	13.3 \pm 4.42

[‡] Flux measured for Western H₂ region only.

[◊] Measurement confined to elliptical region (Fig. 1 [d]).

per cent. Accurate 2 \rightarrow 1S(3), 1 \rightarrow 0S(0), and 1 \rightarrow 0Q(2) line flux measurements are not possible due to the presence of strong sky subtraction residuals at these wavelengths. We confine the calculation of the 1 \rightarrow 0/2 \rightarrow 1S(1) ratio for the central field to the area marked by the ellipse in Fig. 1 (d), where the 2 \rightarrow 1S(1) flux error is smallest. In this region we calculate a 1 \rightarrow 0/2 \rightarrow 1S(1) ratio of 8.3 ± 1.9 prior to extinction correction. We do not detect any 2 \rightarrow 1S(1) flux in the Eastern H₂ region, instead we estimate an upper 2 \rightarrow 1S(1) limit (3σ) for the flux in this region of 1.3×10^{-19} W m⁻², which in turn places a lower limit of ~ 9.5 on the 1 \rightarrow 0/2 \rightarrow 1S(1) ratio for this region.

4 ANALYSIS AND DISCUSSION

The 1 \rightarrow 0/2 \rightarrow 1S(1) ratio is a useful discriminator of excitation mechanisms. Pure fluorescence will yield a value of ≈ 2 while a value ≈ 10 indicates the excitation of the gas is being driven by shocks. However it is worth noting, these values depend on shock velocity and pre-shock gas density, with shocked H₂ capable of producing values as low as ≈ 4 (Smith 1995), while fluorescence can produce values approaching those of shocks (Hollenbach & Natta 1995). The 1 \rightarrow 0/2 \rightarrow 1S(1) ratio values, given above, suggest that shocks might be the main excitation mechanism, which agrees with the detection of shock-excited HCO⁺ in the centre of OH231 noted in the position-velocity diagrams of Sánchez Contreras et al. (2000). In order to determine the intrinsic 1 \rightarrow 0/2 \rightarrow 1S(1) ratio, it is necessary to remove the effects of extinction. In the K -band, it is sometimes possible to derive the level of extinction via the comparison of S-/Q-branch H₂ emission lines (see Smith et al. 2003). Unfortunately, due to poor atmospheric transmission above 2.4 μ m, it was not possible to derive a sensible estimate for extinction using the 1 \rightarrow 0 S(1) and Q(3) lines.

Extinction values of $A_K=3-4$ (mag) for the central region are estimated from the ($K - L'$) colour map of OH231 (Kastner et al. 1998). Although, most likely an overestimate, adjusting the Western 1 \rightarrow 0/2 \rightarrow 1S(1) ratio for these levels of extinction yields, for example, a value of 11.7 ± 2.6 ($A_K=4$). It is clear that any adjustment for extinction will increase the observed 1 \rightarrow 0/2 \rightarrow 1S(1) ratio, pushing it further towards the shock regime.

Using a typical value of 5.3×10^{-22} mag cm² for the extinction per unit column density of hydrogen, $A_V/N(\text{H})$, one can estimate the hydrogen column density implied by an $A_K=4$; yielding a $N(\text{H})$ for the H₂ regions of $\sim 7.2 \times 10^{22}$

¹ PyRAF is a product of the Space Telescope Science Institute, which is operated by AURA for NASA.




Figure 1. Narrow-band images and line maps for OH231; a star symbol indicates the SiO maser position (see text), inset labels display the telescope/instrument information. (a) Continuum-subtracted $H\alpha$ image with the $1\rightarrow 0S(1)$ H_2 contours overlaid in white, (b) $2.12\ \mu\text{m}$ image, showing mainly scattered light, with H_2 contours overlaid, SINFONI North, Central, and South (labeled N, C, S) fields are marked by green squares, inset compass shows orientation, (c) $1\rightarrow 0S(1)$ H_2 line map showing the extent of the emission located along the edge of the $H\alpha$ emission, two knots (A, B) of H_2 emission are indicated by arrows; superposed contours indicate the location of the $H\alpha$ emission, (d) $1\rightarrow 0S(1)$ H_2 line map for the central region showing the position/extent of the region used for $1\rightarrow 0/2\rightarrow 1S(1)$ calculation (ellipse). Colour bar units are in $\text{W m}^{-2}\ \text{px}^{-1}$. The $H\alpha$ data, from the Hubble Legacy Archive (see Acknowledgments), has been continuum-subtracted using the procedure outlined in Bujarrabal et al. (2002). The $2.12\ \mu\text{m}$ data was first published in Matsuura et al. (2006) and has been reprocessed using the ESO pipeline.

cm^{-2} . Using a typical column length of the H_2 emitting regions of $1'' (\approx 2.0 \times 10^{16}\ \text{cm})$, as used in Bujarrabal et al. (2002), we estimate an average density for the H_2 regions of $3.5 \times 10^6\ \text{cm}^{-3}$. This value is in good agreement with Alcolea et al. (2001) who estimate an average central density of $\sim 3.0 \times 10^6\ \text{cm}^{-3}$. The H_2 emission is most likely originating in the dense equatorial regions surrounding the central star.

4.1 Equatorial Region

The $1\rightarrow 0S(1)$ H_2 line map (see Fig. 1 [d]) shows two regions of H_2 oriented at a position angle (PA) of 113° , from brightest to faintest peak. If we assume that the distribution of the H_2 around the central region of OH231 is in a disc configuration then by measuring the major and minor axis we can estimate the inclination angle of the equatorial disc with respect to the plane of the sky from the H_2 data. Measurements for $R_{\text{max}}/R_{\text{min}}$ are determined by superposing a full ellipse onto the H_2 line map. We find $R_{\text{max}} = 4.06''$ and $R_{\text{min}} = 2.76''$ from the centre of the ellipse, and using the relation $i = \sin^{-1}(R_{\text{min}}/R_{\text{max}})$, we find $i = 43^\circ \pm 8$. This result is in good agreement with previously published values.

The location and orientation of this H_2 adds to the already complex picture of the equatorial region of OH231. Some previously reported structures, from smallest to largest, include:

- an equatorial torus of SiO maser emission ($\sim 2R_*$) that might lie on the innermost edge of an expanding SO disc (Sánchez Contreras et al. 2000, 2002);
- a centrally located compact disc of circumstellar material with inner $R = 0.03\text{--}0.04''/40\text{--}50\ \text{AU}$ (Matsuura et al. 2006);
- an opaque flared disc with outer $R = 0.25''/330\ \text{AU}$ revealed in mid-IR images (Jura et al. 2002);
- a slowly expanding disc with characteristic $R = 0.9''/1160\ \text{AU}$, detected via the SO ($J=2_2\rightarrow 1_1$) transition (Sánchez Contreras et al. 2000);
- a torus of OH maser emission with outer $R = 2.5''/3250\ \text{AU}$ (Zijlstra et al. 2001);
- an expanding hollow cylinder of HCO^+ with a characteristic radius equal to the OH torus radius (Sánchez Contreras et al. 2000).
- a halo of scattered light at $R \sim 4''/5200\ \text{AU}$ (Meakin et al. 2003);

From our observations the geometry of the H_2 region is un-




Figure 2. An integrated spectrum of the Western side of the central H₂ showing the detected lines, the position of the Br γ recombination line is shown for reference. Extracted spectra location marked in Fig. 1 (d).

clear, however we offer two possibilities: (1) A disc of H₂: Figure 1 (d) shows what might be interpreted as an incomplete disc of H₂, which fits with the series of concentric disc/tori structures listed above. To understand why we observe only emission from the edges and not the front of the disc, a comparison of the noise in the continuum at the front of the putative disc, to the amplitude of the 1 \rightarrow 0S(1) line peak shows both to be of the same order. We might then attribute the ‘missing’ H₂ in this region to variations in the continuum. We would not expect to observe the back of the disc due to the high level of extinction through the nebula. (2) A shell of H₂: if the H₂ is situated in an axisymmetric shell surrounding the central star, we might explain the geometry of the H₂ regions by assuming a density contrast between the poles and equator. This, combined with an increased column depth at the edge of the shell would manifest itself as two arcs of H₂ emission situated equatorially (as noted by Lowe & Gledhill 2005, for IRAS 19306+1407). This is supported by the detection of a shell of higher density gas and dust at the same location as the H₂ (Meakin et al. 2003). Both scenarios offer plausible explanations for the geometry of the H₂ emitting region, however further observations are needed in order to favour one.

We fit the 1 \rightarrow 0S(1) line profile yielding a $V_{\text{LSR}}=36\pm 17$ km s⁻¹ and FWHM=100 km s⁻¹ for both Eastern and Western regions of H₂ emission, a value consistent with the systemic velocity. Channel maps of the 1 \rightarrow 0S(1) line (Fig. 3) show no significant change in the distribution of the H₂.

In an attempt to explain the lack of reported H₂ in this object, we note two previous studies: (1) Weintraub et al. (1998) give a 3σ upper limit of 10^{-5} ergs cm⁻² s⁻¹ ster⁻¹ (1σ limit = 6.6×10^{-18} W m⁻²) for the surface brightness of the 1 \rightarrow 0S(1) line towards OH231. This places their measurement limit close to the 1 \rightarrow 0S(1) line strength given in Table 2, implying that the 1 \rightarrow 0S(1) line in their observations would be difficult to distinguish from noise, or possibly that their slit position, aligned East-West across inferred central star position, did not include the H₂ sites; (2) Matsuura et al. (2006), using the ISAAC instrument on VLT, do not report any detection of H₂. However, this can be explained due to the orientation of the slit, aligned from South-




Figure 3. ‘Channel maps’ of the four central channels of the 1 \rightarrow 0S(1) line from the central region of OH231. Inset labels show datacube channel and corresponding V_{LSR} velocity. Note: in ch794, the central region (marked by a green box) is residual from the continuum subtraction and not true H₂ emission.

West to North-East along the major axis (private comm.), with a slit-width of $0.8''$, i.e., the central H₂ emission site was not covered.

4.2 Northern Region

Figure 1 (c) shows the line map for the 1 \rightarrow 0S(1) transition in the Northern region. Most notable are the two knots of H₂, labeled A and B for the top and bottom knot, respectively. The H α emission contours are superposed on the H₂ line map. The peak intensity of the H₂ emission knots are slightly offset from the two H α emission knots (Fig. 1 [c]), however this small offset can be accounted for by the motion of the outflow (e.g. $V\sim 150$ km s⁻¹, from Bujarrabal et al. 2002) during the 10 years between both sets of observations. It is most likely that the optical and NIR line emission arise from the same shock event. Weaker 1 \rightarrow 0S(1) emission is noted tracing the H α edge of the bipolar outflow in the lower portion of the 1 \rightarrow 0S(1) line map. The NACO 2.12 μ m image does not show any trace of H₂ emission in this region (see contours in Fig. 1 [b]).

The 1 \rightarrow 0S(1) line is spread over \sim six spectral pixels, the four central channels are presented in Fig. 4. Examination of the line profile and channel maps allows us to probe the kinematics of the H₂ in this region, revealing two main features: 1) knot B which is persistent in all channel maps 2) the reduction of peak H₂ intensity in knot A at velocities ≤ -56 km s⁻¹. Both of these H₂ structures lie in the diffuse extended region (labeled B₁ in Fig. 4 of Bujarrabal et al. 2002) perpendicular to the axial flow, with a quoted H α velocity of 150 km s⁻¹ which is in good agreement with not only our deprojected H₂ velocities ($V_{\text{H}_2}\sim 110$ km s⁻¹) but also with HCO⁺ velocities (Sánchez Contreras et al. 2000).




Figure 4. 1→0S(1) line ‘channel maps’ of the four central channels extracted from the Northern region of OH231. Knot A and B are regions of most intense H₂ emission. Labels as in Fig.3.

In the case of knot A, the spectral line is strongly peaked in a single channel, while in knot B the spectral line peak is spread over two channels. Fitting the line profiles of knots A and B, yields LSR velocities of $V_{\text{LSR}} = -8 \text{ km s}^{-1}$, and -30 km s^{-1} , with deprojected velocities of -75 km s^{-1} and -110 km s^{-1} , respectively. This might indicate that further from the H α bow apex (Fig. 1 [a]), we are starting to see emission originating from the front and back of knot B, while knot A displays a narrower range of velocities, i.e., a single peak in its spectral profile. It is worth noting that due to the slit-length of ISAAC (120ʹ) coupled with the null H₂ detection discussed in §4.1, the H₂ in the Northern region might be confined to the wings of the bow shock.

The existence of fast moving shocked H₂ has previously been noted in other objects, for example, Cox et al. (2003) detect high velocity H₂ $\sim 220\text{--}340 \text{ km s}^{-1}$ (dependent on adopted inclination angle) in the outflows of AFGL 618. It is currently unknown exactly how shocked H₂ can be travelling at this speed without complete dissociation. Further high resolution mapping of the H₂ is necessary in order to resolve the shock surfaces, more accurately measure the H₂ kinematics, and apply shock models to this region.

5 CONCLUSIONS

We have presented VLT/SINFONI integral field observations of OH231, revealing the presence of several ro-vibrational H₂ lines. The main conclusions are:

- The discovery of H₂ emission near the centre of OH231, possibly located at the edge of an axisymmetric shell or an incomplete disc.
- A 1→0/2→1S(1) value of 8.3 ± 1.9 is found for the equatorial H₂, suggesting a collisional excitation mechanism.

- Our observations of the central shell/disc of H₂ show no velocity structure. However, higher S/N and/or velocity resolution data are needed to accurately probe the kinematics in this region.

- We detect fast-moving H₂ ($\sim 110 \text{ km s}^{-1}$, along the bipolar axis) via the 1→0S(1) transition along the North-Western tip of the nebula, a region where a strong H α bow shock exists. Due to the small FOV of our observations, the full extent of this H₂ is unknown.

6 ACKNOWLEDGMENTS

This research is funded by UH studentship and based on observations made with ESO Telescopes at the Paranal Observatory under programme ID 084.D-0487(A) and 072.D-0766(A). We thank Mikako Matsuura for providing ISAAC observation information. This research used the HLA (ID 8326) facilities of the STScI, the ST-ECF and the CADC with the support of the following granting agencies: NASA/NSF, ESA, NRC, CSA.

REFERENCES

- Alcolea J., Bujarrabal V., Sánchez Contreras C., Neri R., Zweigle J., 2001, A&A, 373, 932
- Black J. H., van Dishoeck E. F., 1987, ApJ, 322, 412
- Bujarrabal V., Alcolea J., Sánchez Contreras C., Sahai R., 2002, A&A, 389, 271
- Cohen M., Dopita M. A., Schwartz R. D., Tielens A. G. G. M., 1985, ApJ, 297, 702
- Cox P., Huggins P. J., Maillard J.-P., Muthu C., Bachiller R., Forveille T., 2003, ApJ, 586, L87
- Eisenhauer, F. et al., 2003, Proc. SPIE, 4841, 1548
- Frew D. J., Parker Q. A., 2010, PASA, 27, 129
- Hollenbach D., Natta A., 1995, ApJ, 455, 133
- Jura M., Chen C., Plavchan P., 2002, ApJ, 574, 963
- Jura M., Morris M., 1985, ApJ, 292, 487
- Kastner J. H., Weintraub D. A., Merrill K. M., Gatley I., 1998, AJ, 116, 1412
- Kastner J. H., Weintraub D. A., Zuckerman B., Becklin E. E., McLean I., Gatley I., 1992, ApJ, 398, 552
- Lee C.-F., Hsu M.-C., Sahai R., 2009, ApJ, 696, 1630
- Lowe K. T. E., Gledhill T. M., 2005, ASPC, 343, 282
- Matsuura M. et al., 2006, ApJ, 646, L123
- Meakin C. A., Bieging J. H., Latter W. B., Hora J. L., Tielens A. G. G. M., 2003, ApJ, 585, 482
- Sánchez Contreras C., Bujarrabal V., Neri R., Alcolea J., 2000, A&A, 357, 651
- Sánchez Contreras C., Desmurs J. F., Bujarrabal V., Alcolea J., Colomer F., 2002, A&A, 385, L1
- Sánchez Contreras C., Gil de Paz A., Sahai R., 2004, ApJ, 616, 519
- Smith M. D., 1995, A&A, 296, 789
- Smith M. D., Froebrich D., Eisloffel J., 2003, ApJ, 592, 245
- Weintraub D. A., Huard T., Kastner J. H., Gatley I., 1998, ApJ, 509, 728
- Zijlstra A. A., Chapman J. M., te Lintel Hekkert P., Likkell L., Comeron F., Norris R. P., Molster F. J., Cohen R. J., 2001, MNRAS, 322, 280