

Global Prevalence of Major Depressive Disorder, Generalized Anxiety, Stress and Depression Among Infertile Women: A Systematic Review and Meta-Analysis

Abstract

Background and purpose: In recent years, the global prevalence of infertility has increased among women, this is considered as a public health concern. One of the impacts of infertility is mental health problems in the patients, which can lead to complications such as stress, anxiety and depression. The aim of this study is to investigate the global prevalence of major depressive disorder, general anxiety, stress, and depression in infertile women through a systematic review and meta-analysis.

Methods: To identify studies that have reported the prevalence of major depressive disorder, generalized anxiety, stress and depression in infertile women, the PubMed, Scopus, Web of Science, Embase, ScienceDirect and Google Scholar repositories were systematically searched. Articles published up until February 2023 were included, whilst no lower time-limit was imposed in the search strategy. Heterogeneity of studies was examined using the I^2 test and thus random effects model was used to perform the analysis. Data analysis was conducted within the Comprehensive Meta-Analysis (v.2) software.

Findings: In the review of 44 studies with a sample size of 53,300 infertile female patients, the overall prevalence of major depressive disorder (clinical depression), generalized anxiety, stress, and depression was found to be 22.9%, 13.3%, 78.8%, and 31.6% respectively. It was also found that mental health complications are more prevalent among infertile women in Asia (continent).

Conclusion: Considering the prevalence of mental disorders among infertile women, health policymakers can use the results of the present meta-analysis to pay more attention to the mental health of infertile women and devise suitable interventions and programs to reduce and prevent the spread of psychological disorders among infertile women.

Keywords: Infertile women, Depression, Major Depression, Anxiety, Stress, General Anxiety Disorder, GAD

Introduction

Infertility in the field of reproductive health is considered as a global concern, and is one of the most common chronic disorders, regardless of age (1, 2). Infertility refers to the inability to reproduce; as a definition, infertility is when the female or male reproductive systems fail to conceive after 12 months or more of regular unprotected sexual intercourse (3).

Infertility affects millions of people of reproductive age around the world and affect their communities (4). Estimates show that 48 million couples worldwide have infertility problems (4). The prevalence of infertility has been increasing globally over the last few decades, so that from 1990 to

2017 infertility rate has had an annual increase of 0.370% in women and 0.291% in men (2) and it has reached 12 percent (from 8 per cent) among reproductive couples (5). Most couples have an identifiable cause for infertility, whilst the cause among the rest is unexplained. Female partners are involved in 40-55% of infertility cases and male partners in about 20-40% (4).

Women are more psychologically and emotionally affected by infertility than men (6). The main reason for this is related to the stigma around women not having children, especially in cultures that consider this aspect as the main goal of life and the defining role of women. On the other hand, still in many developing and developed societies, a woman is considered a complete person only when she becomes a mother, therefore, women with infertility are socially isolated, neglected, more prone to divorce, and severely reduced in dignity. They feel a reduction in their ego and value, which can lead to feelings of guilt and that their lives are meaningless (7-9).

Although some women cope with infertility and yet have a positive and profound life (10), the experience of infertility is associated with a wide range of psychological problems such as depression (11, 12), stress, anxiety (13, 14), low self-esteem (15, 16), low psychological adjustment (17), feelings of fear, anger, shame, jealousy, loneliness, despair, emotional instability, insufficient sexual influence, and sexual dysfunction (18). Infertility patients face complex challenges that fall in the biological, psychological, social, and moral domains (19). Accordingly, most infertile women have higher levels of stress, anxiety, and depression (20, 21), in such a way that anxiety and depression in infertile women are comparable to what is experienced among cancer patients (22).

Considering the high rate of infertility globally, and the prevalence of problems related to mental health in infertile people, especially the high prevalence of stress, anxiety and depression in women with infertility, as well as the importance of mental health of infertile women, we decided to conduct a systematic review and meta-analysis on the global prevalence of major depressive disorder, general anxiety, stress and depression among women with infertility. It is strongly believed that the findings from this meta-analysis can provide useful insights for health policymakers to devise appropriate intervention programs.

Materials and method

We conducted our initial search in January 2023 using the PubMed, Web of Science, Google Scholar, Scopus, ScienceDirect and Embase databases. The keywords of Infertile Women, Barren Women, Sterile Women, Generalized Anxiety Disorder, GAD, Major Depression, Stress, and Anxiety and their combinations were used to undertake the searches. To maintain the comprehensiveness of the searches, no restrictions were placed upon year of publication of articles. Subsequently, information from the identified articles were transferred into the EndNote reference management software. Additionally, reference lists used in the identified articles were manually reviewed to ensure that grey and previously unfound, yet relevant, studies are also included. The searches were last updated in February 2023.

Inclusion and Exclusion Criteria

The inclusion criteria for study selection were:

1. Studies that reported the prevalence of major depressive disorder, generalized anxiety, stress and depression in women with infertility,
2. Studies with their full-text available, and
3. Studies that provided sufficient data (sample size, prevalence).

Exclusion criteria were:

1. Case reports, and case series studies,
2. Review studies of any sort,
3. Duplicates, and

4- Studies with insufficient data (lack of information about prevalence and sample size).

Study Selection

The study selection was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Initially, studies that were repeated in different databases were excluded from the systematic review, and only one copy was retained. The initial screening took place by reviewing the titles and abstracts of the studies, and irrelevant articles were removed based on the inclusion and exclusion criteria. Then, the full texts of the remaining articles were evaluated based on inclusion and exclusion criteria, and similarly, irrelevant studies were omitted. To avoid bias, all the steps of reviewing sources and data extraction were completed by two researchers independently. If there was a difference of opinion between the two researchers (reviewers) in some of the articles, consensus was achieved with the support of a third reviewer.

Quality Evaluation

To evaluate the quality of the remaining articles, a checklist appropriate for the inspection of observational studies was adopted. The Strengthening the Reporting of Observational Studies in Epidemiology checklist (STROBE) consists of six headings including: title, abstract, introduction, methods, results, and discussion. In total, this instruction consists of 32 subheadings. These 32 subheadings include different methodological aspects of the study, e.g., title, statement of the problem, objectives of the study, type of study, statistical population of the study, sampling method, determination of the appropriate sample size, definition of variables and procedures, study data collection tools, statistical analysis methods, and findings. An article scored a point once a subheading was fulfilled, and accordingly articles with a score of 16 and above were considered to be of average and high methodological quality respectively. Articles with a score below 16 were deemed to be of poor quality and were therefore excluded from our work.

Data Extraction

Data extraction was completed by two researchers using a different pre-prepared checklist. This checklist includes the following headings: first author's name, year of publication, study location, sample size, age group of women, prevalence of different disorders, and study instrument(s).

Statistical Analysis

The reported results from the selected studies were extracted into the Comprehensive Meta-Analysis (CMS v.2) software, and the heterogeneity of the studies was examined through the I^2 test. Publication bias was assessed using the Egger's test at a significance level of 0.05 and corresponding Funnel plots were drawn.

Findings

Following the searchers, 453 articles were found from the selected databases and 2 further possible related articles were identified through manual search, and details of all found articles were transferred into the EndNote reference management software. Subsequently, 128 articles were excluded due to duplication. In the screening phase, titles and abstracts of the studies were reviewed and 136 articles were excluded based on the inclusion and exclusion criteria. In the eligibility evaluation stage, a further 133 articles were excluded through the review of full texts and

in accordance with the inclusion and exclusion criteria. In the quality evaluation phase, through the study of the full text of the articles and based on the scores obtained from the STROBE checklist, studies that had poor methodological quality were excluded, and finally 44 studies were included for final evaluation. The information of these 44 studies is reported in Tables 1, 2, 3, 4, and 5.

In the studies included in Table 1., the highest prevalence of stress among infertile women (92.71%) is related to the study of Teklemicheal et al. in 2022 which used the Copenhagen Multi-Centre Psychosocial Infertility-Fertility Problem Stress Scales (COMPI-FPPS) tool in the age group of 20 to 48 years (23); and the lowest prevalence Stress in infertile women (53.06%) is reported by Xiaoli et al.'s study in 2016 which used the WHOQOL-100 tool (World Health Organization QoL) in the age group of 23 to 41 years (24). Moreover, the overall pooled prevalence of stress among infertile women is 78.8% (95% CI: 25.8%-97.5%). Two studies used COMPI-FPPS and WHOQOL-100 tools to measure stress, which are presented in Table 1.

In the studies included in Table 2, the highest prevalence of major depression in infertile women (68.90%) is reported in a study by Al-Asadi et al. in 2015 which used the ICD-10 tool in the age group of 16 to 45 years (25). The lowest prevalence of major depression in infertile women (70.5%) reported in the study of Volgsten et al. in 2018 which adopted the Primary Care Evaluation of Mental Disorders (PRIME-MD) tool based on DSM-IV in an average age group of 38.3 ± 3.9 years (26). In our meta-analysis, the overall pooled prevalence of major depression among infertile women is found as 22.9% (95% CI: 11%-41.6%).

In the studies included in Table 3, the highest prevalence of generalized anxiety disorder in infertile women (44.10%) is related to a study conducted by Shabani et al. in 2010 which had adopted the General Health Questionnaire (GHQ) and diagnostic calculation (27); the lowest prevalence of generalized anxiety disorder in Infertile women (80.1%) is reported in Volgsten et al. which was conducted in 2018 using the PRIME-MD tool based on DSM-IV in an average age of 38.3 ± 3.9 years (26). In our meta-analysis, the overall pooled prevalence of generalized anxiety disorder in infertile women is found to be 13.3% (95% CI: 6.5%-25.3%).

In the studies included in Table 4, the highest prevalence of depression in infertile women (82%) is related to the work of Dadhwal et al. in 2022 which used the HDRS tool on a sample with an average age of 29.21 ± 3.74 years (28); the lowest prevalence of depression in women with infertility (60.2%) is in the study of Sejbaek et al. This study was in 2013 using the ICD-8 and ICD-10 tools (29). In our meta-analysis the overall pooled prevalence of depression in infertile women is 31.6% (95% CI: 21%-44.6%).




Figure 1: PRISMA Flow Diagram for Study Selection.

Table 1: Summary of Characteristics of Included Studies Related to the Prevalence of Stress in Women with Infertility.

Author	Year	Country	Age range	Sample size	Prevalence of stress in infertile women	Instrument
Teklemicheal et al (23)	2022	Ethiopia	20-48	96	92.71%	COMPI-FPPS*

Xiaoli et al (24)	2016	China	23-41	81	53.06%	WHOQOL-100**
-------------------	------	-------	-------	----	--------	--------------

* Copenhagen Multi-Centre Psychosocial Infertility-Fertility Problem Stress Scales

** World Health Organization Quality of Life

Table 2: Summary of Characteristics of Included Studies Related to the Prevalence of Major Depression in Women with Infertility.

Author	Year	Country	Age range	Sample size	Prevalence of MD in infertile women	Instrument
Carvalho et al (30)	2021	Brazil	35.89 ± 5.20	90	9.90%	(M.I.N.I.)* Brazilian version 5.0.0, DSM IV**
Volgsten et al (26)	2018	Sweden	38.3 ± 3.9	278	5.70%	(PRIME-MD) *** based on DSM-IV
Al-Asadi et al (25)	2015	Iraq	16-45	251	68.90%	(ICD-10) ****
Saffarieh et al (31)	2020	Iran	18-46	30	16.7%	BDI ***** , STAI *****
Holley et al (32)	2015	USA	23-52	174	39.10%	Composite International Diagnostic Interview Major Depression module, a structured diagnostic interview
Chen et al (33)	2004	Taiwan	24-45	112	17.00%	MINI
Osman et al (34)	2022	Egypt	18-59	371	30.20%	PHQ-9*****

* Mini International Neuropsychiatric Interview

** Diagnostic and Statistical Manual of Mental Disorders, version 4

*** Primary Care Evaluation of Mental Health Disorders

**** International Classification of Diseases-Version 10

***** Beck Depression Inventory

***** State-Trait Anxiety Inventory

***** Patient Health questionnaire

Table 3: Summary of Characteristics of Included Studies Related to the Prevalence of General Anxiety Disorder in Women with Infertility.

Author	Year	Location	Age	Sample size	Prevalence of GAD IN infertile women	Instrument
--------	------	----------	-----	-------------	--------------------------------------	------------

Carvalho et al (30)	2021	Brazil	35.89 ± 5.20	90	8.90%	(M.I.N.I.) Brazilian version 5.0.0, DSM IV
Volgsten et al (26)	2018	Sweden	38.3 ± 3.9	278	1.80%	(PRIME-MD) based on DSM-IV
Gui et al (35)	2021	China	24-48	693	21.80%	GAD-7*
Aghanwa et al (36)	1999	Nigeria	...	37	2.70%	PSE** & clinical evaluation
Chen et al (33)	2004	Taiwan	24-45	112	23.20%	MINI
Shabani et al (27)	2010	Iran	...	353	44.10%	GHQ*** & interviews

*Generalized Anxiety Disorder

** Present State Examination

*** General Health Questionnaire

Table 4: Summary of Characteristics of Included Studies Related to the Prevalence of Depression in Women with Infertility.

Author	Year	Location	Age	Women sample size	Prevalence of depression IN infertile women	Instrument
Cui et al (37)	2021	China	22-47	536	27.90%	HADS*
Yangin et al (38)	2015	Turkey	19-43	102	26.50%	BDI
Dong et al (39)	2021	China	21-44	715	19.30%	PHQ-9
Jin et al (40)	2013	China	22-41	460	14.80%	ZAD**
Lakatos et al (41)	2017	Hungary	33.30 ± 4.85	134	44.80%	BDI
Huang et al (42)	2019	China	35.7 ± 5.29	97	30.00%	BDI-II
Naab et al (43)	2013	Ghana	30-39	203	53%	CES-D***
Sakulsaengprapha et al (44)	2019	Thailand	36.5 ± 4.6	421	3%	HADS
Drosdzol et al (45)	2009	Polish	29.8 +4.1	206	35.44%	BDI
Aghanwa et al (36)	1999	Nigeria	...	37	27.00%	PSE & clinical evaluation
Crawford et al (46)	2017	North Carolina	...	416	41%	NIH PROMIS (short form)****
Kato et al (47)	2021	Japan	24-46	513	54%	QIDS*****
Czyżkowska et al (48)	2016	Poland	18-40	50	78%	BDI

Bondade et al (49)	2018	India	26.73 ± 4.23	100	25%	DSM-5
Alhassan et al (50)	2014	Ghana	30.5±6.3	100	62%	BDI
Audier-Bourgain et al (51)	2021	France	24-42	61	8.20%	HADS
Holley et al (32)	2015	USA	23-52	174	24.00%	CESD
Cho et al (52)	2019	Korea	36.16±4.55	118	37.30%	BDI
Khan et al (53)	2020	Pakistan	28.46±7.68	160	75%	HADS
Suna et al (54)	2016	Turkey	20-40	30	23.30%	BDI
Salomão et al (55)	2018	Brazil	32-38	140	11.42%	HADS
Domar et al (56)	1992	338	37%	BDI
Domar et al (56)	1992	338	25%	CES-D
Sulyman et al (57)	2019	Nigeria	19-43	207	25.60%	HADS
Li et al (58)	2021	China	20-37	202	59.9%	SDS*****
Peyvandi et al (59)	2011	Iran	18-48	200	62.00%	BDI
Carreño Meléndez et al (60)	2007	Spain	19-42	240	32.10%	SDS
Deeks et al (61)	2010	Australia	...	22	67.7%	HADS
Sezgin et al (62)	2016	Turkey	21-47	100	33%	HADS
Wright et al (63)	1991	Canada	29.5 ± 4.5	449	23.20%	Psychiatric Symptom Inventory
Alosaimi et al (64)	2015	Saudi Arabia	31.5	206	26.20%	MINI
Herbert et al (65)	2010	Australia	...	1031	26.70%	CES-D10
Sejbaek et al (29)	2013	Denmark	...	42915	2.60%	ICD-8, ICD-10
Upkong et al (66)	2006	Nigeria	24 – 46	112	42.90%	BDI
Dadhwal et al (28)	2022	India	29.21 ± 3.74	150	58% (only dep), 24%(dep+anx)	HDRS*****
Shabani et al (27)	2010	Iran	...	353	30.40%	GHQ & interviews

* Hospital Anxiety and Depression Scale

** Zung Self-Rating Depression Scale

*** Center for Epidemiologic Studies for Depression

**** National Institutes of Health Patient Reported Outcomes Measurement Information System

***** Quick Inventory of Depressive Symptomatology

***** Self-Rating depression Scale

***** Hamilton Depression Rating Scale

Stress

In the review of 2 studies with a total sample size of 177 people, the I^2 heterogeneity test showed high heterogeneity (I^2 : 96.5), and therefore random effects method was adopted to analyze the results. Based on the meta-analysis, the prevalence of stress among infertile women is found as 78.8% (95%CI: 25.8%-97.5%) (Figure 2). It should be noted that it was not possible to check the publication bias in the studies due to the presence of only 2 studies in this category.




Figure 2: Forest Plot of Prevalence of Stress in Infertile Women Based on Random Effects Method.

Major Depression

In the review of 7 studies with a sample size of 1306 people, the I^2 test showed high heterogeneity (I^2 : 97.3), and accordingly random effects method was adopted to analyze the results. Based on the meta-analysis, the pooled prevalence of major depression in infertile women is found to be 22.9% (95%CI: 11%-41.6%) (Figure 3). Test of publication bias using the Egger's test showed the absence of publication bias among the studies (p : 0.179) (Figure 4).




Figure 3: Forest Plot of Prevalence of Major Depression Based on Random Effects Method.




Figure 4: Funnel Plot for Assessing Publication Bias in the Reviewed Studies Related to Major Depression.

Generalized Anxiety Disorder (GAD)

In the review of 6 studies with a sample size of 1563 people, the I^2 test showed high heterogeneity (I^2 : 95.9), and accordingly, random effects method was adopted to analyze the results. Based on the meta-analysis, the pooled prevalence of Generalized Anxiety Disorder among infertile women is 13.3% (95% CI: 6.5%-25.3%) (Figure 5). Test of publication bias using the Egger's test showed the absence of publication bias in the studies (p : 0.191) (Figure 6).




Figure 5: Forest Plot of Prevalence of Generalized Anxiety Disorder Based on Random Effects Method.




Figure 6: Funnel Plot for Assessing Publication Bias in the Reviewed Studies Related to GAD.

Depression

In the review of 36 studies with a sample size of 51,636 people, the I^2 test showed high heterogeneity (I^2 : 99.4), and therefore, random effects method was used to analyze the results. According to our meta-analysis, the pooled prevalence of depression among women Infertility is 31.6% (95% CI: 21%-44.6%) (Figure 7). Assessment of publication bias using the Egger's test indicated the existence of publication bias in the studies (p : 0.000) (Figure 8).




Figure 7: Forest Plot of Prevalence of Depression Based on Random Effects Method.




Figure 8: Funnel Plot for Assessing Publication Bias in the Reviewed Studies Related to Depression.

Discussion

Infertility is considered as a concern for women in several communities and cultures, where there is a strong emphasis on women being mothers or wives (67-69). Accordingly, there are numerous cases where a woman's infertility has an impact on her marriage and life (70). Since pregnancy and the role of mother are viewed as specific to women, social expectations from women are also higher compared to men. Moreover, the fear of social stigma of infertility is greater among women (71). Considering these, infertility is deemed as a personal crisis that can cause severe mental and emotional stress and pressure on couples and can negatively impact their mental health in various ways (72).

The present study was a systematic review and meta-analysis to analyze reported results of relevant literature on the global prevalence of major depressive disorder, generalized anxiety, stress and depression among infertile women. Based on the results of this study, the overall pooled prevalence of major depressive disorder, generalized anxiety disorder, stress, and depression in infertile women is 22.9%, 13.3%, 78.8%, and 31.6%, respectively.

Many studies have reported high prevalence of symptoms of depression, anxiety and stress among infertile women (73-78). According to a study by Alhassan et al., 62% of infertile women in Ghana suffer from depression (50). The prevalence of depression among Chinese infertile women is reported 69% (79). Depression is also highly prevalent among infertile women in Japan and Gambia (80, 81). Jones et al. (1993) also stated that there is mild to moderate depression in 28.3%, moderate to severe depression in 7.2%, and the most severe form of depression in 1.2% of infertile women (82).

On the other hand, Noorbala et al. argued that 48% of infertile women experience depression (83). The rate of depression among infertile Bahraini women was reported as 32.5%, whereas this figure in infertile Tunisian women was 46.6% (84). Similarly, depression rate among infertile Iranian women was 40% (85, 86). Oddens et al.'s study (1999) stated that 24.9% of infertile women had depression

disorders (87). According to the results of the present study, the overall prevalence of major depressive disorder in infertile women is 22.90% and the prevalence of depression is 31.6%.

In some countries, socio-cultural and religious norms allow husbands to have multiple marriages, and a woman's infertility provides them with a reason for this decision. In addition, having a child, a boy, gives women a sense of pride and security for old age, which is why infertility causes many psychological problems for women. However, in developed countries, psychological disorders are less common due to women's legal rights, their participation in professional life and their contribution to the family economy (8, 72, 88). For this reason, the results of our study provide a lower prevalence of depression disorders reported for some countries, especially developing countries, due to the inclusion of different cultures.

According to literature, patients with infertility experience lower quality of life and higher levels of anxiety (89-92). Lawson et al., Allen, and Kraaji et al. have shown higher levels of anxiety in infertile women (93-95). Another study found that 67% of infertile women suffered from anxiety (79). The prevalence of generalized anxiety disorder in infertile women in Taiwan has been reported as 23.2% (96). Anxiety was investigated in 130 infertile women in China, and the results showed different levels of psychological stress among 83.8% of infertile women; in addition, moderate or severe types were observed in 25% (97). In another study, depression and/or anxiety disorders were present in 33% (Hong Kong) (87), and 32% (Scotland) of infertile women (98). In our study, the overall pooled prevalence of generalized anxiety disorder in infertile women is 13.3%, and the results of these studies are in line with our findings. The lower prevalence of anxiety in our review is due to the fact that we only examined generalized anxiety disorder and did not examine other types of anxiety.

The stress levels of infertile women in developing countries such as Cameroon (99), India (100) and Nigeria (101) have been reported as 84.6%, 80% and 87.2%, respectively. On the other hand, in developed countries such as Sweden, 30.8% of stress has been reported for infertile women (102). Similarly, Luk et al., Dooley et al., and El Kissi et al. showed in their studies that stress has a higher prevalence among infertile women (78, 103, 104). The results of these studies are in line with our findings (78.8%). Since cultural and social issues that cause stress in infertile women is more prevalent in some regions, especially in Asian countries, the results of our research are closer to the studies conducted in the Asia (continent).

There are some limitations in our work. One of the limitations of this study is that several articles were excluded from the review due to their low quality, e.g., by not reporting the prevalence figure. Additionally, the level of anxiety, depression and stress among infertile women can be affected by various factors such as the role of the life partner, economic conditions, etc.

Conclusion

According to the results of the present study, the overall prevalence of major depressive disorder, general anxiety, stress and depression in infertile women is 22.9%, 13.3%, 78.8% and 31.6% respectively. According to the results, women suffering from infertility also experience many psychological disorders. In this way, paying attention to the mental health of infertile women is an crucial, and in addition to the treatment of infertility, taking care of the mental health of infertile women is of special importance. Therefore, health policymakers can potentially use the results of the present meta-analysis to gain insights on the prevalence of mental disorders among infertile women. Moreover, policymakers can plan appropriate intervention programmes in response to the psychological problems of infertile women.

References

1. Tomaiuolo R, Veneruso I, Cariati F, D'Argenio V. Microbiota and human reproduction: the case of female infertility. *High-throughput*. 2020;9(2):12.
2. Sun H, Gong T-T, Jiang Y-T, Zhang S, Zhao Y-H, Wu Q-J. Global, regional, and national prevalence and disability-adjusted life-years for infertility in 195 countries and territories, 1990–2017: results from a global burden of disease study, 2017. *Aging (Albany NY)*. 2019;11(23):10952.
3. World Health Organization (WHO). *International Classification of Diseases, 11th Revision (ICD-11)*. Geneva: WHO2018.
4. WHO: Infertility. [Jul; 2022]. 2022. <https://www.who.int/news-room/fact-sheets/detail/infertility>.
5. Vander Borgh M, Wyns C. Fertility and infertility: Definition and epidemiology. *Clinical biochemistry*. 2018;62:2-10.
6. Maroufizadeh S, Navid B, Omani-Samani R, Amini P. The effects of depression, anxiety and stress symptoms on the clinical pregnancy rate in women undergoing IVF treatment. *BMC research notes*. 2019;12(1):1-4.
7. Hawkey AJ, Ussher JM, Perz J. “If you don’t have a baby, you can’t be in our culture”: Migrant and refugee women’s experiences and constructions of fertility and fertility control. *Women's reproductive health*. 2018;5(2):75-98.
8. Vioreanu A-M. The psychological impact of infertility. *Directions for the development of interventions. Mental Health: Global Challenges Journal*. 2021;4(1).
9. Vioreanu A-M. Using CBT for Depression: A Case Study of a Patient with Depressive Disorder Due to a Medical Condition (Infertility). *Mental Health: Global Challenges Journal*. 2023;6(1):2-15.
10. Kiani Z, Simbar M, Hajian S, Zayeri F. Quality of life among infertile women living in a paradox of concerns and dealing strategies: a qualitative study. *Nursing Open*. 2021;8(1):251-61.
11. Yazdi HZG, Sharbaf HA, Kareshki H, Amirian M. Infertility and psychological and social health of Iranian infertile women: a systematic review. *Iranian journal of psychiatry*. 2020;15(1):67.
12. Omani-Samani R, Maroufizadeh S, Almasi-Hashiani A, Amini P. Prevalence of depression and its determinant factors among infertile patients in Iran based on the PHQ-9. *Middle East Fertility Society Journal*. 2018;23(4):460-3.
13. Zurlo MC, Cattaneo Della Volta MF, Vallone F. Infertility-related stress and psychological health outcomes in infertile couples undergoing medical treatments: testing a multi-dimensional model. *Journal of Clinical Psychology in Medical Settings*. 2020;27:662-76.
14. Maroufizadeh S, Ghaheri A, Almasi-Hashiani A, Mohammadi M, Navid B, Ezabadi Z, et al. The prevalence of anxiety and depression among people with infertility referring to Royan Institute in Tehran, Iran: A cross-sectional questionnaire study. *Middle East fertility society journal*. 2018;23(2):103-6.
15. Jamil S, Shoaib M, Aziz W, Ather MH. Does male factor infertility impact on self-esteem and sexual relationship? *Andrologia*. 2020;52(2):e13460.

16. Malekzadeh F, Navid B, Mohammadi M, Samani RO. Self-esteem and defense mechanisms in infertile couples based on the cause of infertility in Royan Infertility Treatment Center in 2017. *Journal of Shahid Sadoughi University of Medical Sciences*. 2020.
17. Zarif Golbar Yazdi H, Aghamohammadian Sharbaf H, Kareshki H, Amirian M. Psychosocial consequences of female infertility in Iran: a meta-analysis. *Frontiers in psychiatry*. 2020;11:518961.
18. Kim M, Kim HS. Mediator effect of marital intimacy on the relationship between depression and marital satisfaction of infertile women. *Journal of Korean Public Health Nursing*. 2018;32(1):96-108.
19. Massarotti C, Gentile G, Ferreccio C, Scaruffi P, Remorgida V, Anserini P. Impact of infertility and infertility treatments on quality of life and levels of anxiety and depression in women undergoing in vitro fertilization. *Gynecological Endocrinology*. 2019;35(6):485-9.
20. Kiani Z, Simbar M, Hajian S, Zayeri F. The prevalence of depression symptoms among infertile women: a systematic review and meta-analysis. *Fertility research and practice*. 2021;7:1-10.
21. Lakatos E, Szigeti JF, Ujma PP, Sexty R, Balog P. Anxiety and depression among infertile women: a cross-sectional survey from Hungary. *BMC women's health*. 2017;17(1):1-9.
22. Villar RR, Fernández SP, Garea CC, Pillado M, Barreiro VB, Martín CG. Quality of life and anxiety in women with breast cancer before and after treatment. *Revista latino-americana de enfermagem*. 2017;25.
23. Teklemicheal AG, Kassa EM, Weldetensaye EK. Prevalence and correlates of infertility related psychological stress in women with infertility: a cross-sectional hospital based survey. *BMC psychology*. 2022;10(1):91.
24. Xiaoli S, Mei L, Junjun B, Shu D, Zhaolian W, Jin W, et al. Assessing the quality of life of infertile Chinese women: a cross-sectional study. *Taiwanese Journal of Obstetrics and Gynecology*. 2016;55(2):244-50.
25. Al-Asadi JN, Hussein ZB. Depression among infertile women in Basrah, Iraq: Prevalence and risk factors. *JOURNAL OF THE CHINESE MEDICAL ASSOCIATION*. 2015;78(11):673-7.
26. Volgsten H, Schmidt L, Svanberg AS, Ekselius L, Poromaa IS. Psychiatric disorders in women and men up to five years after undergoing assisted reproductive technology treatment ? a prospective cohort study. *HUMAN FERTILITY*. 2019;22(4):277-82.
27. Shabani A, Shabani A. Anxiety and mood disorders in infertile women referred to neuropsychological department due to mental disorder because of infertility. *Iranian Journal of Reproductive Medicine*. 2010;8:11-2.
28. Dadhwal V, Choudhary V, Perumal V, Bhattacharya D. Depression, anxiety, quality of life and coping in women with infertility: A cross-sectional study from India. *International Journal of Gynecology and Obstetrics*. 2022;158(3):671-8.
29. Sejbaek CS, Hageman I, Pinborg A, Hougaard CO, Schmidt L. Incidence of depression and influence of depression on the number of treatment cycles and births in a national cohort of 42 880 women treated with ART. *HUMAN REPRODUCTION*. 2013;28(4):1100-9.

30. Carvalho CF, Mattia MMC, da Silva H, Bredemeier F, Arpini NE, Chapon R, et al. Psychotropic medication use among women seeking assisted reproductive technology (ART) therapy: A cross-sectional study. *JOURNAL OF AFFECTIVE DISORDERS*. 2021;292:386-90.
31. Saffarieh E, Sharami SRY, Nasiri S, Aghaamoo S, Ziari A. Evaluation of depression and anxiety in couples with infertility and related factors. *Tehran University Medical Journal*. 2020;78(8):522-7.
32. Holley SR, Pasch LA, Bleil ME, Gregorich S, Katz PK, Adler NE. Prevalence and predictors of major depressive disorder for fertility treatment patients and their partners. *FERTILITY AND STERILITY*. 2015;103(5):1332-9.
33. Chen TH, Chang SP, Tsai CF, Juang KD. Prevalence of depressive and anxiety disorders in an assisted reproductive technique clinic. *HUMAN REPRODUCTION*. 2004;19(10):2313-8.
34. Osman DM, Ahmed GK, Farghal MM, Ibrahim AK. Prevalence and predictors of depressive symptoms among married Egyptian women: a multicenter primary healthcare study. *BMC PSYCHIATRY*. 2022;22(1).
35. Gui WW, Yang X, Jiang HM, Wu HW, Zeng M, Wen YD, et al. Prevalence of anxiety and its associated factors among infertile patients after 'two-child' policy in Chongqing, China: a cross-sectional study. *REPRODUCTIVE HEALTH*. 2021;18(1).
36. Aghanwa HS, Dare FO, Ogunniyi SO. Sociodemographic factors in mental disorders associated with infertility in nigeria. *Journal of Psychosomatic Research*. 1999;46(2):117-23.
37. Cui C, Wang L, Wang X. Effects of self-esteem on the associations between infertility-related stress and psychological distress among infertile chinese women: A cross-sectional study. *Psychology Research and Behavior Management*. 2021;14:1245-55.
38. Yangin H, Kukulu K, Gulşen S, Aktaş M, Sever B. A survey on the correlation between sexual satisfaction and depressive symptoms during infertility. *Health Care for Women International*. 2016;37(10):1082-95.
39. Dong M, Xu X, Li Y, Wang Y, Jin Z, Tan J. Impact of infertility duration on female sexual health. *Reproductive Biology and Endocrinology*. 2021;19(1).
40. Jin X, Wang GX, Liu SS, Zhang J, Zeng F, Qiu Y, et al. Survey of the Situation of Infertile Women Seeking In Vitro Fertilization Treatment in China. *BIOMED RESEARCH INTERNATIONAL*. 2013;2013.
41. Lakatos E, Szigeti JF, Ujma PP, Sexty R, Balog P. Anxiety and depression among infertile women: a cross-sectional survey from Hungary. *BMC WOMENS HEALTH*. 2017;17.
42. Huang L-H, Kuo C-P, Lu Y-C, Lee M-S, Lee S-H. Association of emotional distress and quality of sleep among women receiving in-vitro fertilization treatment. *Taiwanese Journal of Obstetrics and Gynecology*. 2019;58(1):168-72.
43. Naab F, Brown R, Heidrich S. Psychosocial Health of Infertile Ghanaian Women and Their Infertility Beliefs. *JOURNAL OF NURSING SCHOLARSHIP*. 2013;45(2):132-40.
44. Sakulsaengprapha W, Prechapanich J, Petyim S. Prevalence of anxiety and depression in infertile women in Siriraj hospital by Using Thai hospital anxiety and depression scale. *Thai Journal of Obstetrics and Gynaecology*. 2019;27(1):38-46.

45. Drosdzol A, Skrzypulec V. Depression and anxiety among polish infertile couples-an evaluative prevalence study. *Journal of Psychosomatic Obstetrics and Gynecology*. 2009;30(1):11-20.
46. Crawford NM, Hoff HS, Mersereau JE. Infertile women who screen positive for depression are less likely to initiate fertility treatments. *Human Reproduction*. 2017;32(3):582-7.
47. Kato T, Sampei M, Saito K, Morisaki N, Urayama KY. Depressive symptoms, anxiety, and quality of life of Japanese women at initiation of ART treatment. *SCIENTIFIC REPORTS*. 2021;11(1).
48. Czyzkowska A, Awruk K, Janowski K. Sexual Satisfaction and Sexual Reactivity in Infertile Women: The Contribution of The Dyadic Functioning and Clinical Variables. *INTERNATIONAL JOURNAL OF FERTILITY & STERILITY*. 2016;9(4):465-76.
49. Bondade S, Iyengar RS, Shivakumar BK, Karthik KN. Intimate partner violence and psychiatric comorbidity in infertile women - A cross-sectional hospital based study. *Indian Journal of Psychological Medicine*. 2018;40(6):540-6.
50. Alhassan A, Ziblim AR, Muntaka S. A survey on depression among infertile women in Ghana. *BMC Women's Health*. 2014;14(1).
51. Audier-Bourgain M, Baubet T, Pham-Scottez A, Corcos M, Nicolas I. Eating disorders and sexuality: A quantitative study in a French medically assisted procreation course. *BRAIN AND BEHAVIOR*. 2021;11(8).
52. Cho EY, Sung MH. Effects of irrational parenthood cognition, family support, and resilience on depression of infertile women. *Korean Journal of Women Health Nursing*. 2019;25(1):60-72.
53. Khan I, Khan MM, Fareed A, Noor UI A, Fatima, Khan MK. Determine the Frequency of Depression in Females with Infertility. *PAKISTAN JOURNAL OF MEDICAL & HEALTH SCIENCES*. 2020;14(3):1728-30.
54. Suna KK, Ilay G, Aysenur A, Han GK, Ulku UE, Pasa U, et al. Effects of Infertility Etiology and Depression on Female Sexual Function. *JOURNAL OF SEX & MARITAL THERAPY*. 2016;42(1):27-35.
55. Salomão P, Romão A, Reis R, Navarro P, Lerri M, Rosa-E-Silva A, et al. Sexual function of women under treatment for infertility. *Journal of Sexual Medicine*. 2015;12:60.
56. Domar AD, Zuttermeister PC, Friedman R. THE PSYCHOLOGICAL IMPACT OF INFERTILITY - A COMPARISON WITH PATIENTS WITH OTHER MEDICAL CONDITIONS. *JOURNAL OF PSYCHOSOMATIC OBSTETRICS AND GYNECOLOGY*. 1993;14:45-52.
57. Sulyman D, Ayanda KA, Aminu BM, Dattijo LM. Anxiety and depressive disorders among infertile women attending clinic in a Nigeria teaching hospital. *African Journal of Biomedical Research*. 2019;22(2):157-65.
58. Li G, Jiang Z, Kang X, Ma L, Han X, Fang M. Trajectories and predictors of anxiety and depression amongst infertile women during their first IVF/ICSI treatment cycle. *Journal of Psychosomatic Research*. 2021;142:110357.
59. Peyvandi S, Hosseini SH, Daneshpoor SMM, Mohammadpour RA, Qolami N. The prevalence of depression, anxiety and marital satisfaction and related factors in infertile women referred to infertility clinics of Sari city in 2008. *Journal of Mazandaran University of Medical Sciences*. 2011;20(80):25-32.

60. Carreño Meléndez J, Morales Carmona F, Sánchez Bravo C, Henales Almaraz C, Espindola Hernández JG. [An explanation of depression and anxiety symptoms in sterile women]. *Ginecol Obstet Mex.* 2007;75(3):133-41.
61. Deeks AA, Gibson-Helm ME, Teede HJ. Anxiety and depression in polycystic ovary syndrome: a comprehensive investigation. *FERTILITY AND STERILITY.* 2010;93(7):2421-3.
62. Sezgin H, Hocaoglu C, Guvendag-Guven ES. Disability, psychiatric symptoms, and quality of life in infertile women: a cross-sectional study in Turkey. *Shanghai Archives of Psychiatry.* 2016;28(2):86-94.
63. Wright J, Duchesne C, Sabourin S, Bissonnette F, Benoit J, Girard Y. Psychosocial distress and infertility: men and women respond differently**Supported in part by grant no. 6605-2831-56 from Health and Welfare Canada, Ottawa, Canada; no. RS-1318 from Conseil Québécois de Recherche Sociale, Québec, Québec; and no. 871395 from Fonds de Recherche en Santé du Québec, Montréal, Québec, to Drs. Wright, Duchesne, and Sabourin. *Fertility and Sterility.* 1991;55(1):100-8.
64. Alosaimi FD, Altuwirqi MH, Bukhari M, Abotalib Z, BinSaleh S. Psychiatric disorders among infertile men and women attending three infertility clinics in Riyadh, Saudi Arabia. *Ann Saudi Med.* 2015;35(5):359-67.
65. Herbert DL, Lucke JC, Dobson AJ. Depression: an emotional obstacle to seeking medical advice for infertility. *Fertility and Sterility.* 2010;94(5):1817-21.
66. Upkong D, Orji EO. Mental health of infertile women in Nigeria. *Turk Psikiyatri Dergisi.* 2006;17(4).
67. Atwood JD, Dobkin S. Storm clouds are coming: Ways to help couples reconstruct the crisis of infertility. *Contemporary Family Therapy.* 1992;14(5):385-403.
68. Reed K. The effect of infertility on female sexuality. *Journal of Prenatal & Perinatal Psychology & Health.* 1987;2(1):57.
69. Cook EP. Characteristics of the biopsychosocial crisis of infertility. *Journal of Counseling & Development.* 1987;65(9):465-70.
70. Matthews R, Matthews AM. Infertility and involuntary childlessness: The transition to nonparenthood. *Journal of Marriage and the Family.* 1986;641-9.
71. Chehreh R, Ozgoli G, Abolmaali K, Nasiri M, Mazaheri E. Comparison of the infertility-related stress among couples and its relationship with infertility factors. *International Journal of Women's Health and Reproduction sciences.* 2019;7(3):313-8.
72. Alamin S, Allahyari T, Ghorbani B, Sadeghitabar A, Karami MT. Failure in Identity Building as the Main Challenge of Infertility: A Qualitative Study. *J Reprod Infertil.* 2020;21(1):49-58.
73. Freeman EW, Boxer AS, Rickels K, Tureck R, Mastroianni Jr L. Psychological evaluation and support in a program of in vitro fertilization and embryo transfer. *Fertility and sterility.* 1985;43(1):48-53.
74. Domar AD, Broome A, Zuttermeister PC, Seibel M, Friedman R. The prevalence and predictability of depression in infertile women. *Fertility and sterility.* 1992;58(6):1158-63.

75. Anderheim L, Holter H, Bergh C, Möller A. Does psychological stress affect the outcome of in vitro fertilization? *Hum Reprod.* 2005;20(10):2969-75.
76. Williams KE, Marsh WK, Rasgon NL. Mood disorders and fertility in women: a critical review of the literature and implications for future research. *Human Reproduction Update.* 2007;13(6):607-16.
77. Frederiksen Y, Farver-Vestergaard I, Skovgård NG, Ingerslev HJ, Zachariae R. Efficacy of psychosocial interventions for psychological and pregnancy outcomes in infertile women and men: a systematic review and meta-analysis. *BMJ open.* 2015;5(1):e006592.
78. El Kissi Y, Romdhane AB, Hidar S, Bannour S, Idrissi KA, Khairi H, et al. General psychopathology, anxiety, depression and self-esteem in couples undergoing infertility treatment: a comparative study between men and women. *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 2013;167(2):185-9.
79. Guerra D, Llobera A, Veiga A, Barri PN. Psychiatric morbidity in couples attending a fertility service. *Human reproduction (Oxford, England).* 1998;13(6):1733-6.
80. Ogawa M, Takamatsu K, Horiguchi F. Evaluation of factors associated with the anxiety and depression of female infertility patients. *BioPsychoSocial medicine.* 2011;5:1-5.
81. Sundby J. Infertility in the Gambia: traditional and modern health care. *Patient Education and counseling.* 1997;31(1):29-37.
82. Thiering P, Beaurepaire J, Jones M, Saunders D, Tennant C. Mood state as a predictor of treatment outcome after in vitro fertilization/embryo transfer technology (IVF/ET). *J Psychosom Res.* 1993;37(5):481-91.
83. Noorbala AA, Ramazanzadeh F, Malekafzali H, Abedinia N, Forooshani AR, Shariat M, et al. Effects of a psychological intervention on depression in infertile couples. *International Journal of Gynecology & Obstetrics.* 2008;101(3):248-52.
84. Douki S, Zineb SB, Nacef F, Halbreich U. Women's mental health in the Muslim world: Cultural, religious, and social issues. *Journal of affective disorders.* 2007;102(1-3):177-89.
85. Anvar M, Meshkibaf MH, KOUKABI R. Study of psychiatric disturbance in infertile women. 2006.
86. Ramezanzadeh F, Aghssa MM, Abedinia N, Zayeri F, Khanafshar N, Shariat M, et al. A survey of relationship between anxiety, depression and duration of infertility. *BMC women's health.* 2004;4(1):1-7.
87. Oddens BJ, den Tonkelaar I, Nieuwenhuys H. Psychosocial experiences in women facing fertility problems—a comparative survey. *Human reproduction.* 1999;14(1):255-61.
88. Omani-Samani R, Amini P, Navid B, Sepidarkish M, Maroufizadeh S, Almasi-Hashiani A. Prevalence of sexual dysfunction among infertile women in Iran: A systematic review and meta-analysis. *International Journal of Fertility and Sterility.* 2019;12(4):278-83.
89. Chachamovich JR, Chachamovich E, Ezer H, Fleck MP, Knauth D, Passos EP. Investigating quality of life and health-related quality of life in infertility: a systematic review. *Journal of Psychosomatic Obstetrics & Gynecology.* 2010;31(2):101-10.

90. Keramat A, Masoomi SZ, Mousavi SA, Poorolajal J, Shobeiri F, Hazavhei SMM. Quality of life and its related factors in infertile couples. *Journal of research in health sciences*. 2013;14(1):57-64.
91. Rashidi B, Montazeri A, Ramezanzadeh F, Shariat M, Abedinia N, Ashrafi M. Health-related quality of life in infertile couples receiving IVF or ICSI treatment. *BMC health services research*. 2008;8(1):1-6.
92. Schaller MA, Griesinger G, Banz-Jansen C. Women show a higher level of anxiety during IVF treatment than men and hold different concerns: a cohort study. *Archives of gynecology and obstetrics*. 2016;293:1137-45.
93. Lawson AK, Klock SC, Pavone ME, Hirshfeld-Cytron J, Smith KN, Kazer RR. Prospective study of depression and anxiety in female fertility preservation and infertility patients. *Fertility and sterility*. 2014;102(5):1377-84.
94. Allan HT. The anxiety of infertility: the role of the nurses in the fertility clinic. *Human Fertility*. 2013;16(1):17-21.
95. Kraaij V, Garnefski N, Schroevers MJ, Weijmer J, Helmerhorst F. Cognitive coping, goal adjustment, and depressive and anxiety symptoms in people undergoing infertility treatment: a prospective study. *Journal of Health Psychology*. 2010;15(6):876-86.
96. Chen T-H, Chang S-P, Tsai C-F, Juang K-D. Prevalence of depressive and anxiety disorders in an assisted reproductive technique clinic. *Human reproduction*. 2004;19(10):2313-8.
97. Lu Y, Yang L, Lu G. Mental status and personality of infertile women. *Zhonghua Fu Chan Ke Za Zhi*. 1995;30(1):34-7.
98. Lok IH, Lee DTS, Cheung LP, Chung WS, Lo WK, Haines CJ. Psychiatric morbidity amongst infertile Chinese women undergoing treatment with assisted reproductive technology and the impact of treatment failure. *Gynecologic and obstetric investigation*. 2002;53(4):195-9.
99. Nana P, Wandji J, Fomulu J, Mbu R, Leke R, Woubinwou M. Aspects Psycho-Sociaux chez Patients Infertiles à la Maternité Principale de l'Hôpital Central de Yaoundé, Cameroun. *Clinics in mother and child health*. 2011;8(1).
100. Patel A, Sharma P, Narayan P, Binu V, Dinesh N, Pai PJ. Prevalence and predictors of infertility-specific stress in women diagnosed with primary infertility: A clinic-based study. *Journal of human reproductive sciences*. 2016;9(1):28.
101. Busari A, Agokei SP. Psychosocial factors as predictors of infertility related stress among female secondary school teachers in Ibadan North Local Government Area Oyo State. *Nigeria*; 2017.
102. Volgsten H, Skoog Svanberg A, Ekselius L, Lundkvist Ö, Sundström Poromaa I. Prevalence of psychiatric disorders in infertile women and men undergoing in vitro fertilization treatment. *Human Reproduction*. 2008;23(9):2056-63.
103. Luk BH-K, Loke AY. The impact of infertility on the psychological well-being, marital relationships, sexual relationships, and quality of life of couples: A systematic review. *Journal of sex & marital therapy*. 2015;41(6):610-25.
104. Dooley M, Dineen T, Sarma K, Nolan A. The psychological impact of infertility and fertility treatment on the male partner. *Human fertility*. 2014;17(3):203-9.