

Trends in Antibiotic Use in a UK Secondary Care Prior to and During the COVID-19 Pandemic: A Cross-Sectional Retrospective Study



Rasha Abdelsalam Elshenawy

Department of Pharmacy, School of Life and Medical Sciences, University of Hertfordshire, UK

Conflict of Interest



I have no conflict of interest.

Objectives



- Provide an overview of the research background, explaining its significance and the problem it addresses.
- Describe the methodology, including how data was collected and analysed.
- Highlight key findings and their importance in answering the research questions.
- Explain how the findings impact contemporary pharmacy practice and research.
- Discuss future research directions and potential applications of the findings.

Research background



- Antimicrobial resistance (AMR) is a rapidly escalating global health challenge will cause 39 million deaths between 2025 and 2050.
- To address this, the UK Government developed a 5-year action plan, 'Confronting Antimicrobial Resistance 2024 to 2029,' aimed at optimising antimicrobial use.
- The World Health Organisation (WHO) promotes antimicrobial stewardship (AMS) and developed the AWaRe classification system to guide global AMS implementation. The defined daily dose (DDD) standardises antibiotic use comparisons.
- The COVID-19 pandemic significantly impacted health services, particularly in secondary care, potentially exacerbating AMR due to increased antibiotic use.

Aim of the research study

To evaluate changes in antibiotic use in a UK secondary care setting following the onset of the COVID-19 pandemic in 2020 compared to 2019.



Methodology
























- This cross-sectional retrospective study evaluated antibiotic use among adults at an NHS Foundation Trust in England.
- It focused on patients with respiratory tract infections (RTIs) and pneumonia in 2019 and 2020.
- Data from 640 patient records were extracted using a WHO AWaRe-based tool.
- Study included adults, immunocompromised, pregnant women, and RTI patients.
- Antibiotic use measured by Defined Daily Dose (DDD).
- Registered under ISRCTN 14825813 with HRA ethical approval.
- Public and patient involvement was ensured through review by the Citizens Senate.

Key findings



- The results show a notable increase in the Defined Daily Dose (DDD) of several antibiotics during the COVID-19 pandemic.
- In the "Access" category, amoxicillin usage rose from 1.33 DDD pre-pandemic to 4.33 DDD in 2020. Co-amoxiclav remained nearly constant at around 15 DDD but showed a marked increase compared to other antibiotics.
- In the "Watch" category, azithromycin increased from 3 to 9 DDD, and clarithromycin from 32 to 35.5 DDD. Levofloxacin usage doubled from 13 to 26 DDD.
- The "Reserve" category showed no significant increase, indicating stable consumption.

Trends in antibiotic use prior to pandemic and during the COVID-19 pandemic (2019 and 2020)

| Antibiotic Name | Antibiotic Category | Antibiotics DDD in 2019 | Antibiotics DDD in 2020 | Trend |
|-------------------------------|---------------------|-------------------------|-------------------------|---|
| Amoxicillin | Access | 1.33 | 4.33 |  |
| Co-amoxiclav | Access | 15.9 | 15.2 |  |
| Benzylopenicillin | Access | 0.45 | 0.15 |  |
| Doxycycline | Access | 6 | 2 |  |
| Flucloxacillin | Access | 1.13 | 1.13 |  |
| Gentamicin | Access | 0 | 6.67 |  |
| Metronidazole | Access | 1.8 | 0.4 |  |
| Sulfamethoxazole/trimethoprim | Access | 0.25 | 1.5 |  |
| Cephalexin | Access | 0 | 0.5 |  |
| Azithromycin | Watch | 3 | 9 |  |
| Ceftazidime | Watch | 1 | 0 |  |
| Ceftriaxone | Watch | 0.5 | 1 |  |
| Ciprofloxacin | Watch | 7 | 9 |  |
| Clarithromycin | Watch | 32 | 35.5 |  |
| Levofloxacin | Watch | 13 | 28 |  |
| Meropenem | Watch | 0.67 | 2 |  |
| Piperacillin/tazobactam | Watch | 0.32 | 0.32 |  |
| Teicoplanin | Watch | 1 | 1 |  |
| Vancomycin | Watch | 1.5 | 1 |  |
| Aztreonam | Reserve | 0 | 0.5 |  |
| Cefazidime/Azobactam | Reserve | 1.67 | 0 |  |
| Linezolid | Reserve | 1.05 | 0.3 | |

Relevance & impact of findings to contemporary pharmacy practice/research

- The findings highlight the urgent need for robust antimicrobial stewardship to tackle changes in prescribing behaviours during the COVID-19 pandemic.
- This is particularly important due to the increased use of high-risk antibiotics, such as macrolides and levofloxacin.
- These insights emphasise the crucial role of monitoring and optimising antibiotic use in secondary care to improve patient safety and treatment quality.
- They are highly relevant to current pharmacy practice and research, particularly in the fight against antimicrobial resistance.

Future of the presented research



- It is important to provide sustainable and resilient measures to antimicrobial stewardship for long-term effectiveness.
- Future research is essential to ensure the effective implementation of antimicrobial stewardship and appropriate antibiotic prescribing.
- Expanding the antibiotic dashboard to broader healthcare settings will enhance monitoring.
- Integrating real-time data analytics can improve stewardship interventions.
- Predictive models should be developed to anticipate prescribing trends during crises.
- The dashboard's effectiveness in reducing inappropriate antibiotic use across various settings needs to be assessed.

References



- Abdelsalam Elshenawy R, Umaru N, Aslanpour Z. Impact of COVID-19 on “Start Smart, Then Focus” Antimicrobial Stewardship at One NHS Foundation Trust in England Prior to and during the Pandemic. COVID [Internet]. 2024 Jan 1;4(1):102–16. Available from: <https://www.mdpi.com/2673-8112/4/1/10>.
- Department of Health and Social Care. UK 5-year action plan for antimicrobial resistance 2024 to 2029 [Internet]. GOV.UK. 2024. Available from: <https://www.gov.uk/government/publications/uk-5-year-action-plan-for-antimicrobial-resistance-2024-to-2029>.
- GBD 2021 Antimicrobial Resistance Collaborators (2024). Global burden of antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. The Lancet. doi:[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(24\)01867-1/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)01867-1/fulltext).
- WHO. The WHO AWaRe (Access, Watch, Reserve) antibiotic book - Infographics [Internet]. www.who.int. 2023. Available from: <https://www.who.int/publications/i/item/WHO-MHP-HPS-EML-2022.02>.

THANK YOU!

Rasha Abdelsalam Elshenawy

r.elshenawy@herts.ac.uk

Twitter: [Salam_Rasha](#)

Acknowledge

Prof Zoe Aslanpour

Dr Nkiruka Umaru