

## Correction

# Correction: Filippou et al. Transcriptomic Analysis Reveals Molecular Mechanisms Underpinning Mycovirus-Mediated Hypervirulence in *Beauveria bassiana* Infecting *Tenebrio molitor*. *J. Fungi* 2025, 11, 63

Charalampos Filippou <sup>1,2,3,\*</sup>, Robert H. A. Coutts <sup>2</sup>, Ioly Kotta-Loizou <sup>2,4</sup>, Sam El-Kamand <sup>3</sup> and Alexie Papanicolaou <sup>3,\*</sup>

<sup>1</sup> Department of Medicine, School of Medicine, European University Cyprus, 2404 Nicosia, Cyprus

<sup>2</sup> Department of Clinical, Pharmaceutical and Biological Science, School of Life and Medical Sciences, University of Hertfordshire, Hatfield AL10 9AB, UK; r.coutts@herts.ac.uk (R.H.A.C.); i.kotta-loizou2@herts.ac.uk (I.K.-L.)

<sup>3</sup> Hawkesbury Institute for the Environment, Western Sydney University, Richmond, NSW 2753, Australia; sam.el-kamand@westernsydney.edu.au

<sup>4</sup> Department of Life Sciences, Faculty of Natural Sciences, Imperial College London, London SW7 2AZ, UK

\* Correspondence: c.filippou@euc.ac.cy (C.F.); a.papanicolaou@westernsydney.edu.au (A.P.)

## References

There was an error in the original publication [1]. We failed to mention the origin of the strain used in our publication. This correction requires the addition of the following reference:

27. Quesada-Moraga, E.; Vey, A. Intra-specific variation in virulence and in vitro production of macromolecular toxins active against locust among *Beauveria bassiana* strains and effects of in vivo and in vitro passage on these factors. *Biocontrol Sci. Technol.* **2003**, *13*, 323–340.

A correction has been made to Section 2 Materials and Methods, 2.1. Maintenance of Fungal Isolates, Paragraph 1:

“This strain belongs to the culture collection of the Department of Agronomy of the University of Cordoba (Spain) and was originally isolated from infected specimens of the Moroccan locust *Locusta marocanica* (Thunberg) in the breeding area of La Serena in Badajoz (Spain). This strain has been deposited in the Spanish Type Culture Collection located at the University of Valencia under accession number CECT 20376 [27].”

## Corrected Paragraph

### 2. Materials and Methods

#### 2.1. Maintenance of Fungal Isolates

This study relies on a *B. bassiana* isolate named “EABb 92/11-Dm” which is infected with two mycoviruses, *Beauveria bassiana* non-segmented virus (BbNV) 1 and *Beauveria bassiana* polymycovirus (BbPmV) 1. This strain belongs to the culture collection of the Department of Agronomy of the University of Cordoba (Spain) and was originally isolated from infected specimens of the Moroccan locust *Locusta marocanica* (Thunberg) in the breeding area of La Serena in Badajoz (Spain). This strain has been deposited in the Spanish Type Culture Collection located at the University of Valencia under accession number CECT 20376 [27]. The virus-infected and virus-free isogenic lines used were generated as reported previously [17] and are herein designated BbVI and BbVF, respectively. Two virus-free,



Academic Editor: Chengshu Wang

Received: 15 September 2025

Accepted: 14 October 2025

Published: 30 October 2025

**Citation:** Filippou, C.; Coutts, R.H.A.; Kotta-Loizou, I.; El-Kamand, S.; Papanicolaou, A. Correction: Filippou et al. Transcriptomic Analysis Reveals Molecular Mechanisms Underpinning Mycovirus-Mediated Hypervirulence in *Beauveria bassiana* Infecting *Tenebrio molitor*. *J. Fungi* 2025, *11*, 63. *J. Fungi* 2025, *11*, 782. <https://doi.org/10.3390/jof11110782>

**Copyright:** © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

commercial isolates, ATCC 74040 and GHA, were also used. All isolates were maintained on potato dextrose agar (PDA; Sigma-Aldrich, St. Louis, MO, USA) or Sabouraud dextrose agar (SBA; Sigma-Aldrich, St. Louis, MO, USA) at 25 °C and 80% relative humidity [28]. A cocktail of antibiotics (ampicillin, kanamycin, and streptomycin, each at a final concentration of 100 µg/mL) was used during the cultivation of all strains to prevent bacterial contamination. The use of antibiotics ensured that bacterial contaminants did not interfere with the growth and observations of the fungal strains, thereby maintaining the integrity and accuracy of the experimental results.

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

## Reference

1. Filippou, C.; Coutts, R.H.A.; Kotta-Loizou, I.; El-Kamand, S.; Papanicolaou, A. Transcriptomic Analysis Reveals Molecular Mechanisms Underpinning Mycovirus-Mediated Hypervirulence in *Beauveria bassiana* Infecting *Tenebrio molitor*. *J. Fungi* **2025**, *11*, 63. [\[CrossRef\]](#) [\[PubMed\]](#)

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.