

# 1 Usefulness of the NULL-PLEASE Score to predict survival in out-of- 2 hospital cardiac arrest

3 Dr Ying X. GUE MBChB MRCP<sup>1,2</sup>, Dr Max SAYERS MBBS MRCP<sup>3</sup>, Dr Benjamin T. WHITBY MBBS<sup>4</sup>, Dr  
4 Rahim KANJI MBBS MRCP<sup>2</sup>, Dr Krishma ADATIA MBBS<sup>2</sup>, Dr Robert SMITH MD<sup>3</sup>, Dr William R. DAVIES  
5 MBChB PhD<sup>5</sup>, Dr Aris PERPEROGLU PhD<sup>6</sup>, Prof. Tatjana S. POTPARA MD PhD<sup>7</sup>, Prof. Gregory Y.H. LIP  
6 MD FRCP DFM FACC FESC FEHRA<sup>8\*</sup>, Prof. Diana A. GOROG MD PhD FRCP FESC<sup>1,2,9\*</sup>

- 7
- 8 1. University of Hertfordshire, United Kingdom
- 9
- 10 2. East and North Hertfordshire NHS Trust, Hertfordshire, United Kingdom
- 11
- 12 3. Royal Brompton & Harefield NHS Trust, Harefield, United Kingdom
- 13
- 14 4. University of Cambridge, Cambridge, United Kingdom
- 15
- 16 5. Royal Papworth Hospital NHS Foundation Trust, Cambridge, United Kingdom
- 17
- 18 6. University of Essex, United Kingdom
- 19
- 20 7. Clinical Centre of Serbia & School of Medicine, Belgrade University, Serbia, Belgrade
- 21
- 22 8. Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest  
23 Hospital, Liverpool, United Kingdom; and Aalborg Thrombosis Research Unit, Department of Clinical  
24 Medicine, Aalborg University, Aalborg, Denmark
- 25
- 26 9. National Heart and Lung Institute, Imperial College, London, United Kingdom

27  
28 \* Joint senior authors.

29 Funding: None

30 Declarations of interest: None

31 All authors had access to data and a role in writing the manuscript

32 Running head: NULL-PLEASE score mortality prediction in out-of-hospital cardiac arrest

33 Keywords: Cardiac arrest, prognosis, survival, outcome, risk-score

34  
35 Word count: 2996

36  
37 Correspondence to:  
38 Prof. Diana A Gorog  
39 National Heart & Lung Institute  
40 Imperial College  
41 Dovehouse Street  
42 London SW3 6LY  
43 United Kingdom  
44 Email: [d.gorog@imperial.ac.uk](mailto:d.gorog@imperial.ac.uk)

45 **Abstract**

46 **Purpose:** Out-of-hospital cardiac arrest (OHCA) carries a very high mortality even after  
47 successful cardiopulmonary resuscitation. Currently, information given to relatives regarding  
48 prognosis following resuscitation is often emotive and subjective, and varies with clinician  
49 experience. We aimed to validate the NULL-PLEASE score to predict survival following  
50 OHCA.

51 **Methods:** A multicentre cohort study was conducted, with retrospective and prospective  
52 validation in consecutive unselected patients presenting with OHCA. The NULL-PLEASE  
53 score was calculated by attributing points to the following variables: Non-shockable initial  
54 rhythm, Unwitnessed arrest, Long low-flow period, Long no-flow period, pH<7.2,  
55 Lactate>7.0 mmol/l, End-stage renal failure, Age ≥85 years, Still resuscitation and Extra-  
56 cardiac cause. The primary outcome was in-hospital death.

57 **Results:** We assessed 700 patients admitted with OHCA, of whom 47% survived to  
58 discharge. In 300 patients we performed a retrospective validation, followed by prospective  
59 validation in 400 patients. The NULL-PLEASE score was lower in patients who survived  
60 compared to those who died (0 [IQR 0-1] vs. 4 [IQR 2-4], p<0.0005) and strongly predictive  
61 of in-hospital death (c-statistic 0.874, 95% confidence interval [CI] 0.848-0.899). Patients  
62 with a score ≥3 had a 24-fold increased risk of death (OR 23.6; 95%CI 14.840-37.5,  
63 p<0.0005) compared to those with lower scores. A score ≥3 has a 91% positive predictive  
64 value for in-hospital death, whilst a score <3 predicts a 71% chance of survival.

65 **Conclusion:** The easy-to-use NULL-PLEASE score predicts in-hospital mortality with high  
66 specificity and can help clinicians explain the prognosis to relatives in an easy-to-understand,  
67 objective fashion, to realistically prepare them for the future.

68

69 *Word count: 250*

70 **Abbreviations**

71

72 CI Confidence interval

73 IQR Interquartile range

74 NPV Negative predictive value

75 OHCA Out-of-hospital cardiac arrest

76 OR Odds ratio

77 PPV Positive predictive value

78 ROC Receiver operator characteristic curve

79

## 80 **Introduction**

81 Out-of-hospital cardiac arrest (OHCA) affects 84 per 100,000 population.<sup>1</sup> In ~28% of  
82 individuals, there is return of spontaneous circulation and of these, only 10% survive to 30  
83 days or hospital discharge.<sup>1</sup>

84 The post-cardiac arrest syndrome, comprising of possible brain injury, myocardial  
85 dysfunction, systemic ischaemia/reperfusion response, and the persistent precipitating  
86 pathology, often requires resource-intensive monitoring and lengthy treatment in the  
87 intensive care unit.<sup>2</sup> Despite the numerous ethical issues which may be involved,<sup>3</sup> an accurate  
88 prognostic assessment early in the pathway may be helpful for medical teams to help decision  
89 making, to guide families, and to allow allocation of resources to those that are likely to  
90 benefit most, in an objective fashion.

91 Such a scoring system should have high sensitivity (to predict patients with poor prognosis)  
92 and high specificity (to ensure all patients with potentially good outcomes are treated).<sup>4,5</sup>  
93 Several scores of varying complexity and limited practical application have been developed,  
94 and there is currently no recommended simple scoring system for routine clinical use. Yet,  
95 we believe that both healthcare professionals and families of patients would wish to know,  
96 following OHCA, the likelihood of an individual surviving to hospital discharge. Such a  
97 scoring system may be helpful to healthcare professionals and relatives/friends to provide  
98 objective, realistic and non-emotive prognostification at such a crucial time.

99 Currently available risk scores to predict mortality have important limitations. The OHCA  
100 Score integrates arrest-related and biochemical variables without patient-specific  
101 characteristics,<sup>6</sup> with a c-statistic of 0.88. However, its main limitations is that it is very  
102 difficult to calculate, including complex weighting of characteristics and calculation of the

103 natural logarithm of 3 characteristics, making it unpracticable, and it has only been assessed  
104 in small cohorts.<sup>6</sup> The ACLS score, developed >30y ago, is difficult to calculate and has  
105 relatively poor performance (area-under-the-ROC-curve, AUC 0.786).<sup>7</sup> Similarly, the  
106 Graphic Model is very difficult to compute, requires data that are frequently not available  
107 (such as minutes to start of CPR or defibrillation) and has not been externally validated.<sup>8</sup> The  
108 Prediction Tool is also complex and cumbersome to calculate, and not externally validated.<sup>9</sup>  
109 Some scores have only been evaluated in small cohorts,<sup>6,10</sup> some not prospectively  
110 assessed,<sup>7,8,10</sup> some not externally or prospectively validated,<sup>8-12</sup> and some only predict  
111 survival to 1 month, but not in the hospital setting.<sup>6,9-11</sup> There is therefore an urgent, unmet  
112 need for a simple, easy-to-use clinical scoring system to predict survival to hospital  
113 discharge, with high sensitivity and specificity.

114 The NULL-PLEASE score is a relatively new “futility” score to help identify patients who  
115 are unlikely to survive following OHCA.<sup>13</sup> The score has only been validated to predict death  
116 in the emergency room, with a c-statistic of 0.658.<sup>14</sup> Its usefulness for predicting survival in  
117 hospital has not been assessed.

118 It was our aim to provide independent external validation of the NULL-PLEASE score for  
119 prediction of in-hospital survival, in a large cohort of patients with OHCA.

120

121 **Methods**

122 We performed an external validation of the NULL-PLEASE score in an all-comers  
123 population of consecutive patients presenting with OHCA to three large NHS Trusts in  
124 England (East and North Hertfordshire NHS Trust, Royal Brompton and Harefield Hospitals  
125 NHS Trust and Royal Papworth Hospital, Cambridge) from September 2015 to December  
126 2018, as part of an approved service evaluation with permission from local R&D boards.

127

128 ***NULL-PLEASE Score***

129 The NULL-PLEASE score assigns 2 points to each of the initial arrest characteristics  
130 (Nonshockable rhythm, Unwitnessed arrest, Long no-flow or Long low-flow period) and 1  
131 point to each patient characteristic (blood PH <7.2, Lactate >7.0 mmol/L, End-stage kidney  
132 disease on dialysis, Age ≥85 years, Still resuscitation, and Extra-cardiac cause). Definitions  
133 of individual components of the score are shown in Table 1. As a number of patients did not  
134 have lactate or pH measured on arrival, the performance of a modified version of the scoring  
135 system excluding these variables, namely the NULL-EASE score, was also assessed.

136

137 ***Data collection***

138 Demographics, descriptive data pertaining to the arrest, initial blood results including pH and  
139 lactate, cause of arrest (or presumed cause) and length of hospital stay were documented by  
140 clinicians independent of the research team.

141

142 ***Outcome***

143 The primary outcome was in-hospital death or survival to discharge from hospital. The  
144 secondary outcome was length of stay.

145

146 *Statistical analysis*

147 Categorical variables were summarised as proportion (number and percentage) and  
148 continuous variables as median with interquartile range (IQR). The association of the NULL-  
149 PLEASE score components with the primary outcome was examined using univariate logistic  
150 regression analysis. Odds ratios (ORs) with 95% confidence interval (CI) and p-values were  
151 obtained for each component and the score as a whole. The predictive ability of the NULL-  
152 PLEASE score for the primary outcome was tested using AUC analysis and the c-statistic  
153 reported. The same analysis was performed for patients in whom only the NULL-EASE score  
154 was available.

155 Bootstrap re-sampling<sup>15</sup> was used to assess the predictive ability of the score for new data.

156 This has two steps: at the training step, a part of the data is used to fit a logistic regression  
157 model, and at the testing step, the estimates of the logistic regression model are used to  
158 predict how patients not included in the training set would be classified. The process repeats  
159 a thousand times.

160 A subgroup analysis was performed in patients who had return of spontaneous circulation  
161 following the initial arrest and in patients with myocardial infarction as the presumed cause  
162 of arrest. Significance was taken as  $<0.05$ . Statistical analyses were performed using Stata 15  
163 software (StataCorp, College Station, Texas, USA).

164 **Results**

165 A total of 700 patients were included, 300 in the retrospective and 400 in the prospective  
166 validation cohorts. Of the 700 patients, 332 (47%) survived to hospital discharge.  
167 Blood pH results were unavailable in 196 patients and lactate was unavailable in 232 patients.  
168 The causes of OHCA were myocardial infarction (n=454), pulmonary embolism (n=20),  
169 cerebrovascular accident (n=3), bleeding (n=6), trauma (n=9), other causes (n=117) including  
170 sepsis, electrolyte disturbances, and 91 unknown. The median length of stay was 5 days [IQR  
171 2-10].

172  
173 Baseline characteristics of the 300 patients in the retrospective cohort are shown in Table 2.  
174 The NULL-PLEASE score was significantly lower in survivors compared to those who died  
175 (0[IQR 0-0] vs. 3[IQR 2-5],  $p < 0.0005$ ). On univariate logistic regression analysis (Table 2),  
176 most components of the score were individually significantly associated with in-hospital  
177 mortality, except for gender, end-stage renal failure, extra-cardiac cause and age >85 years,  
178 which were under-represented in this cohort. The NULL-PLEASE score was a strong  
179 predictor of in-hospital death (c-statistic 0.851, 95%CI 0.808-0.895). We chose a NULL-  
180 PLEASE score  $\geq 3$  as the optimal cut-point to predict mortality, with sensitivity 50.4% and  
181 specificity 94.4% (Figure 1A), with a positive predictive value (PPV) of 86.1% for in-  
182 hospital death and negative predictive value (NPV) of 73.6% for survival. Although a score  
183  $\geq 2$  had the best combined sensitivity (78.9%) and specificity (84.2%), the cut-point of 3 was  
184 chosen to improve specificity, to ensure almost all patients with potentially good outcomes  
185 are treated, whilst preserving reasonable sensitivity.

186  
187 Baseline characteristics of the 400 patients included in the prospective validation cohort are  
188 shown in Table 3. The NULL-PLEASE score was significantly lower in those surviving to

189 discharge compared to those who died (0[IQR 0-1] vs. 4[IQR 2-6],  $p<0.0005$ ). On univariate  
190 logistic regression analysis (Table 3), all components of the score were significantly  
191 associated with mortality, except for gender and end-stage renal failure, which were under-  
192 represented. The score was confirmed to be a strong predictor of in-hospital death (c-statistic  
193 0.8797, 95%CI 0.8471-0.912) in this prospective validation cohort. A NULL-PLEASE score  
194  $\geq 3$  had sensitivity 73.5% and specificity 90.3%, with a PPV of 92.3% for in-hospital  
195 mortality and NPV of 68.3% (Table 4).

196

197 Combining the retrospective and the prospective cohorts, the odds of in-hospital death  
198 increased with increasing NULL-PLEASE score (Table 4). Patients with a score  $\geq 3$  had a 24-  
199 fold increased risk of in-hospital death (OR 23.6; 95%CI 14.87-37.40,  $p<0.0005$ ) compared  
200 to patients with lower scores, with PPV 90.6% and NPV 70.9%. Using logistic regression, a  
201 NULL-PLEASE score of 3 was associated with 75% likelihood of death (Figure 1B). Results  
202 of bootstrap resampling indicated that the average specificity and sensitivity of a model with  
203 NULL-PLEASE score  $\geq 3$  when predicting out-of-sample observations was 90.8% and  
204 70.7%, respectively (Table 5).

205

#### 206 *Subgroup of patients with OHCA secondary to myocardial infarction*

207 Myocardial infarction was the cause of death in 454 patients and 249 (55%) survived to  
208 discharge. The score performed well in this group (AUC 0.836, 95%CI 0.80-0.87). Amongst  
209 these patients, those with a NULL-PLEASE score  $\geq 3$  had a 19-times higher risk of death (OR  
210 19.6; 95%CI 10.3-37.1,  $p<0.0005$ ) compared to those with lower scores.

211

#### 212 *The modified NULL-EASE score*

213 Since a number of patients did not have lactate or pH measured on arrival, the usefulness of  
214 the modified NULL-EASE score, was also assessed.

215 In the retrospective cohort, the NULL-EASE score was a strong predictor of death, with  
216 AUC 0.819 (95%CI 0.773-0.866). A score  $\geq 3$  had a sensitivity of 39.84% and specificity of  
217 96.05%. Similarly, in the prospective cohort, the NULL-EASE score showed an AUC 0.860  
218 (95%CI 0.826-0.894). A score  $\geq 3$  had sensitivity of 66.12% and specificity of 90.32%.

219 Combining the retrospective and prospective cohorts, the NULL-EASE score remained a  
220 strong predictor of death (AUC 0.849; 95%CI 0.822-0.876), with a score  $\geq 3$  having  
221 sensitivity of 57.34% and specificity of 93.37%, PPV 90.6% and NPV 66.4%

222

#### 223 *NULL-PLEASE score and length of stay*

224 In patients who achieved return of spontaneous circulation following the initial arrest, the  
225 median length of stay was 6 days (IQR 3-12). Among these, length of stay was significantly  
226 longer in patients who survived compared to those who died in hospital (9[IQR 4-16] vs.  
227 4[IQR 2-7] days,  $p < 0.00005$ ). Using Spearman rank correlation, the NULL-PLEASE score  
228 showed weak positive correlation with length of stay in survivors ( $r = 0.248$ ,  $p < 0.0005$ ) and  
229 moderate negative correlation in patients who died ( $r = -0.472$ ,  $p < 0.0005$ ).

230

231 **Discussion**

232

233 In this independent external validation in a contemporary cohort of OHCA patients, we show  
234 that the NULL-PLEASE score is a strong predictor of in-hospital death, with high sensitivity  
235 and specificity. Individuals with a score  $\geq 3$  had a 24-fold increased risk of death compared to  
236 those with a score of 0-2. A score  $\geq 3$  had a 90.6% PPV for in-hospital death, whilst the NPV  
237 indicates that a patient with a score  $< 3$  has 70.9% chance of survival. Such prognostic  
238 information can be very useful for both healthcare professionals and relatives, can be easily  
239 and quickly calculated, and easily understood by lay individuals.

240 Our study provides the most compulsive data yet in support of a risk score to predict survival  
241 in OHCA, which is extremely easy-to-use, yet has high sensitivity and specificity, high NPV  
242 and PPV, and which has been externally validated, both retrospectively and prospectively, in  
243 a very large cohort. With the utilisation of both arrest- and patient-specific characteristics, the  
244 NULL-PLEASE score includes vital features associated with adverse outcome.<sup>16</sup>

245 Importantly, no risk score calculator will be 100% accurate. Experienced clinicians will  
246 recognise that not infrequently, patients defy expectations and those thought to have no  
247 chance have recovered, whilst some of those predicted to do well, have succumbed.

248 Therefore, such a scoring system can at best serve as an adjunct to decision-making and  
249 cannot be used to make decisions on withdrawal of life-supporting treatment in individual  
250 patients. It can, however, be used to guide and explain prognosis to relatives who may find  
251 that being quoted an objective survival rate based on the score may help better prepare them  
252 for the future. Currently, in our experience, information given to relatives is often varied,  
253 being frequently both emotive and subjective (for example, wishing to convey hope even in  
254 perhaps hopeless scenarios, or predicting gloom to avoid unrealistic expectations by relatives

255 and to prepare them for the worst), and varying with the seniority and experience of the  
256 clinician.

257

258 The great strength of the NULL-PLEASE score is not only its strong prognostic value, but its  
259 simplicity and ease-of-use. It can be calculated on the spot and is easy to interpret. In  
260 comparison, both the OHCA and CAHP scores are difficult to calculate, needing advanced  
261 calculator functions, or nomograms, and are neither easy to calculate, nor clinically-friendly.  
262 Our results support and extend the findings of the initial validation of the NULL-PLEASE  
263 score for death in the emergency room in a small cohort,<sup>14</sup> to now predict survival to hospital  
264 discharge, in a large independent cohort, with subsequent validation. Since some 55% of  
265 OHCAs are attributable to a cardiac cause,<sup>17</sup> the strong performance of the score in this  
266 subgroup is highly pertinent. The individual variables in the univariate analysis were highly  
267 predictive of outcome, with the exception of variables that were under-represented and thus  
268 could not be assessed.

269

270 A NULL-PLEASE score  $\geq 3$  had a specificity of 92.5%, ensuring most patients with  
271 potentially good outcomes are not disadvantaged, with a PPV for in-hospital death of 90.6%  
272 with sensitivity 65.8%. In comparison to other scoring systems, an OHCA score<sup>6</sup>  $\geq 32.5$  has  
273 specificity of only 85% and PPV 96%, sensitivity 46% and specificity 96%. However, the  
274 NULL-PLEASE score achieves superior predictive value, and is much easier-to-use.

275

276 Although routine blood gas analysis is recommended in patients with OHCA, it is frequently  
277 not performed upon arrival, due to the pressures of manpower or time and competing  
278 priorities in an emergency situation. Our sensitivity analysis using the modified NULL-EASE  
279 score showed a PPV of 90.6% for a score  $\geq 3$ , similar to that of the NULL-PLEASE score,

280 although sensitivity was lower at 57.3% and NPV only 66.4%. This highlights the importance  
281 of measuring pH and lactate upon arrival to optimise the performance of the score.

282

283 Although both populations consisted of consecutive all-comers, the retrospective and  
284 prospective cohorts differ in some demographic aspects, for example extracardiac cause of  
285 arrest 1% vs. 30%, and non-shockable rhythm 11% vs. 36%, respectively, with associated  
286 difference in mortality (41% and 61%, respectively). These differences, are almost certainly  
287 due to selection bias in the retrospective cohort, which likely unintentionally excluded  
288 patients who may have died very shortly after admission as these cases may not be logged on  
289 databases, as we observed when collecting prospective data. However, this underscores the  
290 importance of prospective validation of any risk scoring system and specifically the strength  
291 of the prospective validation here, which included more patients with extracardiac arrest and  
292 with non-shockable rhythm, showing the score to be applicable to different clinical  
293 presentations.

294

295 The length of stay in our cohort is short compared to a recent UK cohort managed on the  
296 intensive care unit,<sup>19</sup> reporting a median stay of 12 days. This is likely due to the unselected  
297 nature of our patients, whereas Petrie *et al.* reviewed only patients admitted to the intensive  
298 care unit. Even though our median stay is shorter, it still reflects the very significant health  
299 economic burden that patients with OHCA place on healthcare systems. When resources are  
300 limited, the appropriate allocation of resources to patients that are most likely to survive is  
301 essential. We believe our score may be helpful for identifying likely survivors, when  
302 optimizing use of finite healthcare resources, although this can only serve as a rough guide.  
303 New costly interventions are increasingly subjected to cost-effectiveness evaluations, which

304 will require quantification of the potential benefit, for example the number of additional lives  
305 saved. Our score may also be helpful for this purpose.

306

### 307 **Limitations**

308 There is inherent bias in the studied population, since these individuals already survived to  
309 reach hospital, and we excluded those who died pre-admission. For the variable ‘Still  
310 resuscitation’, meaning ongoing CPR on arrival to hospital, this is very dependent on the  
311 particular healthcare system. We are aware that in some places, CPR is almost always  
312 continued to hospital arrival (meaning almost every OHCA case will have ongoing CPR on  
313 arrival), whereas other systems have prehospital physicians or paramedics who can terminate  
314 resuscitation on scene (meaning that only patients with the highest chance of survival are  
315 transported to hospital with ongoing CPR, resulting in selection bias). The score incorporates  
316 aetiology, namely “E- extra cardiac cause”, which in practical terms is frequently not  
317 available. In most patients myocardial infarction was the cause of OHCA, and whether the  
318 score is equally applicable to patients with other causes of OHCA is unclear. Furthermore,  
319 the cause of death was presumed in many cases, without definitive tests, especially in those  
320 who died shortly after admission, since in the UK, post-mortems are not routinely performed,  
321 with cause of death determined by clinicians based on likelihood, given presentation and  
322 comorbidities. Details pertaining to the circumstances of the OHCA and resuscitation are  
323 based on documentation and approximation during or post-event, which may be commonly  
324 inaccurate.<sup>20,21</sup> In the score, ‘Long no-flow period’ is defined as no bystander CPR prior to  
325 arrival of emergency medical services. However, there are no defined time periods for the no-  
326 flow period, it could therefore range from a few to many minutes. Further, although most  
327 components of the NULL-PLEASE score performed well individually, end-stage kidney

328 disease was under-represented in our cohort and so conclusions cannot be drawn about the  
329 usefulness of this particular component of the score.

330 An important limitation is that this risk score does not provide information on neurological  
331 status on discharge, although there are several available scoring systems to assess the  
332 likelihood of good functional recovery on the intensive care unit.<sup>21,22</sup> Lactate and pH were not  
333 always available, and the score appears to perform less well without inclusion of these. On  
334 the other hand, this reflects real-life scenarios where these measurements are not always  
335 available at the time of decision making, highlighting the relative usefulness of the NULL-  
336 EASE score. Finally, the score is predictive of outcome in the average patient, not the  
337 individual patient. Furthermore, the organization of emergency medical services varies across  
338 countries, and our score may need to be calibrated for each specific system.

### 339 **Conclusion**

340 The NULL-PLEASE score is an easy-to-use clinical scoring system to predict in-hospital  
341 mortality in patients with OHCA, with high specificity and high predictive value for in-  
342 hospital death. It could be used to support the prognostication process for physicians, and can  
343 help clinicians explain the prognosis to relatives in an easy-to-understand, objective fashion,  
344 to realistically prepare them for the future.

## 345 **References**

- 346 1. Gräsner J-T, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, Wnent J,  
347 Tjelmeland IBM, Ortiz FR, Maurer H, Baubin M, Mols P, Hadžibegović I, Ioannides  
348 M, Škulec R, Wissenberg M, Salo A, Hubert H, Nikolaou NI, Lóczi G, Svavarsdóttir  
349 H, Semeraro F, Wright PJ, Clarens C, Pijls R, Cebula G, Correia VG, Cimpoesu D,  
350 Raffay V, Trenkler S. EuReCa ONE-27 Nations, ONE Europe, ONE Registry: A  
351 prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27  
352 countries in Europe. *Resuscitation* 2016;**105**:188–195.
- 353 2. Nolan JP, Soar J, Cariou A, Cronberg T, Moulaert VRM, Deakin CD, Bottiger BW,  
354 Friberg H, Sunde K, Sandroni C. European Resuscitation Council and European  
355 Society of Intensive Care Medicine Guidelines for Post-resuscitation Care 2015.  
356 *Resuscitation* 2015;**95**:202–222.
- 357 3. Bossaert LL, Perkins GD, Askitopoulou H, Raffay VI, Greif R, Haywood KL,  
358 Mentzelopoulos SD, Nolan JP, Voorde P Van de, Xanthos TT, Georgiou M, Lippert  
359 FK, Steen PA. European Resuscitation Council Guidelines for Resuscitation 2015.  
360 *Resuscitation* 2015;**95**:302–311.
- 361 4. Sunde K, Kramer-Johansen J, Pytte M, Steen PA. Predicting survival with good  
362 neurologic recovery at hospital admission after successful resuscitation of out-of-  
363 hospital cardiac arrest: the OHCA score. *Eur Heart J* 2007;**28**:773–773.
- 364 5. Gold B, Puertas L, Davis SP, Metzger A, Yannopoulos D, Oakes DA, Lick CJ,  
365 Gillquist DL, Holm SYO, Olsen JD, Jain S, Lurie KG. Awakening after cardiac arrest  
366 and post resuscitation hypothermia: Are we pulling the plug too early? *Resuscitation*  
367 2014;**85**:211–214.
- 368 6. Adrie C, Cariou A, Mourvillier B, Laurent I, Dabbane H, Hantala F, Rhaoui A,  
369 Thuong M, Monchi M. Predicting survival with good neurological recovery at hospital

- 370 admission after successful resuscitation of out-of-hospital cardiac arrest: the OHCA  
371 score. *Eur Heart J* 2006;**27**:2840–2845.
- 372 7. Eisenberg M, Hallstrom A, Bergner L. The ACLS Score. *JAMA*; 1981;**246**:50.
- 373 8. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from out-  
374 of-hospital cardiac arrest: A graphic model. *Ann Emerg Med* 1993;**22**:1652–1658.
- 375 9. Aschauer S, Dorffner G, Sterz F, Erdogmus A, Laggner A. A prediction tool for initial  
376 out-of-hospital cardiac arrest survivors. *Resuscitation* 2014;**85**:1225–1231.
- 377 10. Thompson RJ, McCullough PA, Kahn JK, O’Neill WW. Prediction of death and  
378 neurologic outcome in the emergency department in out-of-hospital cardiac arrest  
379 survivors. *Am J Cardiol* 1998;**81**:17–21.
- 380 11. Ishikawa S, Niwano S, Imaki R, Takeuchi I, Irie W, Toyooka T, Soma K, Kurihara K,  
381 Izumi T. Usefulness of a simple prognostication score in prediction of the prognoses of  
382 patients with out-of-hospital cardiac arrests. *Int Heart J* 2013;**54**:362–370.
- 383 12. Sladjana A. A prediction survival model for out-of-hospital cardiopulmonary  
384 resuscitations. *J Crit Care* 2011;**26**:223.e11-223.e18.
- 385 13. Ahmad R, Lumley S, Lau YC. NULL-PLEASE: A new ‘Futility score’ in the  
386 management of survivors of out-of-hospital cardiac arrest. *Resuscitation*  
387 2016;**106**:e83.
- 388 14. Potpara TS, Mihajlovic M, Stankovic S, Jozic T, Jozic I, Asanin MR, Ahmad R, Lip  
389 GYH. External Validation of the Simple NULL-PLEASE Clinical Score in Predicting  
390 Outcome of Out-of-Hospital Cardiac Arrest. *Am J Med* 2017;**130**:1464.e13-1464.e21.
- 391 15. Sauerbrei W, Schumacher M. A bootstrap resampling procedure for model building:  
392 Application to the cox regression model. *Stat Med* 1992;**11**:2093–2109.
- 393 16. Sasson C, Rogers MAM, Dahl J, Kellermann AL. Predictors of survival from out-of-  
394 hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual*

- 395 *Outcomes* American Heart Association, Inc.; 2010;**3**:63–81.
- 396 17. Berdowski J, Berg RA, Tijssen JGP, Koster RW. Global incidences of out-of-hospital  
397 cardiac arrest and survival rates: Systematic review of 67 prospective studies.  
398 *Resuscitation* Elsevier; 2010;**81**:1479–1487.
- 399 18. Jabre P, Bougouin W, Dumas F, Carli P, Antoine C, Jacob L, Dahan B, Beganton F,  
400 Empana J-P, Marijon E, Karam N, Loupy A, Lefaucheur C, Jost D, Cariou A, Adnet F,  
401 Rea TD, Jouven X. Early Identification of Patients With Out-of-Hospital Cardiac  
402 Arrest With No Chance of Survival and Consideration for Organ Donation. *Ann Intern*  
403 *Med* 2016;**165**:770.
- 404 19. Petrie J, Easton S, Naik V, Lockie C, Brett SJ, Stümpfle R. Hospital costs of out-of-  
405 hospital cardiac arrest patients treated in intensive care; a single centre evaluation  
406 using the national tariff-based system. *BMJ Open* 2015;**5**:e005797.
- 407 20. Sundermann ML, Salcido DD, Koller AC, Menegazzi JJ. Inaccuracy of patient care  
408 reports for identification of critical resuscitation events during out-of-hospital cardiac  
409 arrest. *Am J Emerg Med* 2015;**33**:95–99.
- 410 21. Maupain C, Bougouin W, Lamhaut L, Deye N, Diehl J-L, Geri G, Perier M-C,  
411 Beganton F, Marijon E, Jouven X, Cariou A, Dumas F. The CAHP (Cardiac Arrest  
412 Hospital Prognosis) score: a tool for risk stratification after out-of-hospital cardiac  
413 arrest. *Eur Heart J* 2016;**37**:3222–3228.
- 414 22. Kiehl EL, Parker AM, Matar RM, Gottbrecht MF, Johansen MC, Adams MP, Griffiths  
415 LA, Dunn SP, Bidwell KL, Menon V, Enfield KB, Gimple LW. C-GRaPH: A  
416 Validated Scoring System for Early Stratification of Neurologic Outcome After Out-  
417 of-Hospital Cardiac Arrest Treated With Targeted Temperature Management. *J Am*  
418 *Heart Assoc* 2017;**6**:e003821.

419

420  
421