LOFAR observations of galaxy clusters in HETDEX Extraction and self-calibration of individual LOFAR targets

R. J. van Weeren¹, T. W. Shimwell^{2, 1}, A. Botteon¹, G. Brunetti³, M. Brüggen⁴, J. M. Boxelaar¹, R. Cassano³, G. Di Gennaro¹, F. Andrade-Santos⁵, E. Bonnassieux⁶, A. Bonafede^{6,3}, V. Cuciti⁴, D. Dallacasa^{3,6}, F. de Gasperin⁴, F. Gastaldello⁷, M. J. Hardcastle⁸, M. Hoeft⁹, R. P. Kraft⁵, S. Mandal¹, M. Rossetti⁷, H. J. A. Röttgering¹, C. Tasse^{10, 11, 12}, and A. G. Wilber^{13, 4}

- Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands
- e-mail: rvweeren@strw.leidenuniv.nl
- 2 ASTRON, The Netherlands Institute for Radio Astronomy, Postbus 2, 7990 AA Dwingeloo, The Netherlands
- INAF-Istituto di Radioastronomia, Via Gobetti 101, I-40129 Bologna, Italy
- University of Hamburg, Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany
- Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138
- 6 Dipartimento di Fisica e Astronomia, Università di Bologna, via P. Gobetti 93/2, I-40129 Bologna, Italy
- INAF-IASF Milano, Via A. Corti 12, I-20133, Milano, Italy
- Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK
- Thüringer Landessternwarte, Sternwarte 5, 07778 Tautenburg, Germany
- ¹⁰ GEPI, Observatoire de Paris, CNRS, Université Paris Diderot, 5 place Jules Janssen, 92190 Meudon, France
- 11 Centre for Radio Astronomy Techniques and Technologies, Department of Physics and Electronics, Rhodes University, Graham-
- stown 6140, South Africa
- 12 USN, Observatoire de Paris, CNRS, PSL, UO, Nançay, France
- ¹³ Curtin Institute of Radio Astronomy, 1 Turner Avenue, Technology Park, Bentley WA 6102, Australia

ABSTRACT

Diffuse cluster radio sources, in the form of radio halos and relics, reveal the presence of cosmic rays and magnetic fields in the intracluster medium (ICM). These cosmic rays are thought to be (re-)accelerated through ICM turbulence and shock waves generated by cluster merger events. Here we characterize the presence of diffuse radio emission in known galaxy clusters in the HETDEX Spring Field, covering 424 deg². For this, we developed a method to extract individual targets from LOFAR observations processed with the LoTSS DDF-pipeline. This procedure enables improved calibration and joint imaging and deconvolution of multiple pointings of selected targets. The calibration strategy can also be used for LOFAR Low-Band Antenna (LBA) and international-baseline observations.

The fraction of Planck PSZ2 clusters with any diffuse radio emission apparently associated with the ICM is $73 \pm 17\%$. We detect a total of 10 radio halos and 12 candidate halos in the HETDEX Spring Field. Five clusters host radio relics. The fraction of radio halos in Planck PSZ2 clusters is $31 \pm 11\%$, and $62 \pm 15\%$ when including the candidate radio halos. Based on these numbers, we expect that there will be at least 183 ± 65 radio halos found in the LoTSS survey in PSZ2 clusters, in agreement with predictions. The integrated flux densities for the radio halos were computed by fitting exponential models to the radio images. From these flux densities, we determine the cluster mass (M_{500}) and Compton Y parameter (Y_{500}) 150 MHz radio power ($P_{150 \text{ MHz}}$) scaling relations for Planck PSZ2-detected radio halos. We find that the slopes of these relations are steeper than those determined from the 1.4 GHz radio powers. However, considering the uncertainties this is not a statistically significant result.

Key words. Galaxies: clusters: — Galaxies: clusters: intracluster medium — large-scale structure of universe — Radiation mechanisms: non-thermal - X-rays: galaxies: clusters

1. Introduction

implies that some form of in-situ particle (re-)acceleration is required to explain the Mpc extent of these sources.

Radio observations have revealed the presence of Mpc-scale radio sources associated with intracluster medium (ICM) in a growing number of galaxy clusters. This indicates that the ICM is filled with cosmic ray (CR) electrons and magnetic fields. Diffuse cluster radio sources are commonly divided into radio relics (radio shocks), giant halos, and mini-halos (for reviews see Feretti et al. 2012; Brunetti & Jones 2014; van Weeren et al. 2019). Importantly, the short lifetime (~ 10^{7-8} yrs) of the CR electrons

Giant radio halos are Mpc-size sources that approximately follow the X-ray emission from the thermal ICM. They are predominantly found in merging clusters (Cassano et al. 2010b). The radio power of giant halos correlates with cluster mass (e.g., Cassano et al. 2013) and often used mass proxies are the cluster's X-ray luminosity (L_X) or integrated Compton Y parameter. The upper limits derived for clusters that are dynamically relaxed are underluminous with respect to these correlations (e.g., Brunetti et al. 2007; Cassano et al. 2013). The fraction of clusters with

radio halos is about 30% (Venturi et al. 2008; Kale et al. 2013, 2015) for $L_{X,0.1-2.4 \text{ keV}} > 5 \times 10^{44} \text{ erg s}^{-1}$ clusters in the range 0.2 < z < 0.4. For the most massive (~ $10^{15} \text{ M}_{\odot}$) clusters the occurrence fraction is as high as ~ 80%, and there is evidence that this fraction decreases for lower mass clusters (Cuciti et al. 2015).

Two main classes of models have been proposed for the origin of radio halos. In the turbulent re-acceleration model, particles are re-accelerated by merger induced magnetohydrodynamical turbulence (Brunetti et al. 2001; Petrosian 2001; Brunetti & Lazarian 2007; Miniati 2015). In the hadronic model, the radio emission is produced by secondary electrons that arise from hadronic collisions (e.g., Dennison 1980; Blasi & Colafrancesco 1999; Dolag & Enßlin 2000). Most observational evidence nowadays is in favor of the turbulent re-acceleration model. This evidence includes the discovery of radio halos with ultra-steep radio spectra (USSRH; e.g., Brunetti et al. 2008) and the non-detection of gamma ray emission from the Coma cluster at the level that would be necessary to generate the observed radio emission (Brunetti et al. 2012, 2013; Ackermann et al. 2016; Brunetti et al. 2017). Models for radio halos invoking a combination of turbulent re-acceleration and generation of secondary particles that are consistent with gamma ray limits have also been proposed (e.g., Brunetti & Lazarian 2011; Pinzke et al. 2017; Brunetti et al. 2017).

Radio mini-halos are smaller sized halos (~200–500 kpc) that are exclusively found in relaxed cool-core clusters. Recently, Giacintucci et al. (2017) found that mini-halos are rather common in massive cool-core clusters; about 80% of such clusters host them. The radio emission from mini-halos surrounds the central AGN associated with the brightest cluster galaxy (BCG). Similar to giant radio halos some form of in-situ acceleration is required to power them (e.g., Gitti et al. 2004; Giacintucci et al. 2014). Mini-halos have been explained by hadronic scenarios (e.g., Pfrommer & Enßlin 2004; Fujita et al. 2007; Keshet & Loeb 2010; Fujita & Ohira 2013), or by turbulent re-acceleration induced by gas sloshing motions (Mazzotta & Giacintucci 2008; ZuHone et al. 2013).

Radio relics are polarized, elongated sources found in galaxy cluster outskirts (e.g., Enßlin et al. 1998). They have sizes that can extend up to about 2 Mpc. High-resolution observations show that radio relics often have filamentary morphologies (e.g. Bagchi et al. 2006; Di Gennaro et al. 2018; Rajpurohit et al. 2020, 2018). These sources trace ICM shock waves with low to moderate Mach numbers (e.g., Finoguenov et al. 2010; Akamatsu & Kawahara 2013; Ogrean & Brüggen 2013; Shimwell et al. 2015). The physical mechanisms by which particles are (re-)accelerated at shocks are still being debated. One possibility to accelerate particles at shocks is via the diffusive shock acceleration mechanism (DSA; e.g., Blandford & Eichler 1987; Jones & Ellison 1991; Malkov & O'C Drury 2001). However, this mechanism is thought to be rather inefficient for weak ICM shocks if particles are accelerated from the thermal pool and it fails to explain the observed radio power and spectral indices in a number of cases (e.g., Pinzke et al. 2013; Vazza & Brüggen 2014; Vazza et al. 2016; van Weeren et al. 2016a; Botteon et al. 2020a). It has therefore been proposed that some form of reacceleration of pre-existing fossil CR electrons takes place (e.g., Markevitch et al. 2005; Giacintucci et al. 2008; Kang & Ryu 2011; Kang et al. 2012; van Weeren et al. 2016a; Botteon et al. 2016a). Observations do provide support for this model, where the fossil CRs originate from the tails and lobes of radio galaxies (Bonafede et al. 2014a; Shimwell et al. 2015; van Weeren et al. 2017). It should be noted, however, that for some relics DSA

of thermal pool electrons seems to be sufficient to explain their luminosity (e.g., Botteon et al. 2016b; Locatelli et al. 2020).

Low-frequency observations provide important information about particle acceleration processes. The turbulent reacceleration model for radio halos predicts that the occurrence rate of halos should be higher at low frequencies (Cassano 2010; Cassano et al. 2012). Radiative losses of CR electrons limit the acceleration by turbulence via second order Fermi mechanisms, causing a break in the energy spectrum of these electrons (Cassano et al. 2004; Cassano & Brunetti 2005; Cassano et al. 2006; Brunetti et al. 2008). This in turn results in a synchrotron spectrum that becomes steeper in the situations where the amount of turbulent energy is smaller (less powerful mergers) or the energy losses are stronger (higher redshift).

Another important role of low-frequency observations is to probe the connection between re-acceleration processes and fossil radio plasma. Because of synchrotron and inverse Compton (IC) losses, as relativistic electrons age they emit almost exclusively at low frequencies. These seed fossil CR particles are a critical ingredient in both shock and turbulent re-acceleration models. Possible examples of revived fossil plasma, by compression (Enßlin & Gopal-Krishna 2001; Enßlin & Brüggen 2002) or other mechanisms (e.g., Markevitch et al. 2005), have been been detected in some clusters (e.g., Slee et al. 2001; van Weeren et al. 2017; de Gasperin et al. 2017; Mandal et al. 2020). However, this is probably just the tip of the iceberg.

Finally, recent LOFAR observations have entered uncharted territories, discovering the existence of radio bridges that connect pairs of massive and pre-merging clusters (Govoni et al. 2019; Botteon et al. 2020c). One possibility is that these bridges originate from second order Fermi acceleration mechanisms powered by turbulence filling these vast regions (Brunetti & Vazza 2020).

The LOFAR Two-metre Sky Survey (LoTSS Shimwell et al. 2017) is a deep 120-168 MHz survey that will cover the entire northern sky when completed. This survey is carried out with the High Band Antenna (HBA) stations of LOFAR (van Haarlem et al. 2013). The nominal sensitivity of this survey is $0.1\ {\rm mJy}\ {\rm beam}^{-1}$ at a resolution of 6". The first data release (DR1) covers 424 deg² in the region of the HETDEX Spring field (Shimwell et al. 2019). Given its unprecedented survey depth at low frequencies, resolution, and sky coverage, LoTSS will play an important role in determining the statistics of diffuse cluster sources. In this paper we present the first results on a full sample of galaxy clusters in the LoTSS DR1-area using improved calibration techniques. This is required to properly study extended low-surface brightness cluster sources. The outline of this paper is as follows. In Sect. 2 we describe the procedure we developed to extract and re-calibrate targets of interest from the LoTSS data products. The sample selection is described in Sect. 3. The results are presented in Sect. 4 and we end with a discussion and conclusions in Sects. 5 and 6.

Throughout this paper we assume a Λ CDM cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, and $\Omega_{\Lambda} = 0.7$. All images are in the J2000 coordinate system.

2. Extraction and re-calibration

Here we utilize LOFAR observations that were taken as part of the LoTSS survey. These are typically 8 hr observations. The pointing centers of these observations are placed with the aim of obtaining close to uniform sensitivity coverage of the northern sky. The LoTSS survey design and the observations are discussed in detail in Shimwell et al. (2017, 2019).

2.1. Extraction

The LoTSS processing pipeline (DDF-PIPELINE) delivers images of the full field of view (FoV) of the Dutch LOFAR HBA stations (Tasse 2014a,b; Shimwell et al. 2019). These images have a resolution of 6" and an r.m.s. noise level of the order of 100 μ Jy beam⁻¹. This is achieved by correcting for the direction dependent effects (DDE) in the LOFAR data (due to the ionosphere and imperfect station beam models). For LoTSS, DDE corrections are applied towards 45 directions (facets). The DDF-PIPELINE is optimized to create images of LOFAR's full FoV and carry out survey science. The latest version 2 of the DDF-PIPELINE, which we employ in this work, is described in Tasse et al. (2020). Version 2 of the pipeline is a major improvement compared to version 1 which was used for public Data Release 1. The images from DR1 are generally not suitable to study extended low-surface brightness cluster sources (see Section 3.7 in Shimwell et al. 2019).

The individual images produced by the DDF-PIPELINE¹ are very large, 20,000 \times 20,000 pixels with a pixel size of 1.5". This makes re-imaging with different settings, e.g., uv-ranges, weighting schemes, and deconvolution algorithms, expensive. The tessellation of the sky into 45 calibration facets is done in a fully automated way. The DDE calibration takes all sources in a facet into account, assuming there are no DDE inside a facet. This means that for certain specific targets of interest the facet layout is not optimal. Experience with various faceting schemes has shown that it is often possible to further improve the quality of the DDE calibration for a specific target (sometimes at the expense of other sources).

To allow flexible re-imaging and optimize calibration towards targets of interest, we extended the DDF-PIPELINE with an optional step. In this step all sources are subtracted, apart from those in a specific user defined region, from the visibility data and after applying their DDE calibration solutions. The model visibility prediction is done with the DDFacet imager (Tasse et al. 2018). An important requirement for extraction is that region of interest needs to be quite small, roughly similar to the size of the original facet, but ideally smaller. This is directly related to the fact that we want to improve the DDE calibration, which was limited by the original facet size and the "incorrect" assumption that the DDE correction is constant across that entire facet. This requirement competes with another requirement that there is sufficient flux density available for calibration. A smaller extraction region will have less flux density available for the calibration.

2.1.1. Phase shifting and averaging

After subtracting all sources (clean components) from the uvdata, apart from those in the region of interest, we phase shift the uv-data. The new phase center is placed at the center of the region of interest, i.e., the region where the sources were kept. In practice we use DS9 (Joye & Mandel 2003) region files which can be easily generated. After phase shifting, the data are averaged in time and frequency. The extracted region has a small angular extent and thus bandwidth and time smearing are not an issue. By default, the data are averaged to 16 s and 0.39 MHz. With these averaging parameters, the size of the dataset is reduced by a factor of 8. The visibility data are compressed using Dysco to further reduce data size by about a factor of four (Offringa 2016). Additional averaging is often allowed from the point of bandwidth and time smearing, but we found that more averaging can hinder ionospheric calibration. Optionally, additional radio frequency interference (RFI) flagging is carried out with AOFlagger on the output data (Offringa 2010; Offringa et al. 2010, 2012).

When the data is shifted to the new phase center we also correct for the LOFAR station beam response in this direction. Additionally, the visibility weights set by the DDF-pipeline (for details see Bonnassieux et al. 2018) are updated. They are multiplied by a factor inversely proportional to the station beam response. In this way, we can optimally combine observations from multiple pointing centers with joint imaging and deconvolution. Combining visibility data in this way is allowed because the beam correction is close to constant across the small region that is extracted².

2.2. Self-calibration

The next step in the "extraction" process for a target of interest is to self-calibrate the data. The starting point of this step are one or more phase shifted and averaged datasets for the target of interest, each one corresponding to a different observation with a potentially different pointing center. The direction independent (DI) full Jones calibrations are carried over from the DDF-pipeline. We do not carry over the DDE solutions to avoid the issue with the "negative haloes" described in Tasse et al. (2020).

The self-calibration steps consist of three rounds of 'tecandphase' calibration with DPPP (van Diepen & Dijkema 2018). This is followed by several rounds of diagonal (i.e., XX and YY) gain calibration using a longer solution interval. The shorter timescale 'tecandphase' solutions are pre-applied when solving for the diagonal gains. This scheme somewhat mimics the facet-calibration scheme (van Weeren et al. 2016b) and the DDF-pipeline which also pre-applies fast phase/TEC solutions before solving for slow gain solutions. The diagonal gain solutions are filtered for outliers with LoSoTo (de Gasperin et al. 2019). The solution intervals are automatically determined based on the amount of apparent compact source flux in the extracted region. Solutions intervals for the 'tecandphase' calibration are between 16 s and 48 s. Solution intervals for the diagonal gains are between 16 min and 48 min. All solution intervals are set per observation since the apparent flux in the target of interest region can differ. Solution intervals along the frequency axis for the diagonal gains are between 2 MHz and 6 MHz. For HBA data, a minimum of 0.3 Jy compact (apparent) source flux is needed for the self-calibration steps to converge well. If needed, all solution intervals can be manually controlled by the user. An overview of the parameters used is given in Table 1.

The imaging is done with WSClean (Offringa et al. 2014) with wideband joint deconvolution mode ('channelsout' 6 default), optionally with multi-scale clean (Offringa & Smirnov 2017). The DDFacet imager (Tasse et al. 2018) can also be used instead of WSClean. As default Briggs weighting -0.5 is used. The imaging includes automatic clean masking from the DDF-pipeline, with a default 5σ threshold. Baseline based averaging is also employed when imaging for performance. An inner uv-cut of 80λ is used in the imaging. For the calibration, a default inner uv-cut of 350λ (corresponding to 0.65°) is employed. This value is a compromise between the number of baselines in

¹ https://github.com/mhardcastle/ddf-pipeline

² If that approximation does not hold we use the Image Domain Gridding algorithm which corrects each phase rotated dataset with the correct beam response (van der Tol et al. 2018).

 Table 1. Default imaging and calibration parameters for HBA extraction

parameter	value
inner uv-range (calibration)	350 ^a
inner uv-range (imaging, λ)	80
clean mask threshold (sigma)	5
pixelsize (arcsec)	1.5
Briggs robust weighting	-0.5
channelsout ^b	6

Notes. ^(*a*) Increased to 750 λ for targets with extended low-surface brightness emission. ^(*b*) WSClean wideband deconvolution setting

the calibration and the increased difficulty of modelling very extended structures.

In Figure 1 we show an example of the self-calibration process. The top panel shows the first image before any self-calibration. Note that here two different observations (L254483 and L719874), with different pointing centers, are jointly imaged and deconvolved. The only calibration that is applied here is the DI calibration from the DDF-pipeline. In the middle panel we show the result after three rounds of 'tecandphase' self-calibration. The radial patterns, caused by the ionosphere disappear after this calibration. In the bottom panel we show the results after additional diagonal gain calibration. The diagonal gain calibration corrects for the imperfect knowledge of the station beam response or other slowly varying gain errors.

In Figure 2 we display two comparisons between the default LoTSS mosaics and newly extracted and re-calibrated targets (in this case the clusters Abell 1430 and 1294). As can be seen, the calibration for the targets of interest has improved. The improvement is the result of shrinking the size of the calibration region so that a nearby bright compact source is better calibrated. The extraction and self-calibration scheme has already been successfully applied in various recent works (e.g., Botteon et al. 2020c,b; Mandal et al. 2020; Hardcastle et al. 2019; Cassano et al. 2019; Gu et al. 2019).

2.2.1. Other usage

The self calibration step has been designed in such a way that it can also be carried out on other (non-LoTSS) LOFAR extracted datasets. It offers full flexibility for the various effects that can be solved for in DPPP, including constraint solves. For example, it can enforce the gains to be smooth along the frequency axis or enforce the same solutions for certain stations. The self calibration step has been successfully used on LOFAR HBA international-baseline data and on the Low Band Antennas (LBA) observations. Figure 3 shows an example of it working on LBA and international baseline data.

3. HETDEX-DR1 area galaxy cluster sample

We applied the extraction scheme described in Section 2.1 to a sample of galaxy clusters that fall inside the LOFAR Data Release 1 area (Shimwell et al. 2017, 2019). We selected clusters from the all-sky PSZ2 Planck catalog (Planck Collaboration et al. 2016) of Sunyaev-Zel'dovich (SZ) sources, see Figure 4. This sample of 26 clusters serves as our primary sample for our statistical investigations and is listed in Table 2.

In addition to the SZ-selected sample, we compiled a secondary sample by visually inspecting the LoTSS images at the location of known clusters for the presence of diffuse emission and possible revived fossil plasma sources. Clusters in this sample with (candidate) extended radio emission are from the Abell and Zwicky (Abell 1958; Zwicky et al. 1961; Corwin 1974), MCXC (Piffaretti et al. 2011), GMBCG (Hao et al. 2010), MaxBCG (Koester et al. 2007), and WHL (Wen et al. 2012) catalogs.

To search for diffuse radio emission we produced images with the emission from compact sources subtracted. This was done by first imaging the calibrated datasets with a uv-cut corresponding to a physical scale of 0.4 Mpc. This model was subsequently subtracted from the visibility data.

3.1. Flux density measurements

The flux density measurements for all sources, except radio halos, were done manually by placing a polygon around the sources and integrating the flux density. The uncertainty on the flux density is given by

$$\sigma_S^2 = N_{\text{beams}} \sigma_{\text{rms}}^2 + \sigma_{\text{sub}}^2 + (f \times S)^2 , \qquad (1)$$

where f = 0.2 is the absolute flux-scale uncertainty (Shimwell et al. 2019), N_{beams} the number of beams covering the source, σ_{rms} the map noise, and σ_{sub} the uncertainty due to compact source subtraction. The first two terms of Eq. 1 represent the statistical uncertainty on the flux density measurement. The uncertainty on the compact source subtraction is given by

$$\sigma_{\rm sub}^2 = \sum_i N_{\rm beams,i} \sigma_{\rm rms}^2 \,, \tag{2}$$

where the sum is taken over all *i* sources that were subtracted in the polygon.

The radio halo integrated flux densities were determined by fitting exponential profiles (Murgia et al. 2009) to the radio images of the form

$$I(r) = I_0 e^{-r/r_e} , (3)$$

where r_e is a characteristic *e*-folding radius and I_0 the central surface brightness. We used the image where compact sources were subtracted. Also, all images were smoothed to a beam size corresponding to a physical scale of 50 kpc at the cluster's redshift. For the fitting we used Halo-FDCA³ described in Boxelaar et al. (2020). Extended sources, such as large tailed radio galaxies, were masked during the fitting (in case they were not fully subtracted). The flux density was integrated to a radius of $3r_e$, as proposed by Murgia et al. (2009). Integrating up to a radius of $3r_e$ results in 80% of the flux density when compared to using an infinite radius. This also avoids to some extent the uncertainties related to the extrapolation of the profile to large radii where the radio surface brightness is below the detection limit. The fitting uses a Markov Chain Monte-Carlo method to determine the radio halo parameters and associated uncertainties. For some clusters, when specifically mentioned in Section 4, we fitted an elliptical exponential model (see Boxelaar et al. 2020). In these cases, a major and minor characteristic *e*-folding radius and position angle are determined. This is done for radio halos that clearly display an elongated, rather than circular, shape. In this work we did not use the skewed (asymmetric) models that are available in Halo-FDCA. A full exploration of radio halo fitting models and methods is beyond the scope of this work.

For the total uncertainty on the radio halo integrated flux density we add the uncertainty from the MCMC (statistical errors) and flux calibration uncertainty in quadrature.

³ https://github.com/JortBox/Halo-FDCA

3.2. X-ray observations

We searched the *Chandra* and *XMM-Newton* archives for the available X-ray observations of the clusters in the sample. When available, data were retrieved and processed with CIAO 4.11 using CalDB v4.8.2 and SAS v16.1.0 following standard data reduction recipes. We used the time periods of the observations cleaned by anomalously high background to produce cluster exposure-corrected images in the 0.5 - 2.0 keV band. These images are used in the paper to investigate the connection between the thermal and non-thermal components in the ICM.



Fig. 1. Images showing the subsequent improvement during selfcalibration on the extracted data of the cluster Abell 1430. The top panel shows the starting point of this process: the DDF-pipeline (v2) direction independent (DI) calibration. The middle panel displays the results after DPPP 'tecandphase' calibration. The bottom panel shows the final image after DPPP diagonal gain calibration.



Fig. 2. Comparison between the LoTSS DR2 mosaics and the extracted and re-calibrated images for the clusters Abell 1430 (top panels) and Abell 1294 (bottom panels). For both clusters optimizing the calibration towards a nearby bright compact source improved the image quality. These errors in the LoTSS DR2 mosaics were caused by the spatially varying ionosphere. Note that the DR2 pipeline could have achieved more or less the same results with a different facet layout (but then at the cost of reducing image quality in other directions).



Fig. 3. *Top row:* Images showing the subsequent improvement during self-calibration on extracted 43–67 MHz LBA data. The left panel shows the starting point of this process: a direction independent calibrated image. The middle panel displays the results after DPPP 'tec' calibration, with all LOFAR core stations forced ("antenna constraint" solve) to having the same solution. The right panel shows the final image after DPPP scalarcomplex gain calibration. *Bottom row:* The same as the top row images but for LOFAR international baseline 120–168 MHz HBA data and without using an "antenna constraint" solve.



Fig. 4. The redshift-mass distribution of PSZ2 clusters. Clusters that are located in the HETDEX DR1 area are indicated with dark red points.

Table 2.	. HETDEX	DR1-area	cluster	sample
----------	----------	----------	---------	--------

Cluster	alternative name(s)	RA	DEC	redshift	M _{500,SZ}
PSZ2 G080.16+57.65	Abell 2018	15 01 08	+47 16 37	0.0878	$2.51^{+0.20}_{-0.21}$
PSZ2 G084.10+58.72		14 49 01	+48 33 24	0.7310	$5.40^{+0.62}_{-0.62}$
PSZ2 G086.93+53.18	WHL J228.466+52.8333	15 14 00	+52 48 14	0.6752	$5.45^{+0.50}_{-0.52}$
PSZ2 G087.39+50.92	[WH2015]0986	15 26 33	+540908	0.7480	$5.16^{+0.32}$
PSZ2 G088.98+55.07		14 59 01	+52.49.01	0.7023	$4.92^{+0.60}$
PSZ2 G089 52+62 34	Abell 1904	14 22 13	+48 29 54	0.0701	$1.83^{+0.19}$
1 522 (000).52 (02.5)	RX I1422 1+4831	112213	110 29 31	0.0701	-0.20
PS72 G095 22+67 41	$RXC 11351 7 \pm 4622$	13 51 45	+46 22 00	0.0625	1 500.21+
1 522 6075.22 + 07.41	MCXC I1351 7+4622	15 51 45	140 22 00	0.0025	-0.22
PS72 G096 14+56 24	Abell 1940	14 35 21	+55.08.29	0 1398	$277^{+0.24}$
15220070.14750.24	PX 11/35 /+5508	17 33 21	155 00 27	0.1570	2.77 - 0.26
DS72 COOS 11 + 56 50	A ball 1020	14 27 25	155 45 02	0 1318	2 82+0.28
F SZ2 0098:44+50:59	AUCH 1920 DV 11427 4 : 5545	14 27 23	+33 43 02	0.1310	$2.03_{-0.26}$
DS72 C000 86 158 45	$\mathbf{KA} \mathbf{J} \mathbf{I} 4 2 7 .4 \mathbf{+} \mathbf{J} 5 4 3 5 4 7 0 4$	1 / 1 / / 2	54 47 01	0 6 1 6 0	6 05+0.48
FSZ2 0099.80+38.43	WHI 1212 607 54 7844	14 14 45	+34 47 01	0.0100	$0.03_{-0.49}$
PC72 C 106 (1 + 66 71)	WHLJ213.097+34.7844	12 20 20	10 00 40	0 2214	4 67+0.55
PSZ2 G100.01+00.71	A h - 11 1750	13 30 29	+49 08 48	0.3314	$4.07_{-0.57}$
PSZ2 G107.10+05.32	Abell 1758	13 32 33	+30 29 09	0.2799	$8.22_{-0.28}$
PSZ2G111.75+70.37	Abell 169/	13 13 03	+46 16 52	0.1830	$4.34_{-0.33}^{+0.32}$
	RXCJ1313.1+4616	12 15 05	51 40 00	0.0000	(
PSZ2G114.31+64.89	Abell 1703	13 15 05	+51 49 02	0.2836	$6.76_{-0.38}^{+0.30}$
PSZ2 G114.99+70.36	Abell 1682	13 06 50	+46 33 27	0.2259	$5.70^{+0.35}_{-0.35}$
PSZ2G118.34+68.79	ZwCl 1259.0+4830	13 01 24	+48 14 31	0.2549	$3.77^{+0.45}_{-0.52}$
	[WH2015]0746				0.50
PSZ2G123.66+67.25	Abell 1622	12 49 41	+49 52 18	0.2838	$4.38^{+0.50}_{-0.52}$
	ZwCl1247.2+5008				
PSZ2 G133.60+69.04	Abell 1550	12 29 02	+47 37 21	0.2540	$5.88^{+0.38}_{-0.42}$
PSZ2G135.17+65.43	WHL J121912.2+505435	12 19 12	+50 54 35	0.5436	$6.00^{+0.57}_{-0.62}$
PSZ2G136.92+59.46	Abell 1436			0.0650	$1.80^{+0.17}_{-0.16}$
	RXC J1200.3+5613				0.10
PSZ2G143.26+65.24	Abell 1430	11 59 17	+49 47 37	0.3634	$7.65^{+0.42}_{-0.44}$
	RXCJ1159.2+4947				0.44
	ZwCl1156.4+5009				
PSZ2G144.33+62.85	Abell 1387	11 49 05	+51 35 08	0.1320	$2.66^{+0.33}_{-0.38}$
	RXC J1149.0+5135				0.50
PSZ2G145.65+59.30	Abell 1294	11 32 42	+54 13 12	0.3475	$4.73^{+0.59}_{-0.64}$
	ZwCl1129.6+5430				-0.04
PSZ2G150.56+58.32	MACS J1115.2+5320	11 15 11	+53 19 39	0.4660	$7.55^{+0.50}_{-0.52}$
	RXCJ1115.2+5320				-0.52
PSZ2G151.62+54.78	RX J105453.3+552102	10 54 52	+55 21 13	0.4864	$5.37^{+0.68}_{-0.75}$
	[WH2015]0472				-0.75
	1RXSJ105453.3+552102				
PSZ2G156.26+59.64	[WH2015] 0485	11 08 30	+50 16 02	0.6175	$6.77^{+0.59}_{-0.60}$
Abell 1156		11 04 56	+47 25 15	0 2091	-0.00
Abell 1314		11 34 49	+490440	0.0335	
Abell 1615		12 47 43	+48 51 57	0.2106	
GMBCG J211.77332+55.09968		14 06 55	+550402	0.2506	
MaxBCG J173.04772+47.81041		11 32 11	+47 48 38	0.2261	
NSC J143825+463744		14 38 46	+46 39 56	0.0357	
RXC J1053.7+5452	MCXC J1053.7+5452	10 53 44	+54 52 20	0.0704	• • •
WHL J125836.8+440111		12 58 37	+44 01 11	0.5339	
WHL J122418.6+490549		12 24 19	+49 05 50	0.1004	
WHL J124143.1+490510		12 41 43	+49 05 10	0.3707	
WHL J132226.8+464630		13 22 27	+46 46 30	0.3718	
WHL J132615.8+485229		13 26 16	+48 52 29	0.2800	
WHL J133936.0+484859		13 39 36	+48 48 59	0.3265	
WHL J134746.8+475214		13 47 47	+47 52 15	0.1695	

4. Results

Below we first describe the LOFAR results for the individual clusters. By default images were made using Robust weighting -0.5 (Briggs 1995). In addition, images at lower resolution were produced using a Gaussian taper to down weight the visibilities from longer baselines. Images with the emission from compact sources subtracted are also shown for some clusters.

The classification of diffuse sources is summarised in Table 3. Descriptions and images of clusters for which no diffuse emission was detected are given in Appendix A. For optical overlays we use Pan-STARRS *gri* images (Chambers et al. 2016).

Cluster	classification	LLS	S 144
		Mpc	mJy
PSZ2 G080.16+57.65	cHalo, Relic	(R) 1.1, (H) ~ 1	(H) 92 ± 32 , (R) 55.8 ± 11.6
PSZ2 G084.10+58.72	cHalo	0.5	4.2 ± 1.2
PSZ2 G086.93+53.18	Halo	0.6	12.4 ± 3.6
PSZ2 G087.39+50.92	AGN-no diffuse	_	_
PSZ2 G088.98+55.07	AGN-no diffuse	_	_
PSZ2 G089.52+62.34	(A) Relic	(A) 0.23	(A+C) 74.6 ± 15.0
	(B) cRelic (A+B: cdRelic)	(B) 0.17	(B) 12.2 ± 2.6
PSZ2 G095.22+67.41	cRelic	0.4	6.8 ± 1.5
PSZ2 G096.14+56.24	AGN-no diffuse	_	_
PSZ2 G098.44+56.59	AGN-no diffuse	_	
PSZ2 G099.86+58.45	Halo	1.0	14.7 ± 3.2
PSZ2 G106.61+66.71	cHalo	0.5	20 ± 4
PSZ2 G107.10+65.32	dHalo	(Hn) 2.0, (Hs) 1.3	(Hn) 123 ± 25 , (Hs) 63 ± 14
	Relic (s)	0.5	20.9 ± 4.3
	cFossil (n: S1, S2)	(\$1) 0.4. (\$2) 0.23	$(S1) 79 \pm 17, (S2) 17.5 \pm 3.6$
PSZ2 G111.75+70.37	Relic	0.7	(R) 106.7 ± 21.4
	Halo	~ 0.6	27.7 ± 6.3
PSZ2 G114.31+64.89	Halo	0.5	91.7 + 18.6
PSZ2 G114 99+70 36	cHalo cFossil		*
PSZ2 G118 34+68 79	cHalo Fossil	(H) ~ 0.4 (F) 0.5	(H) 28 ± 13 (F) 67 ± 13
PSZ2 G123 66+67 25	cFossil	0.26	75 + 16
PSZ2 G133 60+69.04	Halo cdRelic/cFossil	(H) 0 9	(H) $129 + 26$
PSZ2 G135 17+65 43	cHalo	0.5	29.0 ± 6.6
	cRelic/cFossil	0.4	9 14 + 1 9
PSZ2 G136 92+59 46	AGN-no diffuse		
PSZ2 G143.26+65.24	Halo	1.5	(H) 29.8 + 6.6
PSZ2 G144 33+62 85	AGN-no diffuse		
PSZ2 G145 65+59 30	cHalo/cFossil	0.4	67+17
PSZ2 G150 56+58 32	Halo	1.0	712 + 145
PSZ2 G151 62+54 78	AGN-no diffuse		
PSZ2 G156.26+59.64	cHalo	0.5	7.9 + 3.7
RXC 11053 7+5452	Relic	0.75	$\frac{1}{214 + 43}$
Abell 1156	cHalo	0.75	157 ± 68
Abell 1314	Eossil (central)	0.7	13.7 ± 0.8 136.0 ± 27.4
Abell 1615	AGN no diffuse	0.44	130.9 ± 27.4
MaxBCC 1173 04772 + 47 81041	AGN no diffuse		—
NSC 1142925 ± 462744	Upalagaified	0.4	—
CMDCC 1211 77222 + 55 00068	AGN no diffuso	0.4	—
WILL 1125926 8 + 440111	Helo	<u> </u>	<u> </u>
WHI 1122/18 6 + 400540	Fossil	0.0	30.4 ± 11.7 200 ± 42
WHI 112/1/2 1 + /00510		1.2	ムU2 エ 42 ★
WHI 1122226 8 + 464620	allala	1.2	*
WILL 1122615 8 + 485220	ACN no diffuse	0.3	*
vv пL J132013.0+483229	AGIN-IIO UIITUSE	0.32	

Table 3. Cluster radio properties

cHalo

AGN-no diffuse

WHL J133936.0+484859

WHL J134746.8+475214

H = Halo, R=Relic, d=double, s=South, n=North, LLS=largest linear size * radio emission is blended with other sources and therefore no reliable flux density measurement can be obtained

 ~ 0.3

*

4.1. PSZ2 G080.16+57.65, Abell 2018

Abell 2018 is relatively nearby cluster located at z = 0.0878. The Pan-STARRS gri image reveals a number of central galaxies, without a clear dominant BCG. The cluster has a low PSZ2 mass of $M_{500} = 2.51^{+0.20}_{-0.21} \times 10^{14} \text{ M}_{\odot}$. The high-resolution radio image, see Figure 5, shows a complex central region with several AGN being present as well as more extended emission. A low-surface brightness arc-like structure is found about 1.3 Mpc to the east of the optical center of the cluster. This source has an LLS of about 1.1 Mpc. In a low-resolution image (made with a 90" taper), additional faint diffuse emission spanning the central region of the cluster is found. This emission extends all the way to the arc-like structure to the east. This emission has a size of approximately 1.0 Mpc by 1.8 Mpc. The XMM image reveals an EW merging cluster with the main component located at the optical center of the cluster. We classify the arc-like structure to the east as a giant radio relic and the large-scale centrally placed emission as a candidate radio halo. The nature of the central brighter diffuse emission near the AGN remains unclear. It could be related to the radio halo or be revived fossil plasma from the AGNs. For the candidate radio halo we determine an integrated flux density of $S_{144} = 92 \pm 32$ mJy, where we masked the central brighter diffuse emission around the AGN and the extension towards the radio relic in the fitting. We note that this cluster falls above the correlation between cluster mass and radio power (see Section 5.3). Deeper observations are required to shed more light on this point and on the origin of the extension from the central emission towards the relic.

4.2. PSZ2 G084.10+58.72

The XMM-Newton image of this distant (z = 0.73) cluster shows a disturbed system with a 1 Mpc bar-like structure north of the cluster center, see Figure 6. A hint of centrally located diffuse emission is seen in our image with compact sources subtracted. This emission extends on scales of about 0.5 Mpc. We therefore classify this as a candidate radio halo with $S_{144} = 4.2 \pm 1.2$ mJy. Di Gennaro et al. (2020) also reported a hint of diffuse emission in this cluster.

4.3. PSZ2 G086.93+53.18

Our low-resolution LOFAR image reveals diffuse emission with a largest extent of about 0.6 Mpc in this z = 0.6752 cluster, see Figure 7. This emission is centrally located and we therefore classify it as a radio halo with a flux density of 12.2 ± 3.6 mJy. This is consistent within the uncertainties with the value reported by Di Gennaro et al. (2020) who presented the discovery of a radio halo in this cluster. Yuan & Han (2020) classify the cluster as a disturbed system based on Chandra observations.

4.4. PSZ2 G089.52+62.34, Abell 1904

The XMM-Newton image reveals a disturbed cluster, with the ICM being elongated in the NE-SW direction, see Figure 8. In the LOFAR image we detect two arc-like radio sources to the north (A) and SE (B) of the cluster center. At the redshift of the cluster (z = 0.0701), these sources have an LLS of 230 and 170 kpc, respectively. Sources A and B are located at projected distances to cluster center of 350 and 150 kpc, respectively. Despite these relatively small distances, we consider these sources to be peripheral given the ICM distribution. A much fainter elongated source (C) is visible NW of source A. No optical counter-

parts are detected for these sources. A compact double lobed radio galaxy is located just west of source A. Given the peripheral location of source A with respect to the ICM distribution and elongated ICM shape, we classify A as a radio relic. The location of B is somewhat peculiar with respect to the overall ICM elongation. Given that source B is also affected by calibration artefacts from a nearby bright source, we classify source B as a candidate radio relic. If the classification of source B is confirmed this cluster hosts a double radio relic. The (candidate) relics in Abell 1904 are relatively small compared to other wellstudied relics, although we note that, for example, the western relic in ZwCl 0008.8+5215 also has a small LLS of 290 kpc (Di Gennaro et al. 2019).

4.5. PSZ2 G095.22+67.41, RXC J1351.7+4622

PSZ2 G095.22+67.41 is a relatively nearby cluster located at z = 0.0625. The XMM-Newton image reveals that the ICM peak coincides with the location of the BCG, see Figure 9. A faint X-ray extension is visible to the east. This suggests that the cluster is not fully relaxed and is undergoing a merger event in the EW direction.

No central diffuse radio emission is found in our LOFAR image. However, we detect a NS elongated source about 0.8 Mpc to the east of the cluster center. The source has an LLS of about 0.4 Mpc and is located south of a brighter radio galaxy. We classify the source as a relic tracing a shock that could have originated from the EW merger event. This is consistent with the NS extent, highly elongated shape, and peripheral location of the radio source. An alternative explanation is that this source traces (old) AGN plasma from a bright elliptical galaxy (MCG+08-25-051, z = 0.0623) that is located near this source. A faint (~ 0.5 mJy) compact radio counterpart is detected in our LO-FAR image for MCG+08-25-051. However, there is no evidence of jets or lobes originating from this source which then suggests that the emission resulted from a previous episode of AGN activity.

4.6. PSZ2 G099.86+58.45

PSZ2 G099.86+58.45 is a massive cluster ($M_{500} = 6.85^{+0.48}_{-0.49} \times 10^{14} M_{\odot}$) with a global temperature of $8.9^{+2.8}_{-1.1}$ keV (Sereno et al. 2018). The cluster is known to be undergoing a merger event. The discovery of a 1 Mpc radio halo in this distant (z = 0.616) cluster, see Figure 10, was reported by Cassano et al. (2019) based on LOFAR data. From an elliptical fit we determine $S_{144} = 14.7 \pm 3.2$ mJy. This number somewhat lower than the 25.3 ± 5.7 reported by Cassano et al. (2019). This difference is caused by a small extension of the radio halo around compact source B (discussed by Cassano et al. (2019)). If this region is included, the measurements are consistent with each other given the uncertainties.

4.7. PSZ2 G106.61+66.71

The LOFAR image displays two bright, partly blended tailed radio galaxies in the western part of the cluster, see Figure 11. The Chandra X-ray image shows a cluster without a strongly peaked core. In addition, the cluster is elongated in the direction of the two tailed radio galaxies. In the center of the cluster we detect diffuse radio emission with an LLS of about 0.5 Mpc. Within this diffuse emission we find two compact AGN associated with the cluster, including the BCG. We consider a mini-halo origin



Fig. 5. PSZ2 G080.16+57.65 / Abell 2018. Left: Low-resolution 144 MHz radio image made with a 30" taper. Middle: XMM-Newton image with radio contours from a 90" tapered image overlaid. Compact sources were subtracted in this image. Right: Pan-STARRS *gri* color image overlaid with radio contours from the robust -0.5 weighted image. The cluster center position is marked with a white 'X'. Radio contours on both the X-ray and optical images are drawn at $3\sigma_{rms} \times [1, 2, 4, ...]$, with σ_{rms} the r.m.s. map noise.



Fig. 6. PSZ2 G084.10+58.72. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 15" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 7. PSZ2 G086.93+53.18. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 15" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

unlikely given the lack of centrally peaked X-ray emission. The western part of the central diffuse emission has a rather high surface brightness, suggesting a link with a cluster AGN. On the other hand, some of the emission (in particular the eastern part) has a smooth morphology somewhat similar to a radio halo. For the above reasons we list the source as candidate radio halo.

4.8. PSZ2 G107.10+65.32, Abell 1758

Abell 1758 is composed of two main clusters, Abell 1758N and Abell 1758S, see Figure 12. These two clusters appear to be in a pre-merger phase (Rizza et al. 1998; David & Kempner 2004).

The individual clusters themselves are disturbed and already undergoing their own merger events. The presence of a radio halo in A1758N was first reported by Kempner & Sarazin (2001) and further studied by Giovannini et al. (2009) and Venturi et al. (2013).

Botteon et al. (2018) studied this cluster with LOFAR HBA observations, a subset of the data used here. They measured a size of about 2 Mpc for the A1758N radio halo and discovered a 1.6 Mpc radio halo and a ~ 0.5 Mpc relic (labelled R) in A1758S. In addition, a hint of faint emission was detected between A1758N and A1758S. This "radio bridge" was subsequently confirmed by Botteon et al. (2020c). Two bright patches,



Fig. 8. PSZ2 G089.52+62.34 / Abell 1904. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 9. PSZ2 G095.22+67.41, RXC J1351.7+4622. Left: XMM-Newton X-ray image with 30" tapered radio contours (compact sources were removed). Right: Optical image with Robust –0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 10. PSZ2 G099.86+58.45. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 30" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

labelled S1 and S2, were found in A1758N. They are not directly associated with an optical counterpart and possibly trace compressed AGN fossil plasma.

For the radio halo in A1758N we find a flux of 123 ± 25 mJy based on the elliptical model fit. This value is lower than the 307 ± 63 mJy measured earlier by Botteon et al. (2020c). The

reason for this difference is that we excluded the region around sources A and B (see Botteon et al. 2018) and S1 and S2. This particular case highlights the difficulties in classifying and separating the emission from different components with uncertain identifications. For the southern radio halo we determine $S_{144} = 63 \pm 14$ mJy from the elliptical model, consistent with



Fig. 11. PSZ2 G106.61+66.71. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

the value reported by Botteon et al. (2018). For a more comprehensive description of the LOFAR findings we refer the reader to Botteon et al. (2018, 2020c).

4.9. PSZ2 G111.75+70.37, Abell 1697

The XMM-Newton X-ray image shows an elongated cluster without a central concentration. The LOFAR images reveal two prominent radio sources in this cluster, see Figure 13. One is located near the southwestern BCG of the cluster. The second source is more extended and located at the NE periphery of the cluster. This emission is also discussed by Paul et al. (2020) based on LoTSS DR1 images (Shimwell et al. 2019). The source consists of a NW-SE elongated structure with an LLS of about 700 kpc, as well as emission trailing SW towards a second BCG. Compact radio emission from this BCG is detected, but no obvious morphological connection with the extended source is visible. The extended source somewhat resembles the Toothbrush relic (van Weeren et al. 2012). We note that the extended radio source partly overlaps with the nearby irregular dwarf galaxy UGC 8308 (DDO 167; e.g., Tikhonov & Karachentsev 1998). However, we consider an association of the radio source with the cluster more likely. We therefore classify the source as a radio relic. This classification is also consistent with the elongation of the ICM, suggesting a NE-SW merger event. In the low resolution radio images extended emission is visible in the region between the NE and the SW BCG. This emission is classified as a radio halo since it follows the X-ray emission from the ICM. The flux density of the halo is 27.7 ± 6.3 mJy. We note that this emission is also discussed by Paul et al. (2020) ("trailing relic emission") and it is concluded that this emission has an ultrasteep spectrum. We note however that the flux density estimate provided by Paul et al. (2020) is incorrect as the emission is not fully deconvolved in the LoTSS DR1 images and thus measurements directly from those images are complicated. This leads to a large overestimation of the flux density as is discussed by Shimwell et al. (2019). This problem is addressed to some extent in the upcoming DR2.

4.10. PSZ2 G114.31+64.89, Abell 1703

The massive cluster Abell 1703 has been extensively studied at optical wavelengths due to its gravitational lensing properties. It is generally classified as a relaxed cluster (e.g., Umetsu et al. 2011; Richard et al. 2009). The Chandra image shows an overall regular appearance with the cluster being somewhat elongated

in the NNW-SSE direction, see Figure 14. Andrade-Santos et al. (2017) determined the concentration parameters, cuspiness, and central density of the cluster using Chandra X-ray data. These properties, however, point to a non-cool core classification. Furthermore, a recent dynamical analysis from Boschin et al. (2020) indicates that this is a merging cluster consisting of two or three subclumps.

The LOFAR image reveals the presence of central diffuse emission, filling the region around between tailed radio source A and a complex compact region of emission which we labelled B. Source B is likely related to AGN activity. A faint tailed source, labelled C, is located in the SE part of the cluster. The total extent of the diffuse emission is difficult to determine because of A and B. Given the central location and extent of at least 0.5 Mpc, we classify the diffuse emission as a radio halo with a integrated flux density of 91.7 \pm 18.6 mJy based on an elliptical model fit.

4.11. PSZ2 G114.99+70.36, Abell 1682

The presence of diffuse emission in this cluster was first reported by Venturi et al. (2008) and subsequently studied in more detail by Venturi et al. (2011); Macario et al. (2013); Venturi et al. (2013). LOFAR observations show complex diffuse extended emission extending over a region of more than 1 Mpc, see Figure 15. The Chandra X-ray image displays a disturbed ICM indicating that the cluster is undergoing a merger event. The LOFAR results have been presented in Clarke et al. (2019). Some of the structures in the cluster are related to distorted and tailed radio galaxies. A few regions of the diffuse emission show a steep spectral index of $\alpha \sim -2$. These regions could trace AGN fossil plasma that has been re-accelerated (or revived) by shocks and turbulence related to the ongoing merger event. A candidate radio halo is located near the peak of the X-ray emission, labelled (CH). The complexity of the various radio structures in the vicinity prevent us from measuring its properties.

4.12. PSZ2 G118.34+68.79

Several radio galaxies and diffuse emission are detected in this cluster, see Figure 16. An optical image shows two BCGs that are located along a SE-NW axis. The cluster hosts a tailed radio galaxy, labeled A. To the east, a bright patch of emission (B) is found just above the SE BCG. Additional fainter emission is located around B. Given the high surface brightness of B and several nearby radio AGN, B is likely AGN plasma, possibly revived by the passage of a shock.



Fig. 12. PSZ2 G107.10+65.32 / Abell 1758. Left: Robust –0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours. Right: Optical image with Robust –0.5 image radio contours. For more details see the caption of Figure 5.

We also find low-level diffuse emission extending on scales of about 0.4 Mpc in the central regions of the cluster, labelled H. However, it is hard to determine its full spatial extent as it partly blends with other extended radio sources in this region. Since the diffuse emission approximately follows the overall galaxy distribution and has a low surface brightness we classify it as a candidate radio halo. Fitting a circular model we estimate $S_{144} =$ 28 ± 13 mJy, where we masked the region near B and west of A.

4.13. PSZ2 G133.60+69.04, Abell 1550

The existence of diffuse emission in Abell 1550 (z = 0.2540) cluster was first reported by Govoni et al. (2012). Extended radio emission is clearly detected in this cluster by LOFAR. It extends over a significantly larger region than found by Govoni et al., corresponding to an LLS of 1.8 Mpc. This is explained by the shallower depth of the VLA image used Govoni et al. and/or a steep radio spectrum.

The Chandra image shows a roughly roundish ICM distribution. Additional faint X-ray emission is observed NE of the main structure. The main part of the diffuse radio emission, labeled H, traces the X-ray emission from the ICM. We therefore classify it as a giant radio halo with a size of about 0.9 Mpc. Based on the elliptical model fit, we obtain a radio halo integrated flux density of 129 ± 26 mJy. The radio images also show a bright tailed radio galaxy (A) and several structures in the western part of the cluster (D, E, C) that do not have clear optical counterparts. A diffuse patch of emission, labeled B, is placed near a group of galaxies located at the same redshift as the main cluster. A lower resolution radio image shows that B is connected to the main radio halo. Source B seems to be associated with the NE X-ray extension. Sources E, D, and C look to be relics, possibly related to revived AGN fossil plasma. We note that both D and C are connected to the emission from the main radio halo. Based on the extension of the ICM and placement of the diffuse radio sources, the cluster seems to have undergone a merger event in the NE-SW direction. The location of B in the NE, and C-D-E in the SW could indicate that these structures are related to mergers shocks. If this interpretation is correct the cluster hosts a double radio relic. The combination of a double radio relic and a radio is relatively rare (Bonafede et al. 2017).

4.14. PSZ2G135.17+65.43

The LOFAR image uncovers a large number of radio galaxies in this z = 0.5436 cluster which are labeled A to F, see Figure 18. One of these, source E, has a physical extent of about 1 Mpc and we classify it as a giant double-double. Sources A to D display head-tail morphologies. Source G is a peripheral source with an extent of about 400 kpc. We do not identify an optical counterpart for this source and tentatively classify it as a relic or an AGN fossil plasma source. The Chandra image of PSZ2 G135.17+65.43 indicates a non-relaxed cluster without a clear central peak. We also detect central diffuse emission in this cluster with a total extent of about 500 kpc, labeled H. Due to the presence of the tailed radio galaxies A and B, the full extent of H is hard to determine but the emission approximately follows the thermal ICM. Given its central location and extent, we classify H as a candidate radio halo with a flux density of $S_{144} = 29.0 \pm 6.6$ mJy based on the circular model fit.

4.15. PSZ2 G143.26+65.24, Abell 1430

The LOFAR images reveal the presence of diffuse emission in this cluster which is located at z = 0.3634, see Figure 19. The LOFAR observations are discussed in more detail in Hoeft et al. (2020), so we only give a brief overview of the main findings. The diffuse emission extends over a region of about 5' by 2.5', with the emission being elongated in the EW direction. The angular extent corresponds to physical size of 1.5 Mpc by 0.8 Mpc. Our LOFAR image also reveals two tailed radio galaxies, labelled A and B (Figure 19). An optical image displays two subclusters. The western subcluster corresponds to the main structure seen in the Chandra X-ray image, while the eastern cluster is much fainter in the Chandra image. The radio emission spans the full region between the main western and the smaller eastern subcluster. Given the clear correspondence between the radio and X-ray emission of the western subcluster and the large extent of the diffuse emission, we classify the western part of the diffuse radio emission as a giant radio halo. We measure $S_{144} = 29.8 \pm 6.6$ mJy for the radio halo from our fitting. We note that this value might be affected by the diffuse emission around the eastern subcluster since it partly overlaps with the radio halo from the western subcluster. The nature of the diffuse emission around the eastern subcluster is not fully clear but it could be a radio bridge (for a discussion on this see Hoeft et al. 2020). The disturbed character of the cluster, both in optical and X-ray images, indicates that the cluster is undergoing a merger event, consistent with the presence of the giant radio halo. A possible source of seed electrons for the radio halo could be the tail of source A, since it blends into the halo emission.



Fig. 13. PSZ2 G111.75+70.37 / Abell 1697. Top Left: Robust -0.5 radio image. Top right: 10" tapered radio image. Bottom left: XMM-Newton X-ray image with 30" tapered radio contours (compact sources were subtracted). Bottom right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

4.16. PSZ2 G145.65+59.30, Abell 1294

An XMM-Newton image shows a cluster that is elongated in the EW direction, see Figure 20. The galaxy distribution shows a similar elongation and extending all the way to a tailed radio galaxy A. Our LOFAR image reveals faint diffuse emission, with an LSS of about 0.4 Mpc, located at the western part of the cluster. This emission is very faint and close to the detection limit of our observations. A candidate tailed radio galaxy (B) is located just east of the diffuse radio source. The diffuse emission is difficult to classify with the current data in hand. A possibility is that it is related to revived or re-accelerated fossil plasma from source B. On the other hand, it could also be a radio halo, somewhat similar to the "off-axis" radio halo found in Abell 1132 (Wilber et al. 2018). The integrated flux density of the diffuse source is $S_{144} = 6.7 \pm 1.7$ mJy, based on the circular model fit.

RA (J2000)

4.17. PSZ2 G150.56+58.32, MACS J1115.2+5320

MACS J1115.2+5320 is a massive merging cluster located at z = 0.466. A Chandra X-ray image reveals an ongoing merger event along a NW-SE axis, see Figure 21. Mann & Ebeling (2012) classified this system as a possible head-on binary cluster merger. The global temperature of the cluster was measured to be 8.6 ± 1.1 keV by Morandi et al. (2015).

RA (J2000)

The LOFAR image shows a long 770 kpc tailed radio galaxy (A), with the tail pointing NW. Two additional tailed radio galaxies (B, C) have their tails pointing to the SE. Source D is located just SE of the head-tail source A and has a rather complex morphology in our high-resolution image. Near source D, a possible merger induced cold front is visible in the Chandra image. Low surface brightness radio emission is detected in the region north of source A. This emission extends over 1 Mpc and given its size we classify it as a giant radio halo. Fitting an elliptical model, we determine $S_{144} = 71.2 \pm 14.5$ mJy for the radio halo.

A&A proofs: manuscript no. aa



Fig. 14. PSZ2 G114.31+64.89 / Abell 1703. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 15" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 15. PSZ2 G114.99+70.36 / Abell 1682. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 16. PSZ2 G118.34+68.79. Left: Robust -0.5 radio image. Middle: 15" tapered radio image with compact radio sources subtracted. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

4.18. PSZ2G156.26+59.64

A bright compact radio source is detected in this z = 0.6175 cluster which seems to be associated with a BCG, see Figure 22. In our low-resolution images, with compact sources removed, faint diffuse emission is detected in the region above the compact radio source. Only a hint of this emission is detected in our high-resolution image. This diffuse emission has a total extent of about 0.5 Mpc. Given its approximate central location, with respect to the cluster member galaxies as judged from the Pan-STARRS images, and large physical size, we classify this source as a candidate radio halo. For the source we determine a flux density of 7.9 ± 3.7 mJy.

4.19. RXC J1053.7+5452

Peripheral diffuse radio emission in this cluster was first reported by Rudnick & Lemmerman (2009) and subsequently studied by van Weeren et al. (2011). Chandra and Suzaku observations were presented by Itahana et al. (2017) showing the cluster is undergoing a merger event. The peripheral radio source is classified as a relic.

The main LOFAR pointing on this source (P164+55) had to be discarded as it was affected by bad ionospheric conditions. Hence the noise levels in our images are higher than for other clusters. Despite the higher noise, the relic is clearly detected in our LOFAR images (Figure 23) and the source has a similar appearance as in the WSRT observations presented in van Weeren



Fig. 17. PSZ2 G133.60+69.04 / Abell 1550. Top left: Robust -0.5 radio image. Top right: 10" tapered radio image. Bottom left: Chandra X-ray image with 10" tapered radio contours. Bottom right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

et al. (2011). In the LOFAR image, the relic has an LLS of about 0.75 Mpc. A hint of an extension is visible from the northern tip of the relic (near a compact source) towards the west and north. Combining the LOFAR flux density measurement with the one obtained from the WSRT, we obtain $\alpha = -1.17 \pm 0.11$, which is typical for radio relics.

4.20. Abell 1156

A Chandra X-ray image displays a cluster that is elongated in the NS direction (Figure 24). Diffuse radio emission, also elongated in the NS direction, is detected in this cluster with an LLS of 0.7 Mpc. This radio emission does not peak at the cluster center but south of it. However, some faint diffuse emission is also visible north of the cluster center. We list this source as a candidate radio halo, originating from a possible NS merger event. Additional extended radio emission, with a mostly EW elongation is found in the southern periphery of the cluster, its origin is not fully clear but it might be related to AGN activity as a connection to a cluster member galaxy is suggested. The cluster also hosts a prominent head-tail radio source in the north, with the tail extending south. along the direction of the proposed merger axis. By fitting the elliptical model we determine $S_{144} = 15.7 \pm 6.8$ mJy for the candidate radio halo.

4.21. Abell 1314

This nearby low-mass cluster hosts a bright 800 kpc long tailed radio galaxy associated with the galaxy IC 711. The main tails shows a range of complicated linear features, see Figure 25. LOFAR observations of this source are described Wilber et al. (2019). GMRT observations have been presented by Srivastava & Singal (2016); Sebastian et al. (2017). Another smaller bright tailed radio galaxy is associated with IC 708. In addition, a filamentary source is detected near the cluster center. This is a candidate radio phoenix related to the central BCG IC 712. Our new



Fig. 18. PSZ2 G135.17+65.43. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

LOFAR images of Abell 1314 reveal some additional details not visible in the previous LOFAR images presented in Wilber et al. (2019) due to the improved calibration. One of these is a thin elongated structure that connects IC 708 to IC 711. Its origin is unclear. In addition, our images show more clearly the filamentary nature of the phoenix source, with an LLS of 0.44 Mpc.

4.22. NSC J143825+463744

The LOFAR observations for NSC J143825+463744, a nearby z = 0.03586 system, are affected by bad ionospheric conditions. Despite of the poor image quality, central extended emission is detected with a size of about~ 0.4 Mpc, see Figure 26. NSC J143825+463744 was classified as a galaxy group (MLCG 1495) by Gal et al. (2003). It is composed of two dominant galaxies NGC 5722 and NGC 5717. No X-ray emission is detected from this system by ROSAT, confirming it has a low mass. Given the lack of an ICM detection by ROSAT, low mass of the system, and poor calibration, we list it as unclassified.

Note that a small EW extended source, right of the image center, is associated with NGC 5714. NGC 5714 is foreground galaxy and unrelated to MLCG 1495.

4.23. WHL J122418.6+490549

WHL J122418.6+490549 hosts a bright elongated radio source with an LLS of 370 kpc (Figure 27). This radio emission could have originated from the BCG (LEDA 2333420), which is located at the southeastern tip of the elongated source. The cluster is barely detected in a Chandra observation, indicating a low-mass system. The source seems somewhat similar to the revived remnant radio lobe found in low-mass cluster Abell 1931 (Brüggen et al. 2018).

4.24. WHL J124143.1+490510

Faint patchy extended emission is detected in our low resolution images of this cluster, see Figure 28. This emission extends over a region of 1.2 Mpc. Given that the emission is approximately centered on the galaxy distribution we classify it as a candidate radio halo.

Article number, page 20 of 35

4.25. WHL J132226.8+464630

Patchy diffuse radio emission with an extent of about 0.5 Mpc is detected west of the BCG (Figure 29). Given its approximate central location we classify it as a candidate radio halo.

4.26. WHL J133936.0+484859

Extended radio emission is detected in this cluster south of the BCG, see Figure 30. This emission seems to be the extension of a tailed radio galaxy. Additional extended (~ 300 kpc) emission surrounds the BCG. This candidate radio (mini-)halo will need to be confirmed with deeper observations.

4.27. WHL J125836.8+440111

In WHL J125836.8+440111 we find extended radio emission with an LLS of 800 kpc, see Figure 31. The northern part of the emission seems to originate from two radio galaxies. However, the emission extends further south, all the way to a distorted double lobed source. Given the large area that is covered by the extended emission we classify part of this emission as a radio halo.

5. Discussion

5.1. Classification of diffuse radio emission

Our LOFAR observations emphasize the challenge of classifying diffuse cluster radio sources in sensitive low-frequency images. In particular, the distinction between diffuse sources and (old) AGN radio plasma is not always clear. The reason for this is that old AGN plasma also has a steep radio spectrum so this plasma brightens significantly in low frequency images. For example, this manifests itself by the longer tails of tailed radio galaxies. Several of the clusters in our sample show extended AGN emission with an LLS of more than 0.5 Mpc (e.g., PSZ2 G114.99+70.36, PSZ2 G135.17+65.43, Abell 1314, and GMBCG J211.77332+55.09968, see Appendix A.10). AGN with an LLS of a few hundred kpc are even more common. When the emission from tailed and distorted AGN starts to overlap with relics and halos the classification of diffuse emission becomes more difficult.

A second reason for that classification is challenging is related to the origin of the diffuse radio emission itself. Particle re-acceleration models require a source of (mildly) relativistic electrons. The LOFAR images reveal that AGNs provide a rich



Fig. 19. PSZ2 G143.26+65.24 / Abell 1430. Top left: Robust -0.5 radio image. Top right: 10" tapered radio image. Bottom left: Chandra X-ray image with 15" tapered radio contours (compact sources were subtracted). Bottom right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

source of fossil electrons and therefore the observational distinction between plasma directly accelerated by the AGN or in-situ re-accelerated in the ICM becomes more difficult. Spectral measurements and a comparison with the thermal ICM properties will therefore be important to correctly classify the extended radio emission.

5.2. Projections for the completed LoTSS survey.

Based on the number of detected radio halos and relics in the HETDEX area, we can make a prediction for the number of radio halos and relics that will be detected in the completed LoTSS survey. We assume here that the LoTSS survey will have uniform sensitivity over the entire northern sky. We ignore any complications due to differences in the mass and redshift distributions between the PSZ2 clusters in the HETDEX DR1 area in the completed LoTSS survey. The uncertainties are computed based on Poisson statistics.

We first focus on the radio halos in the Planck PSZ2 clusters. We detect a total of 8 radio halos and 8 candidate halos in this sample of 26 clusters (where we count PSZ2G107.10+65.32, consisting of Abell 1758N and Abell 1758S, only once). The number of PSZ2 clusters without any diffuse emission is 7. We should note that for three of these clusters the noise levels were higher because of bad ionospheric conditions. Four PSZ2 clusters host radio relics. The fraction of confirmed radio halos in PSZ2 clusters is $31 \pm 11\%$. Considering that there are 595 PSZ2 clusters above a declination of 0 degrees with confirmed redshifts, we estimate that there will be 183 ± 65 radio halos detected in the LoTSS survey in PSZ2 clusters. Also including candidate radio halos, the detection fraction is $62 \pm 15\%$. The number of PSZ2 halos in the LoTSS survey then becomes 366 ± 92 . The fraction of clusters with some form of diffuse emission is $73 \pm 17\%$. This suggests that 435 ± 99 PSZ2 clusters with diffuse emission will be detected in LoTSS.



Fig. 20. PSZ2 G145.65+59.30 / Abell 1294. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 21. PSZ2 G150.56+58.32 / MACS J1115.2+5320. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours (compact sources were subtracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 22. PSZ2 G156.26+59.64. Left: Robust -0.5 radio image. Middle: 15" tapered radio image with compact sources subtracted. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

We detect one radio halo in the non-PSZ2 clusters and four candidate radio halos. Considering these numbers, and that the region we surveyed covers 424 deg², we expect to find at least 10^2 radio halos in non-PSZ2 clusters in the full LoTSS. Here we assumed that about half of the candidate radio halos are real. The total number of radio halos (PSZ2 plus non-PSZ2) that will be found in the full LoTSS survey will thus be around 400–500, again assuming that half of the candidates are real.

The number of radio halos that our extrapolation predicts for the LoTSS survey agrees with that predicted from reacceleration models (Cassano et al. 2010a, 2012). These models predict that a significant fraction of halos have steep spectra, especially in clusters with smaller masses and higher redshift, and that the occurrence of halos in clusters declines with cluster mass. The large statistics that is expected from LoTSS will allow to test the dependence of the occurrence of radio halos with cluster mass and redshift.

Interestingly, we did not detect clear examples of radio minihalos. Radio mini-halos are more difficult to classify due to their smaller sizes and the sample studied here is not known to host any strong cool-core clusters. Therefore, the lack of radio minihalos might be (partly) related to the properties of our sample. However, there is no consensus that SZ-selected samples are biased against cool-core clusters Eckert et al. (2011); Rossetti et al. (2017); Andrade-Santos et al. (2017). The lack of mini-halos in our sample could also be a reflection of the properties of mini-



Fig. 23. RXC J1053.7+5452. Left: Robust -0.5 radio image. Right: Chandra X-ray image with 10" tapered radio contours. For more details see the caption of Figure 5.



Fig. 24. Abell 1156. Left: 15" tapered radio image. Middle: Chandra X-ray image with 30" tapered radio contours (compact sources were sub-tracted). Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 25. Abell 1314. Left: Robust -0.5 radio image. Middle: XMM-Newton X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

halos at low frequencies. Several low-frequency studies have found the presence of diffuse emission at larger radii, beyond the extent of the mini-halos measured at GHz frequencies. Thus, low-frequency studies would be less likely to report mini-halos if size is used as a criterion. Whether this extended emission beyond the "classical" mini-halo extent can be considered as a part of the mini-halo, or is an unrelated component more similar to that of a giant radio halo, is unclear (Bonafede et al. 2014b; Brunetti & Jones 2014; Kale & Parekh 2016; Venturi et al. 2017; Savini et al. 2018, 2019; Kale et al. 2019).

A further complication for the detection is mini-halos is the presence of a radio bright BCG. The calibration needs to achieve



Fig. 26. NSC J143825+463744. Left: Robust -0.5 radio image. Right: 30" tapered radio image. For more details see the caption of Figure 5.



Fig. 27. WHL J122418.6+490549. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10'' tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 28. WHL J124143.1+490510. Left: 30" tapered radio image with compact sources subtracted. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.





Fig. 29. WHL J132226.8+464630. Left: 15" tapered radio image with compact sources subtracted. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. 30. WHL J133936.0+484859. Left: 10" tapered radio image. Middle: Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

a dynamic range sufficient for a detection. This can be a challenge at low frequencies, see for example Figure A.3.

The fraction of radio relic hosting clusters in PSZ2 is $15 \pm 8\%$. We thus expect 92 ± 46 clusters hosting radio relics in LoTSS. One radio relic is detected in a non-PSZ2 cluster. The expected number of radio relics in the LoTSS survey falls about an order of magnitude below the prediction of Nuza et al. (2012). Predictions are not available yet for more recent models (e.g., Nuza et al. 2017; Brüggen & Vazza 2020).

5.3. P_{150 MHz} scaling relations

It is well established that the radio power of giant halos scales with the X-ray luminosity of clusters (e.g., Liang et al. 2000; Cassano et al. 2006, 2013; Kale et al. 2015). A similar scaling

exists for the integrated Compton Y parameter which traces the ICM integrated pressure along the line of sight (Basu 2012; Cassano et al. 2013). Both the X-ray and SZ measurements are proxies of clusters mass. Therefore the suggestion is that the observed correlations originate from an underlying relation between cluster mass and radio power. The explanation for this relation is that a fraction of the gravitational energy released during a merger event, which scales with cluster host mass, is channeled into the re-acceleration of cosmic rays via turbulence (e.g., Cassano & Brunetti 2005; Cassano et al. 2004).

Traditionally, the radio power is computed at a rest-frame frequency of 1.4 GHz ($P_{1.4 \text{ GHz}}$) and all correlation studies so far have used this quantity. With the new LOFAR radio halo detections it becomes feasible to study this relation at a rest-frame frequency of 150 MHz, with $P_{150 \text{ MHz}}$ given by



Fig. 31. WHL J125836.8+440111. Left: 10" tapered radio image with compact sources subtracted. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

$$P_{150 \text{ MHz}} = \frac{4\pi D_L^2 S_{150 \text{ MHz}}}{(1+z)^{\alpha+1}},$$
(4)

with D_L the luminosity distance of the cluster. Because our integrated flux density measurements are obtained at 144 MHz, $P_{150 \text{ MHz}}$ is only marginally affected by the adopted (unknown) radio spectral index.

In this work, we determine the $M_{500}-P_{150 \text{ MHz}}$ scaling relation for radio halos using clusters from the PSZ2 catalog which provides an SZ-based mass estimate. SZ-based mass selected samples should be less affected by the cluster's dynamical state compared to X-ray selected sample which are biased towards relaxed cool-core clusters (e.g., Eckert et al. 2011; Rossetti et al. 2017; Andrade-Santos et al. 2017). We complement our new LO-FAR measurements with halo detections in the 120-180 MHz range from the literature when reliable measurements are available, see Table 5.3. These literature values come mostly from previous LOFAR and GMRT studies. A few of them are taken from Murchison Widefield Array (MWA) studies. We do not include all MWA detections, only those with high-quality measurements, as indicated from good spectral fits with low (< 2) reduced χ^2 values (George et al. 2017), and from halos not significantly affected by the uncertainties from compact source subtraction.

Following Cassano et al. (2013) we use the following relation

$$\log_{10}\left(\frac{P_{150 \text{ MHz}}}{10^{24.5} \text{WHz}^{-1}}\right) = B \log_{10}\left(\frac{M_{500}}{10^{14.9} \text{M}_{\odot}}\right) + A$$
(5)

between radio power and cluster mass. The best fitting parameters are found using the BCES orthogonal regression algorithm (Akritas & Bershady 1996; Nemmen et al. 2012). Past work has adopted by default BCES bisector fits (which give consistently flatter slopes), although, as is mentioned by Hogg et al. (2010), bisector fits are not recommended. For comparison with previous work we also report the results from the BCES bisector and Y|X (with X the independent variable) fits. Our sample approximately covers the mass range $3 - 10 \times 10^{14} M_{\odot}$. The results are shown in Figure 32 (left panel) and Table 5.3. As is the case for $P_{1.4 \text{ GHz}}$, there is also a clear correlation between between M₅₀₀ and $P_{150 \text{ MHz}}$ and we find a slope of $B = 6.13 \pm 1.11$ (BCES orthogonal). With the candidate halos included, we find somewhat flatter slopes (see Table 5.3), but the differences are not significant considering the uncertainties.

We determine the $Y_{500}-P_{150 \text{ MHz}}$ radio halo scaling relation using the PSZ2 Compton Y parameter. For that we convert the $Y_{5R_{500}}$ values from Planck Collaboration et al. (2016) to Y_{500} using $Y_{500} = 0.56 Y_{5R_{500}}$ (Arnaud et al. 2010). We also convert the units from arcmin² to Mpc² to facilitate the comparison with Cassano et al. (2013). We apply the same fitting methods as for the $M_{500}-P_{150 \text{ MHz}}$ correlation, using a relation of the form

$$\log_{10}\left(\frac{P_{150 \text{ MHz}}}{10^{24.5} \text{ WHz}^{-1}}\right) = B \log_{10}\left(\frac{Y_{500}}{10^{-4} \text{Mpc}^2}\right) + A .$$
(6)

The results are plotted in Figure 32 (right panel) and are reported in Table 5.3. The slope of the best fit BCES orthogonal relation is $B = 3.32 \pm 0.65$. The slopes are slightly flatter when including the candidate radio halos, but the differences are not significant considering the uncertainties (see Table 5.3).

The slopes of the 150 MHz scaling relations we find are steeper than the ones obtained at 1.4 GHz by Cassano et al. (2013). For $Y_{500}-P_{150 \text{ MHz}}$ they reported $B = 2.28 \pm 0.35$ from at 1.4 GHz, compared to $B = 3.32 \pm 0.65$ in this work. For $M_{500}-P_{150 \text{ MHz}}$ they reported a slope of $B = 4.51 \pm 0.78$, compared to our value of $B = 6.13 \pm 1.11$ in this work. However, considering the uncertainties, the slopes at 150 MHz are still consistent with the ones reported at 1.4 GHz.

Statistical models employing turbulent re-acceleration (Cassano et al. 2013) predict $P_{1.4\,\text{GHz}} \propto M_{500}^4$, or steeper depending on the ICM magnetic field strength with respect to the equivalent magnetic strength of the cosmic microwave background. It should be noted that these models make a number of simplifying assumptions. The most important simplification is that there is no spatial dependence of the magnetic field and acceleration efficiency, see for example Cassano et al. (2010a) about these "homogeneous models". Cassano (2010) predicts that the slope of the L_X - $P_{1.4\,\text{GHz}}$ scaling relation should steepen by about 0.4

Table 4. Literature	sample fo	r P _{150 MHz}	scaling relations.
---------------------	-----------	------------------------	--------------------

cluster	alternative name	redshift	flux density	frequency	reference
			mJy	MHz	
PSZ2G108.17-11.56	PSZ1G108.18-11.53	0.336	124 ± 11	147	de Gasperin et al. (2015)
PSZ2 G057.80+88.00	Coma Cluster	0.0231	7200 ± 800	150	Cordey (1985)
PSZ2G149.22+54.18	Abell 1132	0.1351	178 ± 27	144	Wilber et al. (2018)
PSZ2 G058.29+18.55	RXC J1825.3+3026	0.065	163 ± 47	144	Botteon et al. (2019)
PSZ2G151.19+48.27	Abell 959	0.2894	94 ± 14	143	Bîrzan et al. (2019)
PSZ2 G071.39+59.54	RXC J1501.3+4220	0.2917	20.2 ± 2.0	144	Wilber et al. (2019)
PSZ2 G139.62+24.18	PSZ1 G139.61+24.20	0.2671	30 ± 4	144	Savini et al. (2018)
PSZ2 G049.22+30.87	RX J1720.1+2638	0.164	165 ± 25	144	Savini et al. (2019)
PSZ2 G138.32-39.82	RXC J0142.0+2131	0.280	32 ± 6	144	Savini et al. (2019)
PSZ2 G226.18+76.79	Abell 1413	0.143	40 ± 7	144	Savini et al. (2019)
PSZ2 G055.59+31.85	Abell 2261	0.224	165 ± 25^{a}	144	Savini et al. (2019)
PSZ2G180.25+21.03	MACS J0717.5+3745	0.546	370 ± 60	147	Bonafede et al. (2018)
PSZ2 G195.75-24.32	Abell 520	0.201	229.7 ± 34.8	145	Hoang et al. (2019a)
PSZ2G100.14+41.67	Abell 2146	0.232	19.8 ± 5.0^{b}	144	Hoang et al. (2019b)
PSZ2G186.37+37.26	Abell 697	0.282	135 ± 27	153	Macario et al. (2013)
PSZ2 G208.80-30.67	Abell 521	0.247	328 ± 66	153	Macario et al. (2013)
PSZ2 G008.94-81.22	Abell 2744	0.308	415 ± 42	150	George et al. (2017)
PSZ2 G175.69-85.98	Abell 141	0.230	110 ± 11	168	Duchesne et al. (2017)

^a No flux density uncertainty provided, adopted an uncertainty of 15%.

^b A range of flux densities was provided for the radio halo by Hoang et al. (2019b). Adopted the average value of that range, with the uncertainty reflecting the provided range.

due to the ultra-steep spectrum halos visible at low frequencies associated with intermediate mass galaxy clusters. Using the fact that $L_{\rm X} \propto M_{500}^2$, we thus expect a steepening of about 0.8 for the $M_{500}-P_{150~\rm MHz}$ scaling relation. In addition, the scatter around the scaling relations should increase. Interestingly, the slopes we find at 150 MHz are indeed steeper than at 1.4 GHz, in line with this prediction. That said, the uncertainties one the determined slopes are still too large for any firm claim. For this reason, extending the sample size and mass range will be crucially important to confirm this result.

One of the limitations of our presented analysis is that due to the small sample size we did not apply any cut in mass or redshift. For a more detailed comparison with the Cassano et al. (2013), a similar redshift and mass cut should be applied. We also note that our sample contains three clusters (Abell 1413, PSZ1 G139.61+24.20 and RX J1720.1+2638) with diffuse sources that were previously classified as mini-halos. Since the diffuse emission in these clusters turns to be more extended⁴ at low frequencies than the typical scale of mini-halos, we have included them in our sample. Further investigations are also required of how the low-frequency scaling relations and occurrence rates depend on the cluster dynamical state. Another limitation of our derived scaling relations is the inclusion of literature values that were obtained in a variety of ways, adopting different methods to measure the integrated flux density. That said, the literature values do not seem to be strongly biased either high or low when compared to the measurements obtained in this work for clusters with similar masses, and hence this should not have a large effect on the derived slopes. This limitation will be removed in future work when larger LOFAR samples will become available.

A comparison between the 150 MHz and 1.4 GHz relations allows use to investigate the average spectral index of radio halos in this frequency range. We find that when using $\alpha = -1.2$ a good

Table 5. BCES fitted scaling relations for radio halos.

mathad	D		Δ	-
method	D	O_B	A	O_A
$M_{500} - P_{150 \text{ MHz}}$				
orthogonal	6.13	1.11	1.22	0.12
bisector	4.67	0.62	1.11	0.09
Y X	3.84	0.69	1.05	0.09
$M_{500}-P_{150 \text{ MHz}}$ with candidates				
orthogonal	5.00	0.87	1.21	0.10
bisector	4.05	0.55	1.11	0.08
Y X	3.40	0.53	1.04	0.09
Y ₅₀₀ -P _{150 MHz}				
orthogonal	3.32	0.65	0.74	0.12
bisector	2.60	0.53	0.74	0.09
Y X	2.10	0.67	0.74	0.08
Y_{500} – $P_{150 MHz}$ with candidates				
orthogonal	3.05	0.68	0.78	0.10
bisector	2.43	0.42	0.76	0.08
Y X	1.98	0.48	0.74	0.07

match is obtained between the two $M_{500}-P_{150 \text{ MHz}}$ scaling relations. For $Y_{500}-P_{150 \text{ MHz}}$ a slightly flatter spectral index seems to be preferred (the blue line in Figure 32 (right panel) is located mostly above our best fit). Giovannini et al. (2009) obtained a medium spectral index of -1.3 for radio halos between 325 MHz and 1.4 GHz, in reasonable agreement our results. However, for a better assessment of the radio halo spectral indices, detailed cluster-to-cluster comparisons are required. In particular, the used scaling relation from Cassano et al. (2013) did not include USSRHs. This could have a direct effect on the derived spectral indices when comparing to a low-frequency sample which potentially contains a significant number of USSRH.

⁴ This also includes Abell 1413 based on new LoTSS data



Fig. 32. Left: Distribution of clusters in the mass (M_{500}) 150 MHz radio power ($P_{150 \text{ MHz}}$) plane. The black solid line displays the BCES orthogonal fit (candidate halos were excluded). The shaded region shows the 3σ (99.7% confidence) region of the fit. The blue line is the BCES orthogonal fit from Cassano et al. (2013) scaled with a spectral index of -1.2. Right: The distribution of clusters in the Y_{500} 150 MHz radio power plane. The plotted symbols and lines represent the same datasets as in the left panel.

6. Conclusions

We have presented a method to extract and re-calibrate targets of interest from the LoTSS survey based on the DDF-pipeline. This method allows joint imaging of data from multiple pointings and an improvement in calibration accuracy. It also enables fast imaging of targets without covering the full FoV of a LO-FAR observation.

We applied the above scheme to 26 Planck PSZ2 and 15 other clusters located in the HETDEX spring region. In total we detect 10 radio halos. Five clusters host radio relics. We also report 12 candidate radio halos. The occurrence fraction of radio halos in PSZ2 clusters is $31\pm11\%$, and $62\pm15\%$ if we include the candidate radio halos. The fraction of clusters with some form of diffuse radio emission in PSZ2 clusters is $73 \pm 17\%$. The relatively large number of candidate radio halos results from the difficulties of unambiguously classifying the emission. Based on the above numbers, we expect to find at least 183 ± 65 radio halos from the analysis of PSZ2 clusters in the LoTSS survey. Considering all clusters and candidate radio halos, we expect that 400–500 halos will be found in the completed LoTSS survey.

We determined for the first time the radio halo scaling relations between cluster mass, Y_{SZ} , and the 150 MHz radio power ($P_{150 \text{ MHz}}$). The slopes for these scaling relations are slightly steeper than those determined at 1.4 GHz, in line with predictions, but considering the uncertainties this is not a statistically significant result. In a future work, using a larger sample, we will present a more detailed statistical analysis of the properties of the diffuse radio sources in the LoTSS survey.

Acknowledgements. This paper is based on data obtained with the International LOFAR Telescope (ILT) under project codes LC2_038 and LC3_008. LOFAR (van Haarlem et al. 2013) is the LOw Frequency ARray designed and constructed by ASTRON. It has observing, data processing, and data storage facilities in several countries, which are owned by various parties (each with their own funding sources) and are collectively operated by the ILT foundation under a joint scientific policy. The ILT resources have benefited from the following recent major funding sources: CNRS-INSU, Observatoire de Paris and Université d'Orléans, France; BMBF, MIWF-NRW, MPG, Germany; Science Foundation Ireland (SFI), Department of Business, Enterprise and Innovation (DBEI), Ireland; NWO, The Netherlands; The Science and Technology Facilities Council, UK; Ministry of Science and Higher Education, Poland; The Istituto Nazionale di Astrofisica (INAF), Italy. This research made use of the Dutch national e-infrastructure with support of the SURF Cooperative (e-infra

180169) and the LOFAR e-infra group. The Jülich LOFAR Long Term Archive and the GermanLOFAR network are both coordinated and operated by the Jülich Supercomputing Centre (JSC), and computing resources on the supercomputer JUWELS at JSC were provided by the Gauss Centre for Supercomputinge.V. (grant CHTB00) through the John von Neumann Institute for Computing (NIC). This research made use of the University of Hertfordshire high-performance computing facility (http://uhhpc.herts.ac.uk) and the LOFAR-UK computing facility located at the University of Hertfordshire and supported by STFC [ST/P000096/1], and of the Italian LOFAR IT computing infrastructure supported and operated by INAF, and by the Physics Department of Turin University (under an agreement with Consorzio Interuniversitario per la Fisica Spaziale) at the C3S Supercomputing Centre, Italy. This research made use of Astropy⁵, a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013, 2018). RJvW and ABott acknowledge support from the VIDI research programme with project number 639.042.729, which is financed by the Netherlands Organisation for Scientific Research (NWO). GB, RC, and FG acknowledge support from INAF mainstream project "Galaxy Clusters Science with LOFAR" 1.05.01.86.05. ABon and EB ackowledge support from the ERC-Stg DRANOEL n. 714245 and from the Italian MIUR grant FARE "SMS". The Pan-STARRS1 Surveys (PS1) and the PS1 public science archive have been made possible through contributions by the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, the Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation Grant No. AST-1238877, the University of Maryland, Eotvos Lorand University (ELTE), the Los Alamos National Laboratory, and the Gordon and Betty Moore Foundation. This research has made use of data obtained from the Chandra Data Archive and the Chandra Source Catalog, and software provided by the Chandra X-ray Center (CXC) in the application packages CIAO, ChIPS, and Sherpa. Based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA.

References

Abell, G. O. 1958, ApJS, 3, 211
Ackermann, M., Ajello, M., Albert, A., et al. 2016, ApJ, 819, 149
Akamatsu, H. & Kawahara, H. 2013, PASJ, 65, 16
Akritas, M. G. & Bershady, M. A. 1996, ApJ, 470, 706
Andrade-Santos, F., Jones, C., Forman, W. R., et al. 2017, ApJ, 843, 76

⁵ http://www.astropy.org

- R. J. van Weeren et al.: LOFAR observations of galaxy clusters in HETDEX
- Arnaud, M., Pratt, G. W., Piffaretti, R., et al. 2010, A&A, 517, A92
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33
- Bagchi, J., Durret, F., Neto, G. B. L., & Paul, S. 2006, Science, 314, 791
- Basu, K. 2012, MNRAS, 421, L112
- Bîrzan, L., Rafferty, D. A., Cassano, R., et al. 2019, MNRAS, 487, 4775
- Blandford, R. & Eichler, D. 1987, Phys. Rep., 154, 1
- Blasi, P. & Colafrancesco, S. 1999, Astroparticle Physics, 12, 169
- Bonafede, A., Brüggen, M., Rafferty, D., et al. 2018, MNRAS, 478, 2927
- Bonafede, A., Cassano, R., Brüggen, M., et al. 2017, MNRAS, 470, 3465
- Bonafede, A., Intema, H. T., Brüggen, M., et al. 2014a, ApJ, 785, 1
- Bonafede, A., Intema, H. T., Bruggen, M., et al. 2014b, MNRAS, 444, L44
- Bonnassieux, E., Tasse, C., Smirnov, O., & Zarka, P. 2018, A&A, 615, A66
- Boschin, W., Girardi, M., & Gastaldello, F. 2020, MNRAS, 492, 2405
- Botteon, A., Brunetti, G., Ryu, D., & Roh, S. 2020a, A&A, 634, A64
- Botteon, A., Brunetti, G., van Weeren, R. J., et al. 2020b, ApJ, 897, 93
- Botteon, A., Cassano, R., Eckert, D., et al. 2019, A&A, 630, A77
- Botteon, A., Gastaldello, F., Brunetti, G., & Dallacasa, D. 2016a, MNRAS, 460, L84
- Botteon, A., Gastaldello, F., Brunetti, G., & Kale, R. 2016b, MNRAS, 463, 1534
- Botteon, A., Shimwell, T. W., Bonafede, A., et al. 2018, MNRAS, 478, 885
- Botteon, A., van Weeren, R. J., Brunetti, G., et al. 2020c, MNRAS, 499, L11
- Boxelaar, J. M., van Weeren, R. J., & Botteon, A. 2020, submitted
- Briggs, D. S. 1995, in American Astronomical Society Meeting Abstracts, Vol. 187, American Astronomical Society Meeting Abstracts, 112.02
- Brüggen, M., Rafferty, D., Bonafede, A., et al. 2018, MNRAS, 477, 3461
- Brüggen, M. & Vazza, F. 2020, MNRAS, 493, 2306
- Brunetti, G., Blasi, P., Reimer, O., et al. 2012, MNRAS, 426, 956
- Brunetti, G., Giacintucci, S., Cassano, R., et al. 2008, Nature, 455, 944
- Brunetti, G. & Jones, T. W. 2014, International Journal of Modern Physics D,
- 23, 30007
- Brunetti, G. & Lazarian, A. 2007, MNRAS, 378, 245
- Brunetti, G. & Lazarian, A. 2011, MNRAS, 410, 127 Brunetti, G., Rudnick, L., Cassano, R., et al. 2013, A&A, 558, A52
- Brunetti, G., Setti, G., Feretti, L., & Giovannini, G. 2001, MNRAS, 320, 365
- Brunetti, G. & Vazza, F. 2020, Phys. Rev. Lett., 124, 051101
- Brunetti, G., Venturi, T., Dallacasa, D., et al. 2007, ApJ, 670, L5
- Brunetti, G., Zimmer, S., & Zandanel, F. 2017, MNRAS, 472, 1506
- Cassano, R. 2010, A&A, 517, A10+
- Cassano, R., Botteon, A., Di Gennaro, G., et al. 2019, ApJ, 881, L18
- Cassano, R. & Brunetti, G. 2005, MNRAS, 357, 1313
- Cassano, R., Brunetti, G., Norris, R. P., et al. 2012, A&A, 548, A100
- Cassano, R., Brunetti, G., Röttgering, H. J. A., & Brüggen, M. 2010a, A&A, 509, A68+
- Cassano, R., Brunetti, G., & Setti, G. 2004, Journal of Korean Astronomical Society, 37, 589
- Cassano, R., Brunetti, G., & Setti, G. 2006, MNRAS, 369, 1577
- Cassano, R., Ettori, S., Brunetti, G., et al. 2013, ApJ, 777, 141
- Cassano, R., Ettori, S., Giacintucci, S., et al. 2010b, ApJ, 721, L82
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv e-prints, arXiv:1612.05560
- Clarke, A. O., Scaife, A. M. M., Shimwell, T., et al. 2019, A&A, 627, A176 Cordey, R. A. 1985, MNRAS, 215, 437
- Corwin, Jr., H. G. 1974, AJ, 79, 1356
- Cuciti, V., Cassano, R., Brunetti, G., et al. 2015, A&A, 580, A97
- David, L. P. & Kempner, J. 2004, ApJ, 613, 831
- de Gasperin, F., Dijkema, T. J., Drabent, A., et al. 2019, A&A, 622, A5
- de Gasperin, F., Intema, H. T., Shimwell, T. W., et al. 2017, Science Advances, 3. e1701634
- de Gasperin, F., Intema, H. T., van Weeren, R. J., et al. 2015, MNRAS, 453, 3483 Dennison, B. 1980, ApJ, 239, L93
- Di Gennaro, G., van Weeren, R. J., Andrade-Santos, F., et al. 2019, ApJ, 873, 64
- Di Gennaro, G., van Weeren, R. J., Brunetti, G., & et al. 2020, Nature Astronomy
- Di Gennaro, G., van Weeren, R. J., Hoeft, M., et al. 2018, ApJ, 865, 24
- Dolag, K. & Enßlin, T. A. 2000, A&A, 362, 151
- Duchesne, S. W., Johnston-Hollitt, M., Offringa, A. R., et al. 2017, arXiv eprints, arXiv:1707.03517
- Eckert, D., Molendi, S., & Paltani, S. 2011, A&A, 526, A79
- Enßlin, T. A., Biermann, P. L., Klein, U., & Kohle, S. 1998, A&A, 332, 395
- Enßlin, T. A. & Brüggen, M. 2002, MNRAS, 331, 1011
- Enßlin, T. A. & Gopal-Krishna. 2001, A&A, 366, 26
- Feretti, L., Giovannini, G., Govoni, F., & Murgia, M. 2012, A&A Rev., 20, 54
- Finoguenov, A., Sarazin, C. L., Nakazawa, K., Wik, D. R., & Clarke, T. E. 2010,
- ApJ, 715, 1143
- Fujita, Y., Kohri, K., Yamazaki, R., & Kino, M. 2007, ApJ, 663, L61
- Fujita, Y. & Ohira, Y. 2013, MNRAS, 428, 599
- Gal, R. R., de Carvalho, R. R., Lopes, P. A. A., et al. 2003, AJ, 125, 2064

- George, L. T., Dwarakanath, K. S., Johnston-Hollitt, M., et al. 2017, MNRAS, 467,936
- Giacintucci, S., Markevitch, M., Cassano, R., et al. 2017, ApJ, 841, 71
- Giacintucci, S., Markevitch, M., Venturi, T., et al. 2014, ApJ, 781, 9 Giacintucci, S., Venturi, T., Macario, G., et al. 2008, A&A, 486, 347
- Giovannini, G., Bonafede, A., Feretti, L., et al. 2009, A&A, 507, 1257 Gitti, M., Brunetti, G., Feretti, L., & Setti, G. 2004, A&A, 417, 1
- Govoni, F., Ferrari, C., Feretti, L., et al. 2012, A&A, 545, A74
- Govoni, F., Orrù, E., Bonafede, A., et al. 2019, Science, 364, 981
- Gu, L., Akamatsu, H., Shimwell, T. W., et al. 2019, Nature Astronomy, 3, 838
- Hao, J., McKay, T. A., Koester, B. P., et al. 2010, ApJS, 191, 254
- Hardcastle, M. J., Croston, J. H., Shimwell, T. W., et al. 2019, MNRAS, 488, 3416
- Hoang, D. N., Shimwell, T. W., van Weeren, R. J., et al. 2019a, A&A, 622, A20
- Hoang, D. N., Shimwell, T. W., van Weeren, R. J., et al. 2019b, A&A, 622, A21 Hoeft, M., Dumba, C., Drabent, A., & et al. 2020, in prep
- Hogg, D. W., Bovy, J., & Lang, D. 2010, arXiv e-prints, arXiv:1008.4686
- Itahana, M., Takizawa, M., Akamatsu, H., et al. 2017, PASJ, 69, 88
- Jones, F. C. & Ellison, D. C. 1991, Space Science Reviews, 58, 259
- Joye, W. A. & Mandel, E. 2003, in Astronomical Society of the Pacific Conference Series, Vol. 295, Astronomical Data Analysis Software and Systems XII, ed. H. E. Payne, R. I. Jedrzejewski, & R. N. Hook, 489
- Kale, R. & Parekh, V. 2016, MNRAS, 459, 2940
- Kale, R., Shende, K. M., & Parekh, V. 2019, MNRAS, 486, L80 Kale, R., Venturi, T., Giacintucci, S., et al. 2015, A&A, 579, A92
- Kale, R., Venturi, T., Giacintucci, S., et al. 2013, A&A, 557, A99
- Kang, H. & Ryu, D. 2011, ApJ, 734, 18
- Kang, H., Ryu, D., & Jones, T. W. 2012, ApJ, 756, 97
- Kempner, J. C. & Sarazin, C. L. 2001, ApJ, 548, 639 Keshet, U. & Loeb, A. 2010, ApJ, 722, 737

Mann, A. W. & Ebeling, H. 2012, MNRAS, 420, 2120

Mazzotta, P. & Giacintucci, S. 2008, ApJ, 675, L9

Offringa, A. R. 2010, AOFlagger: RFI Software

Offringa, A. R. & Smirnov, O. 2017, MNRAS, 471, 301

Ogrean, G. A. & Brüggen, M. 2013, MNRAS, 433, 1701 Paul, S., Salunkhe, S., Sonkamble, S., et al. 2020, A&A, 633, A59

Pinzke, A., Oh, S. P., & Pfrommer, C. 2013, MNRAS, 435, 1061 Pinzke, A., Oh, S. P., & Pfrommer, C. 2017, MNRAS, 465, 4800

Rajpurohit, K., Hoeft, M., Vazza, F., et al. 2020, A&A, 636, A30

Rudnick, L. & Lemmerman, J. A. 2009, ApJ, 697, 1341

Tasse, C. 2014a, arXiv e-prints, arXiv:1410.8706

Rizza, E., Burns, J. O., Ledlow, M. J., et al. 1998, MNRAS, 301, 328

Rossetti, M., Gastaldello, F., Eckert, D., et al. 2017, MNRAS, 468, 1917

Savini, F., Bonafede, A., Brüggen, M., et al. 2019, A&A, 622, A24 Savini, F., Bonafede, A., Brüggen, M., et al. 2018, MNRAS, 478, 2234

Sereno, M., Giocoli, C., Izzo, L., et al. 2018, Nature Astronomy, 2, 744

Shimwell, T. W., Tasse, C., Hardcastle, M. J., et al. 2019, A&A, 622, A1

Srivastava, S. & Singal, A. K. 2016, arXiv e-prints [arXiv:1610.07783]

Shimwell, T. W., Markevitch, M., Brown, S., et al. 2015, MNRAS, 449, 1486 Shimwell, T. W., Röttgering, H. J. A., Best, P. N., et al. 2017, A&A, 598, A104

Slee, O. B., Roy, A. L., Murgia, M., Andernach, H., & Ehle, M. 2001, AJ, 122,

Article number, page 29 of 35

Sebastian, B., Lal, D. V., & Pramesh Rao, A. 2017, AJ, 154, 169

Pfrommer, C. & Enßlin, T. A. 2004, A&A, 413, 17

Offringa, A. R. 2016, A&A, 595, A99

Petrosian, V. 2001, ApJ, 557, 560

A&A, 534, A109

Miniati, F. 2015, ApJ, 800, 60

MNRAS, 420, 2006

606

A95

1172

- Koester, B. P., McKay, T. A., Annis, J., et al. 2007, ApJ, 660, 239 Liang, H., Hunstead, R. W., Birkinshaw, M., & Andreani, P. 2000, ApJ, 544, 686

Mandal, S., Intema, H. T., van Weeren, R. J., et al. 2020, A&A, 634, A4

Markevitch, M., Govoni, F., Brunetti, G., & Jerius, D. 2005, ApJ, 627, 733

Morandi, A., Sun, M., Forman, W., & Jones, C. 2015, MNRAS, 450, 2261

Nemmen, R. S., Georganopoulos, M., Guiriec, S., et al. 2012, Science, 338, 1445

Nuza, S. E., Hoeft, M., van Weeren, R. J., Gottlöber, S., & Yepes, G. 2012,

Offringa, A. R., McKinley, B., Hurley-Walker, N., et al. 2014, MNRAS, 444,

Offringa, A. R., van de Gronde, J. J., & Roerdink, J. B. T. M. 2012, A&A, 539,

Piffaretti, R., Arnaud, M., Pratt, G. W., Pointecouteau, E., & Melin, J.-B. 2011,

Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, A&A, 594, A27 Rajpurohit, K., Hoeft, M., van Weeren, R. J., et al. 2018, ApJ, 852, 65

Richard, J., Pei, L., Limousin, M., Jullo, E., & Kneib, J. P. 2009, A&A, 498, 37

Nuza, S. E., Gelszinnis, J., Hoeft, M., & Yepes, G. 2017, MNRAS, 470, 240

Offringa, A. R., de Bruyn, A. G., Biehl, M., et al. 2010, MNRAS, 405, 155

Murgia, M., Govoni, F., Markevitch, M., et al. 2009, A&A, 499, 679

- Locatelli, N. T., Rajpurohit, K., Vazza, F., et al. 2020, MNRAS, 496, L48 Malkov, M. A. & O'C Drury, L. 2001, Reports on Progress in Physics, 64, 429
- Macario, G., Venturi, T., Intema, H. T., et al. 2013, A&A, 551, A141

Tasse, C. 2014b, A&A, 566, A127

- Tasse, C., Hardcastle, M. J., O'sullivan, S., & et al. 2020, submitted
- Tasse, C., Hugo, B., Mirmont, M., et al. 2018, A&A, 611, A87
- Tikhonov, N. A. & Karachentsev, I. D. 1998, A&AS, 128, 325
- Umetsu, K., Broadhurst, T., Zitrin, A., Medezinski, E., & Hsu, L.-Y. 2011, ApJ, 729, 127
- van der Tol, S., Veenboer, B., & Offringa, A. R. 2018, A&A, 616, A27
- van Diepen, G. & Dijkema, T. J. 2018, DPPP: Default Pre-Processing Pipeline, Astrophysics Source Code Library
- van Haarlem, M. P., Wise, M. W., Gunst, A. W., et al. 2013, A&A, 556, A2
- van Weeren, R. J., Andrade-Santos, F., Dawson, W. A., et al. 2017, Nature Astronomy, 1, 0005
- van Weeren, R. J., Brüggen, M., Röttgering, H. J. A., et al. 2011, A&A, 533, A35+
- van Weeren, R. J., Brunetti, G., Brüggen, M., et al. 2016a, ApJ, 818, 204
- van Weeren, R. J., de Gasperin, F., Akamatsu, H., et al. 2019, Space Sci. Rev., 215, 16
- van Weeren, R. J., Röttgering, H. J. A., Intema, H. T., et al. 2012, A&A, 546, A124
- van Weeren, R. J., Williams, W. L., Hardcastle, M. J., et al. 2016b, ApJS, 223, 2
- Vazza, F. & Brüggen, M. 2014, MNRAS, 437, 2291
- Vazza, F., Brüggen, M., Wittor, D., et al. 2016, MNRAS, 459, 70
- Venturi, T., Giacintucci, S., & Dallacasa, D. 2011, Journal of Astrophysics and Astronomy, 32, 501
- Venturi, T., Giacintucci, S., Dallacasa, D., et al. 2008, A&A, 484, 327
- Venturi, T., Giacintucci, S., Dallacasa, D., et al. 2013, A&A, 551, A24
- Venturi, T., Rossetti, M., Brunetti, G., et al. 2017, A&A, 603, A125
- Wen, Z. L., Han, J. L., & Liu, F. S. 2012, The Astrophysical Journal Supplement Series, 199, 34
- Wilber, A., Brüggen, M., Bonafede, A., et al. 2019, A&A, 622, A25
- Wilber, A., Brüggen, M., Bonafede, A., et al. 2018, MNRAS, 473, 3536
- Yuan, Z. S. & Han, J. L. 2020, MNRAS, 497, 5485
- ZuHone, J. A., Markevitch, M., Brunetti, G., & Giacintucci, S. 2013, ApJ, 762, 78
- Zwicky, F., Herzog, E., Wild, P., Karpowicz, M., & Kowal, C. T. 1961, Catalogue of galaxies and of clusters of galaxies, Vol. I



Fig. A.1. PSZ2 G087.39+50.92. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

Appendix A: Radio images of clusters without diffuse emission

Appendix A.1: PSZ2 G087.39+50.92

The LOFAR image show a compact AGN associated with central BCG of this distant cluster (Figure A.1). No diffuse radio emission is detected. We note that one of the LoTSS observations had to be discarded due to bad ionospheric conditions.

Appendix A.2: PSZ2 G088.98+55.07

A compact radio source is detected at the center of this distant z = 0.7023 cluster. An optical image with radio contours overlaid is shown in Figure A.2. We do not find evidence for diffuse emission in this cluster. However, we note that the LOFAR observations were affected by bad ionospheric conditions.

Appendix A.3: PSZ2 G096.14+56.24, Abell 1940

No diffuse radio emission is detected Abell 1940. The image dynamic range is limited by the bright double lobed radio source associated with the BCG. An optical image with radio contours overlaid is shown in Figure A.3.

Appendix A.4: PSZ2 G098.44+56.59, Abell 1920

Several tailed radio galaxies are visible in the cluster region, but no diffuse emission is detected (see Figure A.4).

Appendix A.5: PSZ2 G123.66+67.25, Abell 1622

The Chandra X-ray image shows that the cluster consists of two subclusters, see Figure A.5. The main subcluster has a large X-ray extent (~ 1 Mpc) and low surface brightness. A smaller more concentrated subcluster is located south of it. Given the overall roundish morphology of both components, the cluster is likely in a pre-merging state. A additional small substructure is seen to the SW of the southern subcluster. PSZ2 G123.66+67.25 has a



Fig. A.2. PSZ2 G088.98+55.07. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.3. PSZ2 G096.14+56.24 / Abell 1940. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

temperature of 4.76 ± 0.87 keV and is classified as a non-cool core by Morandi et al. (2015).

No Mpc-scale diffuse emission is detected in this z = 0.2838 cluster. A bright tailed radio galaxy (A) is found in the northern part of the main subcluster, with the tail bending east about 200 kpc north of its optical counterpart. A patch of radio plasma (B) with an LLS of 250 kpc is found to the south of the northern subcluster. A compact radio source is located just north of this diffuse source, but there is no clear connection visible in our high-resolution images. We speculate that this the diffuse source is AGN fossil plasma that originated from the compact radio source.





Appendix A.6: PSZ2 G136.92+59.46, Abell 1436

The LOFAR image displays compact radio emission associated with the BCG. The XMM-Newton image reveals an elongated low surface brightness cluster, see Figure A.6. Optical Pan-STARRS images show the BCG to have a double nucleus. Additional radio emission is detected south of the BCG. Given that this emission connects to the BCG it likely originated from the BCG. No diffuse radio emission is detected in this cluster.

Appendix A.7: PSZ2 G144.33+62.85, Abell 1387

No diffuse radio emission is detected in this cluster. Two BCGs seem to be present in this cluster based on optical Pan-STARRS images. A distorted tailed radio galaxy is associated with one of the two BCGs.

Appendix A.8: PSZ2 G151.62+54.78

No diffuse emission is detected in this cluster, see Figure A.8. However, the observations for this cluster were affected by bad ionospheric conditions, compromising the image quality.

Appendix A.9: Abell 1615

This cluster host a complex distorted tailed radio galaxy related to the BCG. This source has a largest extended of 450 kpc. An optical image with radio contours overlaid is shown in Figure A.9.

Appendix A.10: GMBCG J211.77332+55.09968

This cluster hosts large 600 kpc asymmetric radio galaxy associated with the BCG. Extended radio emission is also found NW of the BCG, see Figure A.10. This emission is composed of three distinct radio sources, a compact double lobed source, an extended double lobed source, and a foreground spiral galaxy.

Appendix A.11: MaxBCG J173.04772+47.81041

Extended radio emission, likely related to AGN activity from an elliptical galaxy, is detected in the northern part of the cluster. An optical image with radio contours overlaid is shown in Figure A.11.

Appendix A.12: WHL J132615.8+485229

Elongated radio emission is detected that seems to originate from a tailed radio galaxy south of the cluster center (see Figure A.12). This emission extends all the way to the BCG.

Appendix A.13: WHL J134746.8+475214

This cluster host complex extended radio emission on scales of 500 kpc which seems to be related to AGN activity. An optical image with radio contours overlaid is shown in Figure A.13.



Fig. A.5. PSZ2 G123.66+67.25 / Abell 1622. Left: Robust -0.5 radio image. Middle: Chandra X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.6. PSZ2 G136.92+59.46 / Abell 1436. Left: XMM-Newton X-ray image with 10" tapered radio contours. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.





Fig. A.7. PSZ2 G144.33+62.85, Abell 1387. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.

Fig. A.8. PSZ2 G151.62+54.78. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.9. Abell 1615. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.11. MaxBCG J173.04772+47.81041. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.10. GMBCG J211.77332+55.09968. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.12. WHL J132615.8+485229. Left: Robust -0.5 image radio image. Right: Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.



Fig. A.13. WHL J134746.8+475214. Optical image with Robust -0.5 image radio contours. For more details see the caption of Figure 5.